

97

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7 : C12N 15/56, 15/63, 1/21, 9/24, 15/11		A1	(11) International Publication Number: WO 00/52178
			(43) International Publication Date: 8 September 2000 (08.09.00)
(21) International Application Number: PCT/US00/03542		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 14 February 2000 (14.02.00)		<p>Published</p> <p><i>With international search report.</i></p> <p><i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	
(30) Priority Data: 09/258,892 1 March 1999 (01.03.99) US			
(71) Applicants (for all designated States except US): INSIGHT STRATEGY & MARKETING LTD. [IL/IL]; P.O. Box 2128, 76121 Rehovot (IL). HADASIT MEDICAL RESEARCH SERVICES & DEVELOPMENT LTD. [IL/IL]; P.O. Box 12000, 91120 Jerusalem (IL).			
(71) Applicant (for TJ only): FRIEDMAN, Mark, M. [US/IL]; 1 Alharizi Street, 43406 Raanana (IL).			
(72) Inventors; and (75) Inventors/Applicants (for US only): PECKER, Iris [IL/IL]; 42 Wolfson Street, 75203 Rishon Lezion (IL). VLODAVSKY, Israel [IL/IL]; 34 Arbel Street, 90805 Mevaseret Zion (IL). FEINSTEIN, Elena [IL/IL]; 12/29 Hahagana Street, 76214 Rehovot (IL).			
(74) Common Representative: FRIEDMAN, Mark, M.; c/o Castorina, Anthony, Suite 207, 2001 Jefferson Davis Highway, Arlington, VA 22202 (US).			
(54) Title: POLYNUCLEOTIDE ENCODING A POLYPEPTIDE HAVING HEPARANASE ACTIVITY AND EXPRESSION OF SAME IN GENETICALLY MODIFIED CELLS			
(57) Abstract			
<p>A polynucleotide (<i>hpa</i>) encoding a polypeptide having heparanase activity, vectors including same, genetically modified cells expressing heparanase, a recombinant protein having heparanase activity and antisense oligonucleotides and constructs for modulating heparanase expression.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

POLYNUCLEOTIDE ENCODING A POLYPEPTIDE HAVING
HEPARANASE ACTIVITY AND EXPRESSION OF SAME IN
GENETICALLY MODIFIED CELLS

5 FIELD AND BACKGROUND OF THE INVENTION

The present invention relates to a polynucleotide, referred to hereinbelow as *hpa*, encoding a polypeptide having heparanase activity, vectors (nucleic acid constructs) including same and genetically modified cells expressing heparanase. The invention further relates to a recombinant
10 protein having heparanase activity and to antisense oligonucleotides, constructs and ribozymes for down regulating heparanase activity. In addition, the invention relates to heparanase promoter sequences and their uses.

Heparan sulfate proteoglycans: Heparan sulfate proteoglycans
15 (HSPG) are ubiquitous macromolecules associated with the cell surface and extra cellular matrix (ECM) of a wide range of cells of vertebrate and invertebrate tissues (1-4). The basic HSPG structure includes a protein core to which several linear heparan sulfate chains are covalently attached. These polysaccharide chains are typically composed of repeating hexuronic
20 and D-glucosamine disaccharide units that are substituted to a varying extent with N- and O-linked sulfate moieties and N-linked acetyl groups (1-4). Studies on the involvement of ECM molecules in cell attachment, growth and differentiation revealed a central role of HSPG in embryonic morphogenesis, angiogenesis, neurite outgrowth and tissue repair (1-5).
25 HSPG are prominent components of blood vessels (3). In large blood vessels they are concentrated mostly in the intima and inner media, whereas in capillaries they are found mainly in the subendothelial basement membrane where they support proliferating and migrating endothelial cells and stabilize the structure of the capillary wall. The ability of HSPG to
30 interact with ECM macromolecules such as collagen, laminin and fibronectin, and with different attachment sites on plasma membranes suggests a key role for this proteoglycan in the self-assembly and insolubility of ECM components, as well as in cell adhesion and locomotion. Cleavage of the heparan sulfate (HS) chains may therefore
35 result in degradation of the subendothelial ECM and hence may play a decisive role in extravasation of blood-borne cells. HS catabolism is observed in inflammation, wound repair, diabetes, and cancer metastasis, suggesting that enzymes which degrade HS play important roles in pathologic processes. Heparanase activity has been described in activated

immune system cells and highly metastatic cancer cells (6-8), but research has been handicapped by the lack of biologic tools to explore potential causative roles of heparanase in disease conditions.

Involvement of Heparanase in Tumor Cell Invasion and

5 ***Metastasis:*** Circulating tumor cells arrested in the capillary beds of different organs must invade the endothelial cell lining and degrade its underlying basement membrane (BM) in order to invade into the extravascular tissue(s) where they establish metastasis (9, 10). Metastatic
10 tumor cells often attach at or near the intercellular junctions between adjacent endothelial cells. Such attachment of the metastatic cells is followed by rupture of the junctions, retraction of the endothelial cell borders and migration through the breach in the endothelium toward the exposed underlying BM (9). Once located between endothelial cells and the BM, the invading cells must degrade the subendothelial glycoproteins and
15 proteoglycans of the BM in order to migrate out of the vascular compartment. Several cellular enzymes (e.g., collagenase IV, plasminogen activator, cathepsin B, elastase, etc.) are thought to be involved in degradation of BM (10). Among these enzymes is an endo- β -D-glucuronidase (heparanase) that cleaves HS at specific intrachain sites (6,
20 8, 11). Expression of a HS degrading heparanase was found to correlate with the metastatic potential of mouse lymphoma (11), fibrosarcoma and melanoma (8) cells. Moreover, elevated levels of heparanase were detected in sera from metastatic tumor bearing animals and melanoma patients (8) and in tumor biopsies of cancer patients (12).

25 The control of cell proliferation and tumor progression by the local microenvironment, focusing on the interaction of cells with the extracellular matrix (ECM) produced by cultured corneal and vascular endothelial cells, was investigated previously by the present inventors. This cultured ECM closely resembles the subendothelium *in vivo* in its
30 morphological appearance and molecular composition. It contains collagens (mostly type III and IV, with smaller amounts of types I and V), proteoglycans (mostly heparan sulfate- and dermatan sulfate- proteoglycans, with smaller amounts of chondroitin sulfate proteoglycans), laminin, fibronectin, entactin and elastin (13, 14). The ability of cells to degrade HS
35 in the cultured ECM was studied by allowing cells to interact with a metabolically sulfate labeled ECM, followed by gel filtration (Sephacrose 6B) analysis of degradation products released into the culture medium (11). While intact HSPG are eluted next to the void volume of the column

($K_{av} < 0.2$, $M_r \sim 0.5 \times 10^6$), labeled degradation fragments of HS side chains are eluted more toward the V_t of the column ($0.5 < k_{av} < 0.8$, $M_r = 5-7 \times 10^3$) (11).

The heparanase inhibitory effect of various non-anticoagulant species of heparin that might be of potential use in preventing extravasation of blood-borne cells was also investigated by the present inventors. Inhibition of heparanase was best achieved by heparin species containing 16 sugar units or more and having sulfate groups at both the N and O positions. While O-desulfation abolished the heparanase inhibiting effect of heparin, O-sulfated, N-acetylated heparin retained a high inhibitory activity, provided that the N-substituted molecules had a molecular size of about 4,000 daltons or more (7). Treatment of experimental animals with heparanase inhibitors (e.g., non-anticoagulant species of heparin) markedly reduced (>90%) the incidence of lung metastases induced by B16 melanoma, Lewis lung carcinoma and mammary adenocarcinoma cells (7, 8, 16). Heparin fractions with high and low affinity to anti-thrombin III exhibited a comparable high anti-metastatic activity, indicating that the heparanase inhibiting activity of heparin, rather than its anticoagulant activity, plays a role in the anti-metastatic properties of the polysaccharide (7).

Heparanase activity in the urine of cancer patients: In an attempt to further elucidate the involvement of heparanase in tumor progression and its relevance to human cancer, urine samples for heparanase activity were screened (16a). Heparanase activity was detected in the urine of some, but not all, cancer patients. High levels of heparanase activity were determined in the urine of patients with an aggressive metastatic disease and there was no detectable activity in the urine of healthy donors.

Heparanase activity was also found in the urine of 20% of normal and microalbuminuric insulin dependent diabetes mellitus (IDDM) patients, most likely due to diabetic nephropathy, the most important single disorder leading to renal failure in adults.

Possible involvement of heparanase in tumor angiogenesis: Fibroblast growth factors are a family of structurally related polypeptides characterized by high affinity to heparin (17). They are highly mitogenic for vascular endothelial cells and are among the most potent inducers of neovascularization (17, 18). Basic fibroblast growth factor (bFGF) has been extracted from the subendothelial ECM produced *in vitro* (19) and from basement membranes of the cornea (20), suggesting that ECM may serve as

a reservoir for bFGF. Immunohistochemical staining revealed the localization of bFGF in basement membranes of diverse tissues and blood vessels (21). Despite the ubiquitous presence of bFGF in normal tissues, endothelial cell proliferation in these tissues is usually very low, suggesting that bFGF is somehow sequestered from its site of action. Studies on the interaction of bFGF with ECM revealed that bFGF binds to HSPG in the ECM and can be released in an active form by HS degrading enzymes (15, 20, 22). It was demonstrated that heparanase activity expressed by platelets, mast cells, neutrophils, and lymphoma cells is involved in release of active bFGF from ECM and basement membranes (23), suggesting that heparanase activity may not only function in cell migration and invasion, but may also elicit an indirect neovascular response. These results suggest that the ECM HSPG provides a natural storage depot for bFGF and possibly other heparin-binding growth promoting factors (24, 25). Displacement of bFGF from its storage within basement membranes and ECM may therefore provide a novel mechanism for induction of neovascularization in normal and pathological situations.

Recent studies indicate that heparin and HS are involved in binding of bFGF to high affinity cell surface receptors and in bFGF cell signaling (26, 27). Moreover, the size of HS required for optimal effect was similar to that of HS fragments released by heparanase (28). Similar results were obtained with vascular endothelial cells growth factor (VEGF) (29), suggesting the operation of a dual receptor mechanism involving HS in cell interaction with heparin-binding growth factors. It is therefore proposed that restriction of endothelial cell growth factors in ECM prevents their systemic action on the vascular endothelium, thus maintaining a very low rate of endothelial cells turnover and vessel growth. On the other hand, release of bFGF from storage in ECM as a complex with HS fragment, may elicit localized endothelial cell proliferation and neovascularization in processes such as wound healing, inflammation and tumor development (24, 25).

Expression of heparanase by cells of the immune system:

Heparanase activity correlates with the ability of activated cells of the immune system to leave the circulation and elicit both inflammatory and autoimmune responses. Interaction of platelets, granulocytes, T and B lymphocytes, macrophages and mast cells with the subendothelial ECM is associated with degradation of HS by a specific heparanase activity (6). The enzyme is released from intracellular compartments (e.g., lysosomes,

specific granules, etc.) in response to various activation signals (e.g., thrombin, calcium ionophore, immune complexes, antigens, mitogens, etc.), suggesting its regulated involvement in inflammation and cellular immunity.

5 ***Some of the observations regarding the heparanase enzyme were reviewed in reference No. 6 and are listed hereinbelow:***

First, a proteolytic activity (plasminogen activator) and heparanase participate synergistically in sequential degradation of the ECM HSPG by inflammatory leukocytes and malignant cells.

10 Second, a large proportion of the platelet heparanase exists in a latent form, probably as a complex with chondroitin sulfate. The latent enzyme is activated by tumor cell-derived factor(s) and may then facilitate cell invasion through the vascular endothelium in the process of tumor metastasis.

15 Third, release of the platelet heparanase from α -granules is induced by a strong stimulant (i.e., thrombin), but not in response to platelet activation on ECM.

20 Fourth, the neutrophil heparanase is preferentially and readily released in response to a threshold activation and upon incubation of the cells on ECM.

25 Fifth, contact of neutrophils with ECM inhibited release of noxious enzymes (proteases, lysozyme) and oxygen radicals, but not of enzymes (heparanase, gelatinase) which may enable diapedesis. This protective role of the subendothelial ECM was observed when the cells were stimulated with soluble factors but not with phagocytosable stimulants.

Sixth, intracellular heparanase is secreted within minutes after exposure of T cell lines to specific antigens.

30 Seventh, mitogens (Con A, LPS) induce synthesis and secretion of heparanase by normal T and B lymphocytes maintained *in vitro*. T lymphocyte heparanase is also induced by immunization with antigen *in vivo*.

Eighth, heparanase activity is expressed by pre-B lymphomas and B-lymphomas, but not by plasmacytomas and resting normal B lymphocytes.

35 Ninth, heparanase activity is expressed by activated macrophages during incubation with ECM, but there was little or no release of the enzyme into the incubation medium. Similar results were obtained with human myeloid leukemia cells induced to differentiate to mature macrophages.

Tenth, T-cell mediated delayed type hypersensitivity and experimental autoimmunity are suppressed by low doses of heparanase inhibiting non-anticoagulant species of heparin (30).

Eleventh, heparanase activity expressed by platelets, neutrophils and metastatic tumor cells releases active bFGF from ECM and basement membranes. Release of bFGF from storage in ECM may elicit a localized neovascular response in processes such as wound healing, inflammation and tumor development.

Twelfth, among the breakdown products of the ECM generated by heparanase is a tri-sulfated disaccharide that can inhibit T-cell mediated inflammation *in vivo* (31). This inhibition was associated with an inhibitory effect of the disaccharide on the production of biologically active TNF α by activated T cells *in vitro* (31).

Other potential therapeutic applications: Apart from its involvement in tumor cell metastasis, inflammation and autoimmunity, mammalian heparanase may be applied to modulate: bioavailability of heparin-binding growth factors (15); cellular responses to heparin-binding growth factors (e.g., bFGF, VEGF) and cytokines (IL-8) (31a, 29); cell interaction with plasma lipoproteins (32); cellular susceptibility to certain viral and some bacterial and protozoa infections (33, 33a, 33b); and disintegration of amyloid plaques (34). Heparanase may thus prove useful for conditions such as wound healing, angiogenesis, restenosis, atherosclerosis, inflammation, neurodegenerative diseases and viral infections. Mammalian heparanase can be used to neutralize plasma heparin, as a potential replacement of protamine. Anti-heparanase antibodies may be applied for immunodetection and diagnosis of micrometastases, autoimmune lesions and renal failure in biopsy specimens, plasma samples, and body fluids. Common use in basic research is expected.

The identification of the *hpa* gene encoding for heparanase enzyme will enable the production of a recombinant enzyme in heterologous expression systems. Availability of the recombinant protein will pave the way for solving the protein structure function relationship and will provide a tool for developing new inhibitors.

Viral Infection: The presence of heparan sulfate on cell surfaces have been shown to be the principal requirement for the binding of Herpes Simplex (33) and Dengue (33a) viruses to cells and for subsequent infection of the cells. Removal of the cell surface heparan sulfate by heparanase may

therefore abolish virus infection. In fact, treatment of cells with bacterial heparitinase (degrading heparan sulfate) or heparinase (degrading heparan) reduced the binding of two related animal herpes viruses to cells and rendered the cells at least partially resistant to virus infection (33). There are some indications that the cell surface heparan sulfate is also involved in HIV infection (33b).

Neurodegenerative diseases: Heparan sulfate proteoglycans were identified in the prion protein amyloid plaques of Genstmann-Straussler Syndrome, Creutzfeldt-Jakob disease and Scrape (34). Heparanase may disintegrate these amyloid plaques which are also thought to play a role in the pathogenesis of Alzheimer's disease.

Restenosis and Atherosclerosis: Proliferation of arterial smooth muscle cells (SMCs) in response to endothelial injury and accumulation of cholesterol rich lipoproteins are basic events in the pathogenesis of atherosclerosis and restenosis (35). Apart from its involvement in SMC proliferation (i.e., low affinity receptors for heparin-binding growth factors), HS is also involved in lipoprotein binding, retention and uptake (36). It was demonstrated that HSPG and lipoprotein lipase participate in a novel catabolic pathway that may allow substantial cellular and interstitial accumulation of cholesterol rich lipoproteins (32). The latter pathway is expected to be highly atherogenic by promoting accumulation of apoB and apoE rich lipoproteins (i.e. LDL, VLDL, chylomicrons), independent of feed back inhibition by the cellular sterol content. Removal of SMC HS by heparanase is therefore expected to inhibit both SMC proliferation and lipid accumulation and thus may halt the progression of restenosis and atherosclerosis.

Gene therapy:

The ultimate goal in the management of inherited as well as acquired diseases is a rational therapy with the aim to eliminate the underlying biochemical defects associated with the disease rather than symptomatic treatment. Gene therapy is a promising candidate to meet these objectives. Initially it was developed for treatment of genetic disorders, however, the consensus view today is that it offers the prospect of providing therapy for a variety of acquired diseases, including cancer, viral infections, vascular diseases and neurodegenerative disorders.

The gene-based therapeutic can act either intracellularly, affecting only the cells to which it is delivered, or extracellularly, using the recipient cells as local endogenous factories for the therapeutic product(s). The

application of gene therapy may follow any of the following strategies: (i) prophylactic gene therapy, such as using gene transfer to protect cells against viral infection; (ii) cytotoxic gene therapy, such as cancer therapy, where genes encode cytotoxic products to render the target cells vulnerable to attack by the normal immune response; (iii) biochemical correction, primarily for the treatment of single gene defects, where a normal copy of the gene is added to the affected or other cells.

To allow efficient transfer of the therapeutic genes, a variety of gene delivery techniques have been developed based on viral and non-viral vector systems. The most widely used and most efficient systems for delivering genetic material into target cells are viral vectors. So far, 329 clinical studies (phase I, I/II and II) with over 2,500 patients have been initiated Worldwide since 1989 (50).

The approach of gene addition pose serious barriers. The expression of many genes is tightly regulated and context dependent, so achieving the correct balance and function of expression is challenging. The gene itself is often quite large, containing many exons and introns. The delivery vector is usually a virus, which can infect with a high efficiency but may, on the other hand, induce immunological response and consequently decreases effectiveness, especially upon secondary administration. Most of the current expression vector-based gene therapy protocols fail to achieve clinically significant transgene expression required for treating genetic diseases. Apparently, it is difficult to deliver enough virus to the right cell type to elicit an effective and therapeutic effect (51)

Homologous recombination, which was initially considered to be of limited use for gene therapy because of its low frequency in mammalian cells, has recently emerged as a potential strategy for developing gene therapy. Different approaches have been used to study homologous recombination in mammalian cells; some involve DNA repair mechanisms. These studies aimed at either gene disruption or gene correction and include RNA/DNA chimeric oligonucleotides, small or large homologous DNA fragments, or adeno-associated viral vectors. Most of these studies show a reasonable frequency of homologous recombination, which warrants further *in vivo* testing (52). Homologous recombination-based gene therapy has the potential to develop into a powerful therapeutic modality for genetic diseases. It can offer permanent expression and normal regulation of corrected genes in appropriate cells or organs and probably can be used for treating dominantly inherited diseases such as polycystic kidney disease.

Genomic sequences function in regulation of gene expression:

The efficient expression of therapeutic genes in target cells or tissues is an important component of efficient and safe gene therapy. The expression of genes is driven by the promoter region upstream of the coding sequence, although regulation of expression may be supplemented by farther upstream or downstream DNA sequences or DNA in the introns of the gene. Since this important information is embedded in the DNA, the description of gene structure is crucial to the analysis of gene regulation. Characterization of cell specific or tissue specific promoters, as well as other tissue specific regulatory elements enables the use of such sequences to direct efficient cell specific, or developmental stage specific gene expression. This information provides the basis for targeting individual genes and for control of their expression by exogenous agents, such as drugs. Identification of transcription factors and other regulatory proteins required for proper gene expression will point at new potential targets for modulating gene expression, when so desired or required.

Efficient expression of many mammalian genes depends on the presence of at least one intron. The expression of mouse thymidylate synthase (TS) gene, for example, is greatly influenced by intron sequences. The addition of almost any of the introns from the mouse TS gene to an intronless TS minigene leads to a large increase in expression (42). The involvement of intron 1 in the regulation of expression was demonstrated for many other genes. In human factor IX (hFIX), intron 1 is able to increase the expression level about 3 fold more as compared to that of the hFIX cDNA (43). The expression enhancing activity of intron 1 is due to efficient functional splicing sequences, present in the precursor mRNA. By being efficiently assembled into spliceosome complexes, transcripts with splicing sequences may be better protected in the nucleus from random degradations, than those without such sequences (44).

A forward-inserted intron1-carrying hFIX expression cassette suggested to be useful for directed gene transfer, while for retroviral-mediated gene transfer system, reversely-inserted intron 1-carrying hFIX expression cassette was considered (43).

A highly conserved cis-acting sequence element was identified in the first intron of the mouse and rat c-Ha-ras, and in the first exon of Ha- and Ki-ras genes of human, mouse and rat. This cis-acting regulatory sequence confers strong transcription enhancer activity that is differentially modulated by steroid hormones in metastatic and nonmetastatic

subpopulations. Perturbations in the regulatory activities of such cis-acting sequences may play an important role in governing oncogenic potency of Ha-ras through transcriptional control mechanisms (45).

5 Intron sequences affect tissue specific, as well as inducible gene expression. A 182 bp intron 1 DNA segment of the mouse Col2a1 gene contains the necessary information to confer high-level, temporally correct, chondrocyte expression on a reporter gene in intact mouse embryos, while Col2a1 promoter sequences are dispensable for chondrocyte expression (46). In CollA1 gene the intron plays little or no role in constitutive
10 expression of collagen in the skin, and in cultured cells derived from the skin, however, in the lungs of young mice, intron deletion results in decrease of expression to less than 50 % (47).

A classical enhancer activity was shown in the 2 kb intron fragment in bovine beta-casein gene. The enhancer activity was largely dependent on
15 the lactogenic hormones, especially prolactin. It was suggested that several elements in the intron-1 of the bovine beta-casein gene cooperatively interact not only with each other but also with its promoter for hormonal induction (48).

Identification and characterization of regulatory elements in genomic
20 non-coding sequences, such as introns, provides a tool for designing and constructing novel vectors for tissue specific, hormone regulated or any other defined expression pattern, for gene therapy. Such an expression cassette was developed, utilizing regulatory elements from the human cytokeratin 18 (K18) gene, including 5' genomic sequences and one of its
25 introns. This cassette efficiently expresses reporter genes, as well as the human cystic fibrosis transmembrane conductance regulator (CFTR) gene, in cultured lung epithelial cells (49).

Alternative splicing:

Alternative splicing of pre mRNA is a powerful and versatile
30 regulatory mechanism that can effect quantitative control of gene expression and functional diversification of proteins. It contributes to major developmental decisions and also to a fine-tuning of gene function. Genetic and biochemical approaches have identified cis-acting regulatory elements and trans-acting factors that control alternative splicing of specific mRNAs.
35 This mechanism results in the generation of variant isoforms of various proteins from a single gene. These include cell surface molecules such as CD44, receptors, cytokines such as VEGF and enzymes. Products of

alternatively spliced transcripts differ in their expression pattern, substrate specificity and other biological parameters.

The FGF receptor RNA undergoes alternative splicing which results in the production of several isoforms, which exhibit different ligand binding specificities. The alternative splicing is regulated in a cell specific manner (53).

Alternative spliced mRNAs are often correlated with malignancy. An increase in specific splice variant of tyrosinase was identified in murine melanomas (54). Multiple splicing variants of estrogen receptor are present in individual human breast tumors. CD44 has various isoform, some are characteristic of malignant tissues.

Identification of tumor specific alternative splice variants provide new tool for cancer diagnostics. CD44 variants have been used for detection of malignancy in urine samples from patients with urothelial cancer by competitive RT-PCR (55). CD44 exon 6 was suggested as prognostic indicator of metastasis in breast cancer (56).

Different enzymes or polypeptides generated by alternative splicing may have different function or catalytic specificity. The identification and characterization of the enzyme forms, which are involved in pathological processes, is crucial for the design of appropriate and efficient drugs.

Modulation of gene expression – Antisense technology:

An antisense oligonucleotide (e.g., antisense oligodeoxyribonucleotide) may bind its target nucleic acid either by Watson-Crick base pairing or Hoogsteen and anti-Hoogsteen base pairing (64). According to the Watson-Crick base pairing, heterocyclic bases of the antisense oligonucleotide form hydrogen bonds with the heterocyclic bases of target single-stranded nucleic acids (RNA or single-stranded DNA), whereas according to the Hoogsteen base pairing, the heterocyclic bases of the target nucleic acid are double-stranded DNA, wherein a third strand is accommodated in the major groove of the B-form DNA duplex by Hoogsteen and anti-Hoogsteen base pairing to form a triple helix structure.

According to both the Watson-Crick and the Hoogsteen base pairing models, antisense oligonucleotides have the potential to regulate gene expression and to disrupt the essential functions of the nucleic acids in cells. Therefore, antisense oligonucleotides have possible uses in modulating a wide range of diseases in which gene expression is altered.

Since the development of effective methods for chemically synthesizing oligonucleotides, these molecules have been extensively used

in biochemistry and biological research and have the potential use in medicine, since carefully devised oligonucleotides can be used to control gene expression by regulating levels of transcription, transcripts and/or translation.

5 Oligodeoxyribonucleotides as long as 100 base pairs (bp) are routinely synthesized by solid phase methods using commercially available, fully automated synthesis machines. The chemical synthesis of oligoribonucleotides, however, is far less routine. Oligoribonucleotides are also much less stable than oligodeoxyribonucleotides, a fact which has
10 contributed to the more prevalent use of oligodeoxyribonucleotides in medical and biological research, directed at, for example, the regulation of transcription or translation levels.

Gene expression involves few distinct and well regulated steps. The first major step of gene expression involves transcription of a messenger
15 RNA (mRNA) which is an RNA sequence complementary to the antisense (i.e., -) DNA strand, or, in other words, identical in sequence to the DNA sense (i.e., +) strand, composing the gene. In eukaryotes, transcription occurs in the cell nucleus.

The second major step of gene expression involves translation of a
20 protein (e.g., enzymes, structural proteins, secreted proteins, gene expression factors, etc.) in which the mRNA interacts with ribosomal RNA complexes (ribosomes) and amino acid activated transfer RNAs (tRNAs) to direct the synthesis of the protein coded for by the mRNA sequence.

Initiation of transcription requires specific recognition of a promoter
25 DNA sequence located upstream to the coding sequence of a gene by an RNA-synthesizing enzyme -- RNA polymerase. This recognition is preceded by sequence-specific binding of one or more transcription factors to the promoter sequence. Additional proteins which bind at or close to the promoter sequence may trans upregulate transcription via cis elements
30 known as enhancer sequences. Other proteins which bind to or close to the promoter, but whose binding prohibits the action of RNA polymerase, are known as repressors.

There are also evidence that in some cases gene expression is downregulated by endogenous antisense RNA repressors that bind a
35 complementary mRNA transcript and thereby prevent its translation into a functional protein.

Thus, gene expression is typically upregulated by transcription factors and enhancers and downregulated by repressors.

However, in many disease situation gene expression is impaired. In many cases, such as different types of cancer, for various reasons the expression of a specific endogenous or exogenous (e.g., of a pathogen such as a virus) gene is upregulated. Furthermore, in infectious diseases caused by pathogens such as parasites, bacteria or viruses, the disease progression depends on expression of the pathogen genes, this phenomenon may also be considered as far as the patient is concerned as upregulation of exogenous genes.

Most conventional drugs function by interaction with and modulation of one or more targeted endogenous or exogenous proteins, e.g., enzymes. Such drugs, however, typically are not specific for targeted proteins but interact with other proteins as well. Thus, a relatively large dose of drug must be used to effectively modulate a targeted protein.

Typical daily doses of drugs are from 10^{-5} - 10^{-1} millimoles per kilogram of body weight or 10^{-3} - 10 millimoles for a 100 kilogram person. If this modulation instead could be effected by interaction with and inactivation of mRNA, a dramatic reduction in the necessary amount of drug could likely be achieved, along with a corresponding reduction in side effects. Further reductions could be effected if such interaction could be rendered site-specific. Given that a functioning gene continually produces mRNA, it would thus be even more advantageous if gene transcription could be arrested in its entirety.

Given these facts, it would be advantageous if gene expression could be arrested or downmodulated at the transcription level.

The ability of chemically synthesizing oligonucleotides and analogs thereof having a selected predetermined sequence offers means for downmodulating gene expression. Three types of gene expression modulation strategies may be considered.

At the transcription level, antisense or sense oligonucleotides or analogs that bind to the genomic DNA by strand displacement or the formation of a triple helix, may prevent transcription (64).

At the transcript level, antisense oligonucleotides or analogs that bind target mRNA molecules lead to the enzymatic cleavage of the hybrid by intracellular RNase H (65). In this case, by hybridizing to the targeted mRNA, the oligonucleotides or oligonucleotide analogs provide a duplex hybrid recognized and destroyed by the RNase H enzyme. Alternatively, such hybrid formation may lead to interference with correct splicing (66).

As a result, in both cases, the number of the target mRNA intact transcripts ready for translation is reduced or eliminated.

At the translation level, antisense oligonucleotides or analogs that bind target mRNA molecules prevent, by steric hindrance, binding of essential translation factors (ribosomes), to the target mRNA, a phenomenon known in the art as hybridization arrest, disabling the translation of such mRNAs (67).

Thus, antisense sequences, which as described hereinabove may arrest the expression of any endogenous and/or exogenous gene depending on their specific sequence, attracted much attention by scientists and pharmacologists who were devoted at developing the antisense approach into a new pharmacological tool (68).

For example, several antisense oligonucleotides have been shown to arrest hematopoietic cell proliferation (69), growth (70), entry into the S phase of the cell cycle (71), reduced survival (72) and prevent receptor mediated responses (73). For use of antisense oligonucleotides as antiviral agents the reader is referred to reference 74.

For efficient *in vivo* inhibition of gene expression using antisense oligonucleotides or analogs, the oligonucleotides or analogs must fulfill the following requirements (i) sufficient specificity in binding to the target sequence; (ii) solubility in water; (iii) stability against intra- and extracellular nucleases; (iv) capability of penetration through the cell membrane; and (v) when used to treat an organism, low toxicity.

Unmodified oligonucleotides are impractical for use as antisense sequences since they have short *in vivo* half-lives, during which they are degraded rapidly by nucleases. Furthermore, they are difficult to prepare in more than milligram quantities. In addition, such oligonucleotides are poor cell membrane penetrators (75).

Thus it is apparent that in order to meet all the above listed requirements, oligonucleotide analogs need to be devised in a suitable manner. Therefore, an extensive search for modified oligonucleotides has been initiated.

For example, problems arising in connection with double-stranded DNA (dsDNA) recognition through triple helix formation have been diminished by a clever "switch back" chemical linking, whereby a sequence of polypurine on one strand is recognized, and by "switching back", a homopurine sequence on the other strand can be recognized. Also, good

helix formation has been obtained by using artificial bases, thereby improving binding conditions with regard to ionic strength and pH.

In addition, in order to improve half-life as well as membrane penetration, a large number of variations in polynucleotide backbones have
5 been done, nevertheless with little success.

Oligonucleotides can be modified either in the base, the sugar or the phosphate moiety. These modifications include, for example, the use of methylphosphonates, monothiophosphates, dithiophosphates, phosphoramidates, phosphate esters, bridged phosphorothioates, bridged
10 phosphoramidates, bridged methylenephosphonates, dephospho internucleotide analogs with siloxane bridges, carbonate bridges, carboxymethyl ester bridges, carbonate bridges, carboxymethyl ester bridges, acetamide bridges, carbamate bridges, thioether bridges, sulfoxyl bridges, sulfono bridges, various "plastic" DNAs, α -anomeric bridges and
15 borane derivatives. For further details the reader is referred to reference 76.

International patent application WO 89/12060 discloses various building blocks for synthesizing oligonucleotide analogs, as well as oligonucleotide analogs formed by joining such building blocks in a defined sequence. The building blocks may be either "rigid" (i.e., containing a ring
20 structure) or "flexible" (i.e., lacking a ring structure). In both cases, the building blocks contain a hydroxy group and a mercapto group, through which the building blocks are said to join to form oligonucleotide analogs. The linking moiety in the oligonucleotide analogs is selected from the group consisting of sulfide (-S-), sulfoxide (-SO-), and sulfone (-SO₂-). However,
25 the application provides no data supporting the specific binding of an oligonucleotide analog to a target oligonucleotide.

International patent application WO 92/20702 describe an acyclic oligonucleotide which includes a peptide backbone on which any selected chemical nucleobases or analogs are stringed and serve as coding characters
30 as they do in natural DNA or RNA. These new compounds, known as peptide nucleic acids (PNAs), are not only more stable in cells than their natural counterparts, but also bind natural DNA and RNA 50 to 100 times more tightly than the natural nucleic acids cling to each other (77). PNA oligomers can be synthesized from the four protected monomers containing
35 thymine, cytosine, adenine and guanine by Merrifield solid-phase peptide synthesis. In order to increase solubility in water and to prevent aggregation, a lysine amide group is placed at the C-terminal.

Thus, antisense technology requires pairing of messenger RNA with an oligonucleotide to form a double helix that inhibits translation. The concept of antisense-mediated gene therapy was already introduced in 1978 for cancer therapy. This approach was based on certain genes that are crucial in cell division and growth of cancer cells. Synthetic fragments of genetic substance DNA can achieve this goal. Such molecules bind to the targeted gene molecules in RNA of tumor cells, thereby inhibiting the translation of the genes and resulting in dysfunctional growth of these cells. Other mechanisms has also been proposed. These strategies have been used, with some success in treatment of cancers, as well as other illnesses, including viral and other infectious diseases. Antisense oligonucleotides are typically synthesized in lengths of 13-30 nucleotides. The life span of oligonucleotide molecules in blood is rather short. Thus, they have to be chemically modified to prevent destruction by ubiquitous nucleases present in the body. Phosphorothioates are very widely used modification in antisense oligonucleotide ongoing clinical trials (57). A new generation of antisense molecules consist of hybrid antisense oligonucleotide with a central portion of synthetic DNA while four bases on each end have been modified with 2'O-methyl ribose to resemble RNA. In preclinical studies in laboratory animals, such compounds have demonstrated greater stability to metabolism in body tissues and an improved safety profile when compared with the first-generation unmodified phosphorothioate (Hybridon Inc. news). Dosens of other nucleotide analogs have also been tested in antisense technology.

RNA oligonucleotides may also be used for antisense inhibition as they form a stable RNA-RNA duplex with the target, suggesting efficient inhibition. However, due to their low stability RNA oligonucleotides are typically expressed inside the cells using vectors designed for this purpose. This approach is favored when attempting to target a mRNA that encodes an abundant and long-lived protein (57).

Recent scientific publications have validated the efficacy of antisense compounds in animal models of hepatitis, cancers, coronary artery restenosis and other diseases. The first antisense drug was recently approved by the FDA. This drug Fomivirsen, developed by Isis, is indicated for local treatment of cytomegalovirus in patients with AIDS who are intolerant of or have a contraindication to other treatments for CMV retinitis or who were insufficiently responsive to previous treatments for CMV retinitis (Pharmacotherapy News Network).

Several antisense compounds are now in clinical trials in the United States. These include locally administered antivirals, systemic cancer therapeutics. Antisense therapeutics has the potential to treat many life-threatening diseases with a number of advantages over traditional drugs. Traditional drugs intervene after a disease-causing protein is formed. Antisense therapeutics, however, block mRNA transcription/translation and intervene before a protein is formed, and since antisense therapeutics target only one specific mRNA, they should be more effective with fewer side effects than current protein-inhibiting therapy.

A second option for disrupting gene expression at the level of transcription uses synthetic oligonucleotides capable of hybridizing with double stranded DNA. A triple helix is formed. Such oligonucleotides may prevent binding of transcription factors to the gene's promoter and therefore inhibit transcription. Alternatively, they may prevent duplex unwinding and, therefore, transcription of genes within the triple helical structure.

Another approach is the use of specific nucleic acid sequences to act as decoys for transcription factors. Since transcription factors bind specific DNA sequences it is possible to synthesize oligonucleotides that will effectively compete with the native DNA sequences for available transcription factors *in vivo*. This approach requires the identification of gene specific transcription factor (57).

Indirect inhibition of gene expression was demonstrated for matrix metalloproteinase genes (MMP-1, -3, and -9), which are associated with invasive potential of human cancer cells. E1AF is a transcription activator of MMP genes. Expression of E1AF antisense RNA in HSC3AS cells showed decrease in mRNA and protein levels of MMP-1, -3, and -9. Moreover, HSC3AS showed lower invasive potential *in vitro* and *in vivo*. These results imply that transfection of antisense inhibits tumor invasion by down-regulating MMP genes (58).

Ribozymes:

Ribozymes are being increasingly used for the sequence-specific inhibition of gene expression by the cleavage of mRNAs encoding proteins of interest. The possibility of designing ribozymes to cleave any specific target RNA has rendered them valuable tools in both basic research and therapeutic applications. In the therapeutics area, ribozymes have been exploited to target viral RNAs in infectious diseases, dominant oncogenes in cancers and specific somatic mutations in genetic disorders. Most notably, several ribozyme gene therapy protocols for HIV patients are

already in Phase 1 trials (62). More recently, ribozymes have been used for transgenic animal research, gene target validation and pathway elucidation. Several ribozymes are in various stages of clinical trials. ANGIOZYME was the first chemically synthesized ribozyme to be studied in human clinical trials. ANGIOZYME specifically inhibits formation of the VEGF-r (Vascular Endothelial Growth Factor receptor), a key component in the angiogenesis pathway. Ribozyme Pharmaceuticals, Inc., as well as other firms have demonstrated the importance of anti-angiogenesis therapeutics in animal models. HEPTAZYME, a ribozyme designed to selectively destroy Hepatitis C Virus (HCV) RNA, was found effective in decreasing Hepatitis C viral RNA in cell culture assays (Ribozyme Pharmaceuticals, Incorporated - WEB home page).

Gene disruption in animal models:

The emergence of gene inactivation by homologous recombination methodology in embryonic stem cells has revolutionized the field of mouse genetics. The availability of a rapidly growing number of mouse null mutants has represented an invaluable source of knowledge on mammalian development, cellular biology and physiology, and has provided many models for human inherited diseases. Animal models are required for an effective drug delivery development program and evaluation of gene therapy approach. The improvement of the original knockout strategy, as well as exploitation of exogenous enzymatic systems that are active in the recombination process, has been considerably extended the range of genetic manipulations that can be produced. Additional methods have been developed to provide versatile research tools: Double replacement method, sequential gene targeting, conditional cell type specific gene targeting, single copy integration method, inducible gene targeting, gene disruption by viral delivery, replacing one gene with another, the so called knock-in method and the induction of specific balanced chromosomal translocation. It is now possible to introduce a point mutation as a unique change in the entire genome, therefore allowing very fine dissection of gene function *in vivo*. Furthermore, the advent of methods allowing conditional gene targeting opens the way for analysis of consequence of a particular mutation in a defined organ and at a specific time during the life of the experimental animal (59).

DNA vaccination:

Observations in the early 1990s that plasmid DNA could directly transfect animal cells *in vivo* sparked exploration of the use of DNA

plasmids to induce immune response by direct injection into animal of DNA encoding antigenic protein. When a DNA vaccine plasmid enters the eukaryotic cell, the protein it encodes is transcribed and translated within the cell. In the case of pathogens, these proteins are presented to the immune system in their native form, mimicking the presentation of antigens during a natural infection. DNA vaccination is particularly useful for the induction of T cell activation. It was applied for viral and bacterial infectious diseases, as well as for allergy and for cancer. The central hypothesis behind active specific immunotherapy for cancer is that tumor cells express unique antigens that should stimulate the immune system. The first DNA vaccine against tumor was carcino-embryonic antigen (CEA). DNA vaccinated animals expressed immunoprotection and immunotherapy of human CEA-expressing syngeneic mouse colon and breast carcinoma (61). In a mouse model of neuroblastoma, DNA immunization with HuD resulted in tumor growth inhibition with no neurological disease (60). Immunity to the brown locus protein, gp⁷⁵ tyrosinase-related protein-1, associated with melanoma, was investigated in a syngeneic mouse model. Priming with human gp75 DNA broke tolerance to mouse gp75. Immunity against mouse gp75 provided significant tumor protection (60).

Glycosyl hydrolases:

Glycosyl hydrolases are a widespread group of enzymes that hydrolyze the o-glycosidic bond between two or more carbohydrates or between a carbohydrate and a noncarbohydrate moiety. The enzymatic hydrolysis of glycosidic bond occurs by using major one or two mechanisms leading to overall retention or inversion of the anomeric configuration. In both mechanisms catalysis involves two residues: a proton donor and a nucleophile. Glycosyl hydrolyses have been classified into 58 families based on amino acid similarities. The glycosyl hydrolyses from families 1, 2, 5, 10, 17, 30, 35, 39 and 42 act on a large variety of substrates, however, they all hydrolyze the glycosidic bond in a general acid catalysis mechanism, with retention of the anomeric configuration. The mechanism involves two glutamic acid residues, which are the proton donors and the nucleophile, with an asparagine always preceding the proton donor. Analyses of a set of known 3D structures from this group revealed that their catalytic domains, despite the low level of sequence identity, adopt a similar (α/β) 8 fold with the proton donor and the nucleophile located at the C-terminal ends of strands β 4 and β 7, respectively. Mutations

in the functional conserved amino acids of lysosomal glycosyl hydrolases were identified in lysosomal storage diseases.

Lysosomal glycosyl hydrolases including β -glucuronidase, β -mannosidase, β -glucocerebrosidase, β -galactosidase and α -L iduronidase, are all exo-glycosyl hydrolases, belong to the GH-A clan and share a similar catalytic site. However, many endo-glucanases from various organisms, such as bacterial and fungal xylanases and cellulases share this catalytic domain.

Genomic sequence of hpa gene and its implications:

It is well established that heparanase activity is correlated with cancer metastasis. This correlation was demonstrated at the level of enzymatic activity as well as the levels of protein and *hpa* cDNA expression in highly metastatic cancer cells as compared with non-metastatic cells. As such, inhibition of heparanase activity is desirable, and has been attempted by several means. The genomic region, encoding the *hpa* gene and the surrounding, provides a new powerful tool for regulation of heparanase activity at the level of gene expression. Regulatory sequences may reside in noncoding regions both upstream and downstream the transcribed region as well as in intron sequences. A DNA sequence upstream of the transcription start site contains the promoter region and potential regulatory elements. Regulatory factors, which interact with the promoter region may be identified and be used as potential drugs for inhibition of cancer, metastasis and inflammation. The promoter region can be used to screen for inhibitors of heparanase gene expression. Furthermore, the *hpa* promoter can be used to direct cell specific, particularly cancer cell specific, expression of foreign genes, such as cytotoxic or apoptotic genes, in order to specifically destroy cancer cells.

Cancer and yet unknown related genetic disorders may involve rearrangements and mutations in the heparanase gene, either in coding or non-coding regions. Such mutations may affect expression level or enzymatic activity. The genomic sequence of *hpa* enables the amplification of specific genomic DNA fragments, identification and diagnosis of mutations.

There is thus a widely recognized need for, and it would be highly advantageous to have genomic, cDNA and composite polynucleotides encoding a polypeptide having heparanase activity, vectors including same, genetically modified cells expressing heparanase and a recombinant protein

having heparanase activity, as well as antisense oligonucleotides, constructs and ribozymes which can be used for down regulation heparanase activity.

SUMMARY OF THE INVENTION

5 Cloning of the human *hpa* gene which encodes heparanase, and expression of recombinant heparanase by transfected host cells is reported herein, as well as downregulation of heparanase activity by antisense technology.

10 A purified preparation of heparanase isolated from human hepatoma cells was subjected to tryptic digestion and microsequencing. The YGPDVGQPR (SEQ ID NO:8) sequence revealed was used to screen EST databases for homology to the corresponding back translated DNA sequence. Two closely related EST sequences were identified and were thereafter found to be identical. Both clones contained an insert of 1020 bp
15 which included an open reading frame of 973 bp followed by a 27 bp of 3' untranslated region and a Poly A tail. Translation start site was not identified.

Cloning of the missing 5' end of *hpa* was performed by PCR amplification of DNA from placenta Marathon RACE cDNA composite
20 using primers selected according to the EST clones sequence and the linkers of the composite. A 900 bp PCR fragment, partially overlapping with the identified 3' encoding EST clones was obtained. The joined cDNA fragment (*hpa*), 1721 bp long (SEQ ID NO:9), contained an open reading frame which encodes a polypeptide of 543 amino acids (SEQ ID NO:10)
25 with a calculated molecular weight of 61,192 daltons.

Cloning an extended 5' sequence was enabled from the human SK-hep1 cell line by PCR amplification using the Marathon RACE. The 5' extended sequence of the SK-hep1 *hpa* cDNA was assembled with the sequence of the *hpa* cDNA isolated from human placenta (SEQ ID NO:9).
30 The assembled sequence contained an open reading frame, SEQ ID NOs: 13 and 15, which encodes, as shown in SEQ ID NOs:14 and 15, a polypeptide of 592 amino acids with a calculated molecular weight of 66,407 daltons.

The ability of the *hpa* gene product to catalyze degradation of heparan sulfate in an *in vitro* assay was examined by expressing the entire
35 open reading frame of *hpa* in insect cells, using the Baculovirus expression system. Extracts and conditioned media of cells infected with virus containing the *hpa* gene, demonstrated a high level of heparan sulfate degradation activity both towards soluble ECM-derived HSPG and intact

ECM. This degradation activity was inhibited by heparin, which is another substrate of heparanase. Cells infected with a similar construct containing no *hpa* gene had no such activity, nor did non-infected cells. The ability of heparanase expressed from the extended 5' clone towards heparin was demonstrated in a mammalian expression system.

The expression pattern of *hpa* RNA in various tissues and cell lines was investigated using RT-PCR. It was found to be expressed only in tissues and cells previously known to have heparanase activity.

A panel of monochromosomal human/CHO and human/mouse somatic cell hybrids was used to localize the human heparanase gene to human chromosome 4. The newly isolated heparanase sequence can be used to identify a chromosome region harboring a human heparanase gene in a chromosome spread.

A human genomic library was screened and the human locus harboring the heparanase gene isolated, sequenced and characterized. Alternatively spliced heparanase mRNAs were identified and characterized. The human heparanase promoter has been isolated, identified and positively tested for activity. The mouse heparanase promoter has been isolated and identified as well. Antisense heparanase constructs were prepared and their influence on cells *in vitro* tested. A predicted heparanase active site was identified. And finally, the presence of sequences hybridizing with human heparanase sequences was demonstrated for a variety of mammals and for an avian.

According to one aspect of the present invention there is provided an isolated nucleic acid comprising a genomic, complementary or composite polynucleotide sequence encoding a polypeptide having heparanase catalytic activity.

According to further features in preferred embodiments of the invention described below, the polynucleotide or a portion thereof is hybridizable with SEQ ID NOs: 9, 13, 42, 43 or a portion thereof at 68 °C in 6 x SSC, 1 % SDS, 5 x Denharts, 10 % dextran sulfate, 100 µg/ml salmon sperm DNA, and ³²p labeled probe and wash at 68 °C with 3 x SSC and 0.1 % SDS.

According to still further features in the described preferred embodiments the polynucleotide or a portion thereof is at least 60 % identical with SEQ ID NOs: 9, 13, 42, 43 or portions thereof as determined using the Bestfit procedure of the DNA sequence analysis software package

developed by the Genetic Computer Group (GCG) at the university of Wisconsin (gap creation penalty - 12, gap extension penalty - 4).

According to still further features in the described preferred embodiments the polypeptide is as set forth in SEQ ID NOs:10, 14, 44 or
5 portions thereof.

According to still further features in the described preferred embodiments the polypeptide is at least 60 % homologous to SEQ ID NOs:10, 14, 44 or portions thereof as determined with the Smith-Waterman algorithm, using the Bioaccelerator platform developed by Compugene
10 (gapop: 10.0, gapext: 0.5, matrix: blosum62).

According to additional aspects of the present invention there are provided a nucleic acid construct (vector) comprising the isolated nucleic acid described herein and a host cell comprising the construct.

According to a further aspect of the present invention there is
15 provided an antisense oligonucleotide comprising a polynucleotide or a polynucleotide analog of at least 10 bases being hybridizable *in vivo*, under physiological conditions, with a portion of a polynucleotide strand encoding a polypeptide having heparanase catalytic activity.

According to an additional aspect of the present invention there is
20 provided a method of *in vivo* downregulating heparanase activity comprising the step of *in vivo* administering the antisense oligonucleotide herein described.

According to yet an additional aspect of the present invention there is provided a pharmaceutical composition comprising the antisense
25 oligonucleotide herein described and a pharmaceutically acceptable carrier.

According to still an additional aspect of the present invention there is provided a ribozyme comprising the antisense oligonucleotide described herein and a ribozyme sequence.

According to a further aspect of the present invention there is
30 provided an antisense nucleic acid construct comprising a promoter sequence and a polynucleotide sequence directing the synthesis of an antisense RNA sequence of at least 10 bases being hybridizable *in vivo*, under physiological conditions, with a portion of a polynucleotide strand encoding a polypeptide having heparanase catalytic activity.

35 According to further features in preferred embodiments of the invention described below, the polynucleotide strand encoding the polypeptide having heparanase catalytic activity is as set forth in SEQ ID NOs: 9, 13, 42 or 43.

According to still further features in the described preferred embodiments the polypeptide having heparanase catalytic activity is as set forth in SEQ ID NOs: 10, 14 or 44.

According to still a further aspect of the present invention there is provided a method of *in vivo* downregulating heparanase activity comprising the step of *in vivo* administering the antisense nucleic acid construct herein described.

According to yet a further aspect of the present invention there is provided a pharmaceutical composition comprising the antisense nucleic acid construct herein described and a pharmaceutically acceptable carrier.

According to a further aspect of the present invention there is provided a nucleic acid construct comprising a polynucleotide sequence functioning as a promoter, the polynucleotide sequence is derived from SEQ ID NO:42 and includes at least nucleotides 2535-2635 thereof or from SEQ ID NO:43 and includes at least nucleotides 320-420.

According to a further aspect of the present invention there is provided a method of expressing a polynucleotide sequence comprising the step of ligating the polynucleotide sequence into the nucleic acid construct described above, downstream of the polynucleotide sequence derived from SEQ ID NOs:42 or 43.

According to a further aspect of the present invention there is provided a recombinant protein comprising a polypeptide having heparanase catalytic activity.

According to further features in preferred embodiments of the invention described below, the polypeptide includes at least a portion of SEQ ID NOs:10, 14 or 44.

According to still further features in the described preferred embodiments the protein is encoded by a polynucleotide hybridizable with SEQ ID NOs: 9, 13, 42, 43 or a portion thereof at 68 °C in 6 x SSC, 1 % SDS, 5 x Denharts, 10 % dextran sulfate, 100 µg/ml salmon sperm DNA, and ³²p labeled probe and wash at 68 °C with 3 x SSC and 0.1 % SDS.

According to still further features in the described preferred embodiments the protein is encoded by a polynucleotide at least 60 % identical with SEQ ID NOs: 9, 13, 42, 43 or portions thereof as determined using the Bestfit procedure of the DNA sequence analysis software package developed by the Genetic Computer Group (GCG) at the university of Wisconsin (gap creation penalty - 12, gap extension penalty - 4).

According to a further aspect of the present invention there is provided a pharmaceutical composition comprising, as an active ingredient, the recombinant protein herein described.

According to a further aspect of the present invention there is provided a method of identifying a chromosome region harboring a heparanase gene in a chromosome spread comprising the steps of (a) hybridizing the chromosome spread with a tagged polynucleotide probe encoding heparanase; (b) washing the chromosome spread, thereby removing excess of non-hybridized probe; and (c) searching for signals associated with the hybridized tagged polynucleotide probe, wherein detected signals being indicative of a chromosome region harboring a heparanase gene.

According to a further aspect of the present invention there is provided a method of *in vivo* eliciting anti-heparanase antibodies comprising the steps of administering a nucleic acid construct including a polynucleotide segment corresponding to at least a portion of SEQ ID NOs:9, 13 or 43 and a promoter for directing the expression of said polynucleotide segment *in vivo*. Accordingly, there is provided also a DNA vaccine for *in vivo* eliciting anti-heparanase antibodies comprising a nucleic acid construct including a polynucleotide segment corresponding to at least a portion of SEQ ID NOs:9, 13 or 43 and a promoter for directing the expression of said polynucleotide segment *in vivo*.

The present invention can be used to develop new drugs to inhibit tumor cell metastasis, inflammation and autoimmunity. The identification of the *hpa* gene encoding for heparanase enzyme enables the production of a recombinant enzyme in heterologous expression systems. Additional features, advantages, uses and applications of the present invention in biological science and in diagnostic and therapeutic medicine are described hereinafter.

BRIEF DESCRIPTION OF THE DRAWINGS

The invention herein described, by way of example only, with reference to the accompanying drawings, wherein:

FIG. 1 presents nucleotide sequence and deduced amino acid sequence of *hpa* cDNA. A single nucleotide difference at position 799 (A to T) between the EST (Expressed Sequence Tag) and the PCR amplified cDNA (reverse transcribed RNA) and the resulting amino acid substitution (Tyr to Phe) are indicated above and below the substituted unit,

respectively. Cysteine residues and the poly adenylation consensus sequence are underlined. The asterisk denotes the stop codon TGA.

FIG. 2 demonstrates degradation of soluble sulfate labeled HSPG substrate by lysates of High Five cells infected with pFhpa2 virus. Lysates of High Five cells that were infected with pFhpa2 virus (●) or control pF2 virus (□) were incubated (18 h, 37 °C) with sulfate labeled ECM-derived soluble HSPG (peak I). The incubation medium was then subjected to gel filtration on Sepharose 6B. Low molecular weight HS degradation fragments (peak II) were produced only during incubation with the pFhpa2 infected cells, but there was no degradation of the HSPG substrate (✧) by lysates of pF2 infected cells.

FIGs. 3a-b demonstrate degradation of soluble sulfate labeled HSPG substrate by the culture medium of pFhpa2 and pFhpa4 infected cells. Culture media of High Five cells infected with pFhpa2 (3a) or pFhpa4 (3b) viruses (●), or with control viruses (□) were incubated (18 h, 37 °C) with sulfate labeled ECM-derived soluble HSPG (peak I, ✧). The incubation media were then subjected to gel filtration on Sepharose 6B. Low molecular weight HS degradation fragments (peak II) were produced only during incubation with the hpa gene containing viruses. There was no degradation of the HSPG substrate by the culture medium of cells infected with control viruses.

FIG. 4 presents size fractionation of heparanase activity expressed by pFhpa2 infected cells. Culture medium of pFhpa2 infected High Five cells was applied onto a 50 kDa cut-off membrane. Heparanase activity (conversion of the peak I substrate, ✧ into peak II HS degradation fragments) was found in the high (> 50 kDa) (●), but not low (< 50 kDa) (○) molecular weight compartment.

FIGs. 5a-b demonstrate the effect of heparin on heparanase activity expressed by pFhpa2 and pFhpa4 infected High Five cells. Culture media of pFhpa2 (5a) and pFhpa4 (5b) infected High Five cells were incubated (18 h, 37 °C) with sulfate labeled ECM-derived soluble HSPG (peak I, ✧) in the absence (●) or presence (Δ) of 10 μg/ml heparin. Production of low molecular weight HS degradation fragments was completely abolished in the presence of heparin, a potent inhibitor of heparanase activity (6, 7).

FIGs. 6a-b demonstrate degradation of sulfate labeled intact ECM by virus infected High Five and Sf21 cells. High Five (6a) and Sf21 (6b) cells were plated on sulfate labeled ECM and infected (48 h, 28 °C) with pFhpa4 (●) or control pF1 (□) viruses. Control non-infected Sf21 cells (R) were

plated on the labeled ECM as well. The pH of the cultured medium was adjusted to 6.0 - 6.2 followed by 24 h incubation at 37 °C. Sulfate labeled material released into the incubation medium was analyzed by gel filtration on Sepharose 6B. HS degradation fragments were produced only by cells infected with the *hpa* containing virus.

FIG. 7a-b demonstrate degradation of sulfate labeled intact ECM by virus infected cells. High Five (7a) and Sf21 (7b) cells were plated on sulfate labeled ECM and infected (48 h, 28 °C) with pF*hpa*4 (●) or control pF1 (□) viruses. Control non-infected Sf21 cells (R) were plate on labeled ECM as well. The pH of the cultured medium was adjusted to 6.0 - 6.2, followed by 48 h incubation at 28 °C. Sulfate labeled degradation fragments released into the incubation medium was analyzed by gel filtration on Sepharose 6B. HS degradation fragments were produced only by cells infected with the *hpa* containing virus.

FIGs. 8a-b demonstrate degradation of sulfate labeled intact ECM by the culture medium of pF*hpa*4 infected cells. Culture media of High Five (8a) and Sf21 (8b) cells that were infected with pF*hpa*4 (●) or control pF1 (□) viruses were incubated (48 h, 37 °C, pH 6.0) with intact sulfate labeled ECM. The ECM was also incubated with the culture medium of control non-infected Sf21 cells (R). Sulfate labeled material released into the reaction mixture was subjected to gel filtration analysis. Heparanase activity was detected only in the culture medium of pF*hpa*4 infected cells.

FIGs. 9a-b demonstrate the effect of heparin on heparanase activity in the culture medium of pF*hpa*4 infected cells. Sulfate labeled ECM was incubated (24 h, 37 °C, pH 6.0) with culture medium of pF*hpa*4 infected High Five (9a) and Sf21 (9b) cells in the absence (●) or presence (V) of 10 µg/ml heparin. Sulfate labeled material released into the incubation medium was subjected to gel filtration on Sepharose 6B. Heparanase activity (production of peak II HS degradation fragments) was completely inhibited in the presence of heparin.

FIGs. 10a-b demonstrate purification of recombinant heparanase on heparin-Sepharose. Culture medium of Sf21 cells infected with pF*hpa*4 virus was subjected to heparin-Sepharose chromatography. Elution of fractions was performed with 0.35 - 2 M NaCl gradient (◇). Heparanase activity in the eluted fractions is demonstrated in Figure 10a (●). Fractions 15-28 were subjected to 15 % SDS-polyacrylamide gel electrophoresis followed by silver nitrate staining. A correlation is demonstrated between a

major protein band (MW ~ 63,000) in fractions 19 - 24 and heparanase activity.

FIGs. 11a-b demonstrate purification of recombinant heparanase on a Superdex 75 gel filtration column. Active fractions eluted from heparin-Sepharose (Figure 10a) were pooled, concentrated and applied onto Superdex 75 FPLC column. Fractions were collected and aliquots of each fraction were tested for heparanase activity (C, Figure 11a) and analyzed by SDS-polyacrylamide gel electrophoresis followed by silver nitrate staining (Figure 11b). A correlation is seen between the appearance of a major protein band (MW ~ 63,000) in fractions 4 - 7 and heparanase activity.

FIGs. 12a-e demonstrate expression of the *hpa* gene by RT-PCR with total RNA from human embryonal tissues (12a), human extra-embryonal tissues (12b) and cell lines from different origins (12c-e). RT-PCR products using *hpa* specific primers (I), primers for GAPDH housekeeping gene (II), and control reactions without reverse transcriptase demonstrating absence of genomic DNA or other contamination in RNA samples (III). M- DNA molecular weight marker VI (Boehringer Mannheim). For 12a: lane 1 - neutrophil cells (adult), lane 2 - muscle, lane 3 - thymus, lane 4 - heart, lane 5 - adrenal. For 12b: lane 1 - kidney, lane 2 - placenta (8 weeks), lane 3 - placenta (11 weeks), lanes 4-7 - mole (complete hydatidiform mole), lane 8 - cytotrophoblast cells (freshly isolated), lane 9 - cytotrophoblast cells (1.5 h *in vitro*), lane 10 - cytotrophoblast cells (6 h *in vitro*), lane 11 - cytotrophoblast cells (18 h *in vitro*), lane 12 - cytotrophoblast cells (48 h *in vitro*). For 12c: lane 1 - JAR bladder cell line, lane 2 - NCITT testicular tumor cell line, lane 3 - SW-480 human hepatoma cell line, lane 4 - HTR (cytotrophoblasts transformed by SV40), lane 5 - HPTLP-I hepatocellular carcinoma cell line, lane 6 - EJ-28 bladder carcinoma cell line. For 12d: lane 1 - SK-hep-1 human hepatoma cell line, lane 2 - DAMI human megakaryocytic cell line, lane 3 - DAMI cell line + PMA, lane 4 - CHRF cell line + PMA, lane 5 - CHRF cell line. For 12e: lane 1 - ABAE bovine aortic endothelial cells, lane 2 - 1063 human ovarian cell line, lane 3 - human breast carcinoma MDA435 cell line, lane 4 - human breast carcinoma MDA231 cell line.

FIG. 13 presents a comparison between nucleotide sequences of the human *hpa* and a mouse EST cDNA fragment (SEQ ID NO:12) which is 80 % homologous to the 3' end (starting at nucleotide 1066 of SEQ ID NO:9) of the human *hpa*. The aligned termination codons are underlined.

FIG. 14 demonstrates the chromosomal localization of the *hpa* gene. PCR products of DNA derived from somatic cell hybrids and of genomic DNA of hamster, mouse and human of were separated on 0.7 % agarose gel following amplification with *hpa* specific primers. Lane 1 – Lambda DNA digested with *Bst*EII, lane 2 – no DNA control, lanes 3 – 29, PCR amplification products. Lanes 3-5 – human, mouse and hamster genomic DNA, respectively. Lanes 6-29, human monochromosomal somatic cell hybrids representing chromosomes 1-22 and X and Y, respectively. Lane 30 – Lambda DNA digested with *Bst*EII. An amplification product of approximately 2.8 Kb is observed only in lanes 5 and 9, representing human genomic DNA and DNA derived from cell hybrid carrying human chromosome 4, respectively. These results demonstrate that the *hpa* gene is localized in human chromosome 4.

FIG. 15 demonstrates the genomic exon-intron structure of the human *hpa* locus (top) and the relative positions of the lambda clones used as sequencing templates to sequence the locus (below). The vertical rectangles represent exons (E) and the horizontal lines therebetween represent introns (I), upstream (U) and downstream (D) regions. Continuous lines represent DNA fragments, which were used for sequence analysis. The discontinuous line in lambda 6 represent a region, which overlaps with lambda 8 and hence was not analyzed. The plasmid contains a PCR product, which bridges the gap between L3 and L6.

FIG. 16 presents the nucleotide sequence of the genomic region of the *hpa* gene. Exon sequences appear in upper case and intron sequences in lower case. The deduced amino acid sequence of the exons is printed below the nucleotide sequence. Two predicted transcription start sites are shown in bold.

FIG. 17 presents an alignment of the amino acid sequences of human heparanase, mouse and partial sequences of rat homologues. The human and the mouse sequences were determined by sequence analysis of the isolated cDNAs. The rat sequence is derived from two different EST clones, which represent two different regions (5' and 3') of the rat *hpa* cDNA. The human sequence and the amino acids in the mouse and rat homologues, which are identical to the human sequence, appear in bold.

FIG. 18 presents a heparanase Zoo blot. Ten micrograms of genomic DNA from various sources were digested with *Eco*RI and separated on 0.7 % agarose – TBE gel. Following electrophoresis, the was gel treated with HCl and than with NaOH and the DNA fragments were downward

transferred to a nylon membrane (Hybond N+, Amersham) with 0.4 N NaOH. The membrane was hybridized with a 1.6 Kb DNA probe that contained the entire *hpa* cDNA. Lane order: H – Human; M – Mouse; Rt – Rat; P – Pig; Cw – Cow; Hr – Horse; S – Sheep; Rb – Rabbit; D – Dog; Ch – Chicken; F – Fish. Size markers (Lambda *Bst*II) are shown on the left

FIG. 19 demonstrates the secondary structure prediction for heparanase performed using the PHD server – Profile network Prediction Heidelberg. H – helix, E – extended (beta strand), The glutamic acid predicted as the proton donor is marked by asterisk and the possible nucleophiles are underlined.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention is of a polynucleotide or nucleic acid, referred to hereinbelow interchangeably as *hpa*, *hpa* cDNA or *hpa* gene or identified by its SEQ ID NOs, encoding a polypeptide having heparanase activity, vectors or nucleic acid constructs including same and which are used for over-expression or antisense inhibition of heparanase, genetically modified cells expressing same, recombinant protein having heparanase activity, antisense oligonucleotides and ribozymes for heparanase modulation, and heparanase promoter sequences which can be used to direct the expression of desired genes.

Before explaining at least one embodiment of the invention in detail, it is to be understood that the invention is not limited in its application to the details of construction and the arrangement of the components set forth in the following description or illustrated in the drawings. The invention is capable of other embodiments or of being practiced or carried out in various ways. Also, it is to be understood that the phraseology and terminology employed herein is for the purpose of description and should not be regarded as limiting.

Cloning of the human and mouse *hpa* genes, cDNAs and genomic sequence (for human), encoding heparanase and expressing recombinant heparanase by transfected cells is reported herein. These are the first mammalian heparanase genes to be cloned.

A purified preparation of heparanase isolated from human hepatoma cells was subjected to tryptic digestion and microsequencing.

The YGPDVGQPR (SEQ ID NO:8) sequence revealed was used to screen EST databases for homology to the corresponding back translated

DNA sequences. Two closely related EST sequences were identified and were thereafter found to be identical.

Both clones contained an insert of 1020 bp which includes an open reading frame of 973 bp followed by a 3' untranslated region of 27 bp and a Poly A tail, whereas a translation start site was not identified.

Cloning of the missing 5' end was performed by PCR amplification of DNA from placenta Marathon RACE cDNA composite using primers selected according to the EST clones sequence and the linkers of the composite.

A 900 bp PCR fragment, partially overlapping with the identified 3' encoding EST clones was obtained. The joined cDNA fragment (*hpa*), 1721 bp long (SEQ ID NO:9), contained an open reading frame which encodes, as shown in Figure 1 and SEQ ID NO:11, a polypeptide of 543 amino acids (SEQ ID NO:10) with a calculated molecular weight of 61,192 daltons.

A single nucleotide difference at position 799 (A to T) between the EST clones and the PCR amplified cDNA was observed. This difference results in a single amino acid substitution (Tyr to Phe) (Figure 1). Furthermore, the published EST sequences contained an unidentified nucleotide, which following DNA sequencing of both the EST clones was resolved into two nucleotides (G and C at positions 1630 and 1631 in SEQ ID NO:9, respectively).

The ability of the *hpa* gene product to catalyze degradation of heparan sulfate in an *in vitro* assay was examined by expressing the entire open reading frame in insect cells, using the Baculovirus expression system.

Extracts and conditioned media of cells infected with virus containing the *hpa* gene, demonstrated a high level of heparan sulfate degradation activity both towards soluble ECM-derived HSPG and intact ECM, which was inhibited by heparin, while cells infected with a similar construct containing no *hpa* gene had no such activity, nor did non-infected cells.

The expression pattern of *hpa* RNA in various tissues and cell lines was investigated using RT-PCR. It was found to be expressed only in tissues and cells previously known to have heparanase activity.

Cloning an extended 5' sequence was enabled from the human SK-hep1 cell line by PCR amplification using the Marathon RACE. The 5' extended sequence of the SK-hep1 *hpa* cDNA was assembled with the sequence of the *hpa* cDNA isolated from human placenta (SEQ ID NO:9).

The assembled sequence contained an open reading frame, SEQ ID NOs: 13 and 15, which encodes, as shown in SEQ ID NOs:14 and 15, a polypeptide of 592 amino acids, with a calculated molecular weight of 66,407 daltons. This open reading frame was shown to direct the expression of catalytically active heparanase in a mammalian cell expression system. The expressed heparanase was detectable by anti heparanase antibodies in Western blot analysis.

A panel of monochromosomal human/CHO and human/mouse somatic cell hybrids was used to localize the human heparanase gene to human chromosome 4. The newly isolated heparanase sequence can therefore be used to identify a chromosome region harboring a human heparanase gene in a chromosome spread.

The *hpa* cDNA was then used as a probe to screen a human genomic library. Several phages were positive. These phages were analyzed and were found to cover most of the *hpa* locus, except for a small portion which was recovered by bridging PCR. The *hpa* locus covers about 50,000 bp. The *hpa* gene includes 12 exons separated by 11 introns.

RT-PCR performed on a variety of cells revealed alternatively spliced *hpa* transcripts.

The amino acid sequence of human heparanase was used to search for homologous sequences in the DNA and protein databases. Several human EST's were identified, as well as mouse sequences highly homologous to human heparanase. The following mouse EST's were identified AA177901, AA674378, AA67997, AA047943, AA690179, AI122034, all sharing an identical sequence and correspond to amino acids 336-543 of the human heparanase sequence. The entire mouse heparanase cDNA was cloned, based on the nucleotide sequence of the mouse EST's using Marathon cDNA libraries. The mouse and the human *hpa* genes share an average homology of 78 % between the nucleotide sequences and 81 % similarity between the deduced amino acid sequences. *hpa* homologous sequences from rat were also uncovered (EST's AI060284 and AI237828).

Homology search of heparanase amino acid sequence against the DNA and the protein databases and prediction of its protein secondary structure enabled to identify candidate amino acids that participate in the heparanase active site.

Expression of *hpa* antisense in mammalian cell lines resulted in about five fold decrease in the number of recoverable cells as compared to controls.

Human *Hpa* cDNA was shown to hybridize with genomic DNAs of a variety of mammalian species and with an avian.

The human and mouse *hpa* promoters were identified and the human promoter was tested positive in directing the expression of a reporter gene.

Thus, according to the present invention there is provided an isolated nucleic acid comprising a genomic, complementary or composite polynucleotide sequence encoding a polypeptide having heparanase catalytic activity.

The phrase "composite polynucleotide sequence" refers to a sequence which includes exonal sequences required to encode the polypeptide having heparanase activity, as well as any number of intronal sequences. The intronal sequences can be of any source and typically will include conserved splicing signal sequences. Such intronal sequences may further include cis acting expression regulatory elements.

The term "heparanase catalytic activity" or its equivalent term "heparanase activity" both refer to a mammalian endoglycosidase hydrolyzing activity which is specific for heparan or heparan sulfate proteoglycan substrates, as opposed to the activity of bacterial enzymes (heparinase I, II and III) which degrade heparin or heparan sulfate by means of β -elimination (37).

According to a preferred embodiment of the present invention the polynucleotide or a portion thereof is hybridizable with SEQ ID NOs: 9, 13, 42, 43 or a portion thereof at 68 °C in 6 x SSC, 1 % SDS, 5 x Denharts, 10 % dextran sulfate, 100 μ g/ml salmon sperm DNA, and ³²p labeled probe and wash at 68 °C with 3, 2, 1, 0.5 or 0.1 x SSC and 0.1 % SDS.

According to another preferred embodiment of the present invention the polynucleotide or a portion thereof is at least 60 %, preferably at least 65 %, more preferably at least 70 %, more preferably at least 75 %, more preferably at least 80 %, more preferably at least 85 %, more preferably at least 90 %, most preferably, 95-100 % identical with SEQ ID NOs: 9, 13, 42, 43 or portions thereof as determined using the Bestfit procedure of the DNA sequence analysis software package developed by the Genetic Computer Group (GCG) at the university of Wisconsin (gap creation penalty - 12, gap extension penalty - 4 - which are the default parameters).

According to another preferred embodiment of the present invention the polypeptide encoded by the polynucleotide sequence is as set forth in SEQ ID NOs:10, 14, 44 or portions thereof having heparanase catalytic activity. Such portions are expected to include amino acids Asp-Glu 224-225 (SEQ ID NO:10), which can serve as proton donors and glutamic acid 343 or 396 which can serve as a nucleophile.

According to another preferred embodiment of the present invention the polypeptide encoded by the polynucleotide sequence is at least 60 %, preferably at least 65 %, more preferably at least 70 %, more preferably at least 75 %, more preferably at least 80 %, more preferably at least 85 %, more preferably at least 90 %, most preferably, 95-100 % homologous (both similar and identical acids) to SEQ ID NOs:10, 14, 44 or portions thereof as determined with the Smith-Waterman algorithm, using the Bioaccelerator platform developed by Compugene (gapop: 10.0, gapext: 0.5, matrix: blosum62, see also the description to Figure 17).

Further according to the present invention there is provided a nucleic acid construct comprising the isolated nucleic acid described herein. The construct may and preferably further include an origin of replication and

The construct or vector can be of any type. It may be a phage which infects bacteria or a virus which infects eukaryotic cells. It may also be a plasmid, phagemid, cosmid, bacmid or an artificial chromosome.

Further according to the present invention there is provided a host cell comprising the nucleic acid construct described herein. The host cell can be of any type. It may be a prokaryotic cell, an eukaryotic cell, a cell line, or a cell as a portion of an organism. The polynucleotide encoding heparanase can be permanently or transiently present in the cell. In other words, genetically modified cells obtained following stable or transient transfection, transformation or transduction are all within the scope of the present invention. The polynucleotide can be present in the cell in low copy (say 1-5 copies) or high copy number (say 5-50 copies or more). It may be integrated in one or more chromosomes at any location or be present as an extrachromosomal material.

The present invention is further directed at providing a heparanase over-expression system which includes a cell overexpressing heparanase catalytic activity. The cell may be a genetically modified host cell transiently or stably transfected or transformed with any suitable vector which includes a polynucleotide sequence encoding a polypeptide having

heparanase activity and a suitable promoter and enhancer sequences to direct over-expression of heparanase. However, the overexpressing cell may also be a product of an insertion (e.g., via homologous recombination) of a promoter and/or enhancer sequence downstream to the endogenous
5 heparanase gene of the expressing cell, which will direct over-expression from the endogenous gene.

The term "over-expression" as used herein in the specification and claims below refers to a level of expression which is higher than a basal level of expression typically characterizing a given cell under otherwise
10 identical conditions.

According to another aspect the present invention provides an antisense oligonucleotide comprising a polynucleotide or a polynucleotide analog of at least 10, preferably 11-15, more preferably 16-17, more preferably 18, more preferably 19-25, more preferably 26-35, most
15 preferably 35-100 bases being hybridizable *in vivo*, under physiological conditions, with a portion of a polynucleotide strand encoding a polypeptide having heparanase catalytic activity. The antisense oligonucleotide can be used for downregulating heparanase activity by *in vivo* administration thereof to a patient. As such, the antisense oligonucleotide according to the
20 present invention can be used to treat types of cancers which are characterized by impaired (over) expression of heparanase, and are dependent on the expression of heparanase for proliferating or forming metastases.

The antisense oligonucleotide can be DNA or RNA or even include
25 nucleotide analogs, examples of which are provided in the Background section hereinabove. The antisense oligonucleotide according to the present invention can be synthetic and is preferably prepared by solid phase synthesis. In addition, it can be of any desired length which still provides specific base pairing (e.g., 8 or 10, preferably more, nucleotides long) and it
30 can include mismatches that do not hamper base pairing under physiological conditions.

Further according to the present invention there is provided a pharmaceutical composition comprising the antisense oligonucleotide herein described and a pharmaceutically acceptable carrier. The carrier can
35 be, for example, a liposome loadable with the antisense oligonucleotide.

According to a preferred embodiment of the present invention the antisense oligonucleotide further includes a ribozyme sequence. The ribozyme sequence serves to cleave a heparanase RNA molecule to which

the antisense oligonucleotide binds, to thereby downregulate heparanase expression.

Further according to the present invention there is provided an antisense nucleic acid construct comprising a promoter sequence and a polynucleotide sequence directing the synthesis of an antisense RNA sequence of at least 10 bases being hybridizable *in vivo*, under physiological conditions, with a portion of a polynucleotide strand encoding a polypeptide having heparanase catalytic activity. Like the antisense oligonucleotide, the antisense construct can be used for downregulating heparanase activity by *in vivo* administration thereof to a patient. As such, the antisense construct, like the antisense oligonucleotide, according to the present invention can be used to treat types of cancers which are characterized by impaired (over) expression of heparanase, and are dependent on the expression of heparanase for proliferating or forming metastases.

Thus, further according to the present invention there is provided a pharmaceutical composition comprising the antisense construct herein described and a pharmaceutically acceptable carrier. The carrier can be, for example, a liposome loadable with the antisense construct.

Formulations for topical administration may include, but are not limited to, lotions, ointments, gels, creams, suppositories, drops, liquids, sprays and powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable. Coated condoms, stents, active pads, and other medical devices may also be useful. Compositions for oral administration include powders or granules, suspensions or solutions in water or non-aqueous media, sachets, capsules or tablets. Thickeners, diluents, flavorings, dispersing aids, emulsifiers or binders may be desirable. Formulations for parenteral administration may include, but are not limited to, sterile aqueous solutions which may also contain buffers, diluents and other suitable additives.

Dosing is dependent on severity and responsiveness of the condition to be treated, but will normally be one or more doses per day, week or month with course of treatment lasting from several days to several months or until a cure is effected or a diminution of disease state is achieved. Persons ordinarily skilled in the art can easily determine optimum dosages, dosing methodologies and repetition rates.

Further according to the present invention there is provided a nucleic acid construct comprising a polynucleotide sequence functioning as a promoter, the polynucleotide sequence is derived from SEQ ID NO:42 and

includes at least nucleotides 2135-2635, preferably 2235-2635, more preferably 2335-2635, more preferably 2435-2635, most preferably 2535-2635 thereof, or SEQ ID NO:43 and includes at least nucleotides 1-420, preferably 120-420, more preferably 220-420, most preferably 320-420, thereof. These nucleotides are shown in the example section that follows to direct the synthesis of a reporter gene in transformed cells. Thus, further according to the present invention there is provided a method of expressing a polynucleotide sequence comprising the step of ligating the polynucleotide sequence downstream to either of the promoter sequences described herein. Heparanase promoters can be isolated from a variety of mammalian and other species by cloning genomic regions present 5' to the coding sequence thereof. This can be readily achievable by one ordinarily skilled in the art using the heparanase polynucleotides described herein, which are shown in the Examples section that follows to participate in efficient cross species hybridization.

Further according to the present invention there is provided a recombinant protein comprising a polypeptide having heparanase catalytic activity. The protein according to the present invention include modifications known as post translational modifications, including, but not limited to, proteolysis (e.g., removal of a signal peptide and of a pro- or preprotein sequence), methionine modification, glycosylation, alkylation (e.g., methylation), acetylation, etc. According to preferred embodiments the polypeptide includes at least a portion of SEQ ID NOs:10, 14 or 44, the portion has heparanase catalytic activity. According to preferred embodiments of the present invention the protein is encoded by any of the above described isolated nucleic acids. Further according to the present invention there is provided a pharmaceutical composition comprising, as an active ingredient, the recombinant protein described herein.

The recombinant protein may be purified by any conventional protein purification procedure close to homogeneity and/or be mixed with additives. The recombinant protein may be manufactured using any of the genetically modified cells described above, which include any of the expression nucleic acid constructs described herein. The recombinant protein may be in any form. It may be in a crystallized form, a dehydrated powder form or in solution. The recombinant protein may be useful in obtaining pure heparanase, which in turn may be useful in eliciting anti-heparanase antibodies, either poly or monoclonal antibodies, and as a

screening active ingredient in an anti-heparanase inhibitors or drugs screening assay or system.

Further according to the present invention there is provided a method of identifying a chromosome region harboring a human heparanase gene in a chromosome spread. the method is executed implementing the following method steps, in which in a first step the chromosome spread (either interphase or metaphase spread) is hybridized with a tagged polynucleotide probe encoding heparanase. The tag is preferably a fluorescent tag. In a second step according to the method the chromosome spread is washed, thereby excess of non-hybridized probe is removed. Finally, signals associated with the hybridized tagged polynucleotide probe are searched for, wherein detected signals being indicative of a chromosome region harboring the human heparanase gene. One ordinarily skilled in the art would know how to use the sequences disclosed herein in suitable labeling reactions and how to use the tagged probes to detect, using *in situ* hybridization, a chromosome region harboring a human heparanase gene.

Further according to the present invention there is provided a method of *in vivo* eliciting anti-heparanase antibodies comprising the steps of administering a nucleic acid construct including a polynucleotide segment corresponding to at least a portion of SEQ ID NOs:9, 13 or 43 and a promoter for directing the expression of said polynucleotide segment *in vivo*. Accordingly, there is provided also a DNA vaccine for *in vivo* eliciting anti-heparanase antibodies comprising a nucleic acid construct including a polynucleotide segment corresponding to at least a portion of SEQ ID NOs:9, 13 or 43 and a promoter for directing the expression of said polynucleotide segment *in vivo*. The vaccine optionally further includes a pharmaceutically acceptable carrier, such as a virus, liposome or an antigen presenting cell. Alternatively, the vaccine is employed as a naked DNA vaccine

The present invention can be used to develop treatments for various diseases, to develop diagnostic assays for these diseases and to provide new tools for basic research especially in the fields of medicine and biology.

Specifically, the present invention can be used to develop new drugs to inhibit tumor cell metastasis, inflammation and autoimmunity. The identification of the *hpa* gene encoding for the heparanase enzyme enables the production of a recombinant enzyme in heterologous expression systems.

Furthermore, the present invention can be used to modulate bioavailability of heparin-binding growth factors, cellular responses to heparin-binding growth factors (e.g., bFGF, VEGF) and cytokines (e.g., IL-8), cell interaction with plasma lipoproteins, cellular susceptibility to viral, protozoa and some bacterial infections, and disintegration of neurodegenerative plaques. Recombinant heparanase offers a potential treatment for wound healing, angiogenesis, restenosis, atherosclerosis, inflammation, neurodegenerative diseases (such as, for example, Genstmann-Straussler Syndrome, Creutzfeldt-Jakob disease, Scrape and Alzheimer's disease) and certain viral and some bacterial and protozoa infections. Recombinant heparanase can be used to neutralize plasma heparin, as a potential replacement of protamine.

As used herein, the term "modulate" includes substantially inhibiting, slowing or reversing the progression of a disease, substantially ameliorating clinical symptoms of a disease or condition, or substantially preventing the appearance of clinical symptoms of a disease or condition. A "modulator" therefore includes an agent which may modulate a disease or condition. Modulation of viral, protozoa and bacterial infections includes any effect which substantially interrupts, prevents or reduces any viral, bacterial or protozoa activity and/or stage of the virus, bacterium or protozoon life cycle, or which reduces or prevents infection by the virus, bacterium or protozoon in a subject, such as a human or lower animal.

As used herein, the term "wound" includes any injury to any portion of the body of a subject including, but not limited to, acute conditions such as thermal burns, chemical burns, radiation burns, burns caused by excess exposure to ultraviolet radiation such as sunburn, damage to bodily tissues such as the perineum as a result of labor and childbirth, including injuries sustained during medical procedures such as episiotomies, trauma-induced injuries including cuts, those injuries sustained in automobile and other mechanical accidents, and those caused by bullets, knives and other weapons, and post-surgical injuries, as well as chronic conditions such as pressure sores, bedsores, conditions related to diabetes and poor circulation, and all types of acne, etc.

Anti-heparanase antibodies, raised against the recombinant enzyme, would be useful for immunodetection and diagnosis of micrometastases, autoimmune lesions and renal failure in biopsy specimens, plasma samples, and body fluids. Such antibodies may also serve as neutralizing agents for heparanase activity.

The genomic heparanase sequences described herein can be used to construct knock-in and knock-out constructs. Such constructs include a fragment of 10-20 Kb of a heparanase locus and a negative and a positive selection markers and can be used to provide heparanase knock-in and knock-out animal models by methods known to the skilled artisan. Such animal models can be used for studying the function of heparanase in developmental processes, and in normal as well as pathological processes. They can also serve as an experimental model for testing drugs and gene therapy protocols. The complementary heparanase sequence (cDNA) can be used to derive transgenic animals, overexpressing heparanase for same. Alternatively, if cloned in the antisense orientation, the complementary heparanase sequence (cDNA) can be used to derive transgenic animals under-expressing heparanase for same.

The heparanase promoter sequences described herein and other cis regulatory elements linked to the heparanase locus can be used to regulated the expression of genes. For example, these promoters can be used to direct the expression of a cytotoxic protein, such as TNF, in tumor cells. It will be appreciated that heparanase itself is abnormally expressed under the control of its own promoter and other cis acting elements in a variety of tumors, and its expression is correlated with metastasis. It is also abnormally highly expressed in inflammatory cells. The introns of the heparanase gene can be used for the same purpose, as it is known that introns, especially upstream introns include cis acting element which affect expression. A heparanase promoter fused to a reporter protein can be used to study/monitor its activity.

The polynucleotide sequences described herein can also be used to provide DNA vaccines which will elicit in vivo anti heparanase antibodies. Such vaccines can therefore be used to combat inflammatory and cancer.

Antisense oligonucleotides derived according to the heparanase sequences described herein, especially such oligonucleotides supplemented with ribozyme activity, can be used to modulate heparanase expression. Such oligonucleotides can be from the coding region, from the introns or promoter specific. Antisense heparanase nucleic acid constructs can similarly function, as well known in the art.

The heparanase sequences described herein can be used to study the catalytic mechanism of heparanase. Carefully selected site directed mutagenesis can be employed to provide modified heparanase proteins

having modified characteristics in terms of, for example, substrate specificity, sensitivity to inhibitors, etc.

While studying heparanase expression in a variety of cell types alternatively spliced transcripts were identified. Such transcripts if found characteristic of certain pathological conditions can be used as markers for such conditions. Such transcripts are expected to direct the synthesis of heparanases with altered functions.

Additional objects, advantages, and novel features of the present invention will become apparent to one ordinarily skilled in the art upon examination of the following examples, which are not intended to be limiting. Additionally, each of the various embodiments and aspects of the present invention as delineated hereinabove and as claimed in the claims section below finds experimental support in the following examples.

EXAMPLES

Generally, the nomenclature used herein and the laboratory procedures in recombinant DNA technology described below are those well known and commonly employed in the art. Standard techniques are used for cloning, DNA and RNA isolation, amplification and purification. Generally enzymatic reactions involving DNA ligase, DNA polymerase, restriction endonucleases and the like are performed according to the manufacturers' specifications. These techniques and various other techniques are generally performed according to Sambrook et al., Molecular Cloning--A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y. (1989), which is incorporated herein by reference. Other general references are provided throughout this document. The procedures therein are believed to be well known in the art and are provided for the convenience of the reader. All the information contained therein is incorporated herein by reference.

The following protocols and experimental details are referenced in the Examples that follow:

Purification and characterization of heparanase from a human hepatoma cell line and human placenta: A human hepatoma cell line (Sk-hep-1) was chosen as a source for purification of a human tumor-derived heparanase. Purification was essentially as described in U.S. Pat. No. 5,362,641 to Fuks, which is incorporated by reference as if fully set forth

herein. Briefly, 500 liter, 5×10^{11} cells were grown in suspension and the heparanase enzyme was purified about 240,000 fold by applying the following steps: (i) cation exchange (CM-Sephadex) chromatography performed at pH 6.0, 0.3-1.4 M NaCl gradient; (ii) cation exchange (CM-Sephadex) chromatography performed at pH 7.4 in the presence of 0.1% CHAPS, 0.3-1.1 M NaCl gradient; (iii) heparin-Sepharose chromatography performed at pH 7.4 in the presence of 0.1% CHAPS, 0.35-1.1 M NaCl gradient; (iv) ConA-Sepharose chromatography performed at pH 6.0 in buffer containing 0.1 % CHAPS and 1 M NaCl, elution with 0.25 M α -methyl mannoside; and (v) HPLC cation exchange (Mono-S) chromatography performed at pH 7.4 in the presence of 0.1 % CHAPS, 0.25-1 M NaCl gradient.

Active fractions were pooled, precipitated with TCA and the precipitate subjected to SDS polyacrylamide gel electrophoresis and/or tryptic digestion and reverse phase HPLC. Tryptic peptides of the purified protein were separated by reverse phase HPLC (C8 column) and homogeneous peaks were subjected to amino acid sequence analysis.

The purified enzyme was applied to reverse phase HPLC and subjected to N-terminal amino acid sequencing using the amino acid sequencer (Applied Biosystems).

Cells: Cultures of bovine corneal endothelial cells (BCECs) were established from steer eyes as previously described (19, 38). Stock cultures were maintained in DMEM (1 g glucose/liter) supplemented with 10 % newborn calf serum and 5 % FCS. bFGF (1 ng/ml) was added every other day during the phase of active cell growth (13, 14).

Preparation of dishes coated with ECM: BCECs (second to fifth passage) were plated into 4-well plates at an initial density of 2×10^5 cells/ml, and cultured in sulfate-free Fisher medium plus 5 % dextran T-40 for 12 days. $\text{Na}_2^{35}\text{SO}_4$ (25 $\mu\text{Ci/ml}$) was added on day 1 and 5 after seeding and the cultures were incubated with the label without medium change. The subendothelial ECM was exposed by dissolving (5 min., room temperature) the cell layer with PBS containing 0.5 % Triton X-100 and 20 mM NH_4OH , followed by four washes with PBS. The ECM remained intact, free of cellular debris and firmly attached to the entire area of the tissue culture dish (19, 22).

To prepare soluble sulfate labeled proteoglycans (peak I material), the ECM was digested with trypsin (25 $\mu\text{g/ml}$, 6 h, 37 °C), the digest was concentrated by reverse dialysis and the concentrated material was applied

onto a Sepharose 6B gel filtration column. The resulting high molecular weight material ($K_{av} < 0.2$, peak I) was collected. More than 80 % of the labeled material was shown to be composed of heparan sulfate proteoglycans (11, 39).

5 **Heparanase activity:** Cells (1×10^6 /35-mm dish), cell lysates or conditioned media were incubated on top of ^{35}S -labeled ECM (18 h, 37°C) in the presence of 20 mM phosphate buffer (pH 6.2). Cell lysates and conditioned media were also incubated with sulfate labeled peak I material (10-20 μl). The incubation medium was collected, centrifuged ($18,000 \times g$,
10 4 $^\circ\text{C}$, 3 min.), and sulfate labeled material analyzed by gel filtration on a Sepharose CL-6B column (0.9 x 30 cm). Fractions (0.2 ml) were eluted with PBS at a flow rate of 5 ml/h and counted for radioactivity using Bio-fluor scintillation fluid. The excluded volume (V_o) was marked by blue dextran and the total included volume (V_t) by phenol red. The latter was
15 shown to comigrate with free sulfate (7, 11, 23). Degradation fragments of HS side chains were eluted from Sepharose 6B at $0.5 < K_{av} < 0.8$ (peak II) (7, 11, 23). A nearly intact HSPG released from ECM by trypsin - and, to a lower extent, during incubation with PBS alone - was eluted next to V_o ($K_{av} < 0.2$, peak I). Recoveries of labeled material applied on the columns
20 ranged from 85 to 95 % in different experiments (11). Each experiment was performed at least three times and the variation of elution positions (K_{av} values) did not exceed ± 15 %.

Cloning of hpa cDNA: cDNA clones 257548 and 260138 were obtained from the I.M.A.G.E Consortium (2130 Memorial Parkway SW,
25 Huntsville, AL 35801). The cDNAs were originally cloned in *EcoRI* and *NotI* cloning sites in the plasmid vector pT3T7D-Pac. Although these clones are reported to be somewhat different, DNA sequencing demonstrated that these clones are identical to one another. Marathon RACE (rapid amplification of cDNA ends) human placenta (poly-A) cDNA
30 composite was a gift of Prof. Yossi Shiloh of Tel Aviv University. This composite is vector free, as it includes reverse transcribed cDNA fragments to which double, partially single stranded adapters are attached on both sides. The construction of the specific composite employed is described in reference 39a.

35 Amplification of hp3 PCR fragment was performed according to the protocol provided by Clontech laboratories. The template used for amplification was a sample taken from the above composite. The primers used for amplification were:

First step: 5'-primer: AP1: 5'-CCATCCTAATACGACTCACT ATAGGGC-3', SEQ ID NO:1; 3'-primer: HPL229: 5'-GTAGTGATGCCA TGTAAGTGAATC-3', SEQ ID NO:2.

Second step: nested 5'-primer: AP2: 5'-ACTCACTATAGGGCTCG AGCGGC-3', SEQ ID NO:3; nested 3'- primer: HPL171: 5'-GCATCTTAGCCGTCTTCTTCG-3', SEQ ID NO:4. The HPL229 and HPL171 were selected according to the sequence of the EST clones. They include nucleotides 933-956 and 876-897 of SEQ ID NO:9, respectively.

PCR program was 94 °C - 4 min., followed by 30 cycles of 94 °C - 40 sec., 62 °C - 1 min., 72 °C - 2.5 min. Amplification was performed with Expand High Fidelity (Boehringer Mannheim). The resulting ca. 900 bp hp3 PCR product was digested with *Bfr*I and *Pvu*II. Clone 257548 (*phpa*1) was digested with *Eco*RI, followed by end filling and was then further digested with *Bfr*I. Thereafter the *Pvu*II - *Bfr*I fragment of the hp3 PCR product was cloned into the blunt end - *Bfr*I end of clone *phpa*1 which resulted in having the entire cDNA cloned in pT3T7-pac vector, designated *phpa*2.

RT-PCR: RNA was prepared using TRI-Reagent (Molecular research center Inc.) according to the manufacturer instructions. 1.25 µg were taken for reverse transcription reaction using MuMLV Reverse transcriptase (Gibco BRL) and Oligo (dT)₁₅ primer, SEQ ID NO:5, (Promega). Amplification of the resultant first strand cDNA was performed with *Taq* polymerase (Promega). The following primers were used:

HPU-355: 5'-TTCGATCCCAAGAAGGAATCAAC-3', SEQ ID NO:6, nucleotides 372-394 in SEQ ID NOs:9 or 11.
HPL-229: 5'-GTAGTGATGCCATGTAAGTGAATC-3', SEQ ID NO:7, nucleotides 933-956 in SEQ ID NOs:9 or 11.

PCR program: 94 °C - 4 min., followed by 30 cycles of 94 °C - 40 sec., 62 °C - 1 min., 72 °C - 1 min.

Alternatively, total RNA was prepared from cell cultures using Tri-reagent (Molecular Research Center, Inc.) according to the manufacturer recommendation. Poly A+ RNA was isolated from total RNA using mRNA separator (Clontech). Reverse transcription was performed with total RNA using Superscript II (GibcoBRL). PCR was performed with Expand high fidelity (Boehringer Mannheim). Primers used for amplification were as follows:

Hpu-685, 5'-GAGCAGCCAGGTGAGCCCAAGAT-3', SEQ ID NO:24

Hpu-355, 5'-TTCGATCCCAAGAAGGAATCAAC-3', SEQ ID NO:25

Hpu 565, 5'-AGCTCTGTAGATGTGCTATACAC-3', SEQ ID NO:26

Hpl 967, 5'-TCAGATGCAAGCAGCAACTTTGGC-3', SEQ ID NO:27

Hpl 171, 5'-GCATCTTAGCCGTCTTTCTTCG-3', SEQ ID NO:28

5 Hpl 229, 5'-GTAGTGATGCCATGTAAGTGAATC-3', SEQ ID NO:29

PCR reaction was performed as follows: 94 °C 3 minutes, followed by 32 cycles of 94 °C 40 seconds, 64 °C 1 minute, 72 °C 3 minutes, and one cycle 72 °C, 7 minutes.

Expression of recombinant heparanase in insect cells: Cells, High Five and Sf21 insect cell lines were maintained as monolayer cultures in SF900II-SFM medium (GibcoBRL).

Recombinant Baculovirus: Recombinant virus containing the *hpa* gene was constructed using the Bac to Bac system (GibcoBRL). The transfer vector pFastBac was digested with *Sa*I and *Not*I and ligated with a 1.7 kb fragment of *phpa2* digested with *Xho*I and *Not*I. The resulting plasmid was designated pFast*hpa2*. An identical plasmid designated pFast*hpa4* was prepared as a duplicate and both independently served for further experimentations. Recombinant bacmid was generated according to the instructions of the manufacturer with pFast*hpa2*, pFast*hpa4* and with pFastBac. The latter served as a negative control. Recombinant bacmid DNAs were transfected into Sf21 insect cells. Five days after transfection recombinant viruses were harvested and used to infect High Five insect cells, 3 x 10⁶ cells in T-25 flasks. Cells were harvested 2 - 3 days after infection. 4 x 10⁶ cells were centrifuged and resuspended in a reaction buffer containing 20 mM phosphate citrate buffer, 50 mM NaCl. Cells underwent three cycles of freeze and thaw and lysates were stored at -80 °C. Conditioned medium was stored at 4 °C.

Partial purification of recombinant heparanase: Partial purification of recombinant heparanase was performed by heparin-Sepharose column chromatography followed by Superdex 75 column gel filtration. Culture medium (150 ml) of Sf21 cells infected with pF*hpa4* virus was subjected to heparin-Sepharose chromatography. Elution of 1 ml fractions was performed with 0.35 - 2 M NaCl gradient in presence of 0.1 % CHAPS and 1 mM DTT in 10 mM sodium acetate buffer, pH 5.0. A 25 µl sample of each fraction was tested for heparanase activity. Heparanase activity was eluted at the range of 0.65 - 1.1 M NaCl (fractions 18-26, Figure 10a). 5 µl of each fraction was subjected to 15 % SDS-polyacrylamide gel electrophoresis followed by silver nitrate staining.

Active fractions eluted from heparin-Sepharose (Figure 10a) were pooled and concentrated (x 6) on YM3 cut-off membrane. 0.5 ml of the concentrated material was applied onto 30 ml Superdex 75 FPLC column equilibrated with 10 mM sodium acetate buffer, pH 5.0, containing 0.8 M NaCl, 1 mM DTT and 0.1 % CHAPS. Fractions (0.56 ml) were collected at a flow rate of 0.75 ml/min. Aliquots of each fraction were tested for heparanase activity and were subjected to SDS-polyacrylamide gel electrophoresis followed by silver nitrate staining (Figure 11b).

PCR amplification of genomic DNA: 94 °C 3 minutes, followed by 32 cycles of 94 °C 45 seconds, 64 °C 1 minute, 68 °C 5 minutes, and one cycle at 72 °C, 7 minutes. Primers used for amplification of genomic DNA included:

GHpu-L3 5'-AGGCACCCTAGAGATGTTCCAG-3', SEQ ID NO:30

GHpl-L6 5'-GAAGATTTCTGTTTCCATGACGTG-3', SEQ ID NO:31.

Screening of genomic libraries: A human genomic library in Lambda phage EMBLE3 SP6/T7 (Clontech, Paulo Alto, CA) was screened. 5 x 10⁵ plaques were plated at 5 x 10⁴ pfu/plate on NZCYM agar/top agarose plates. Phages were absorbed on nylon membranes in duplicates (Qiagen). Hybridization was performed at 65 °C in 5 x SSC, 5 x Denhart's, 10 % dextran sulfate, 100 µg/ml Salmon sperm, ³²p labeled probe (10⁶ cpm/ml). A 1.6 kb fragment, containing the entire *hpa* cDNA was labeled by random priming (Boehringer Mannheim). Following hybridization membranes were washed once with 2 x SSC, 0.1 % SDS at 65 °C for 20 minutes, and twice with 0.2 x SSC, 0.1 % SDS at 65 °C for 15 minutes. Hybridizing plaques were picked, and plated at 100 pfu/plate. Hybridization was performed as above and single isolated positive plaques were picked.

Phage DNA was extracted using a Lambda DNA extraction kit (Qiagen). DNA was digested with *Xho*I and *Eco*RI, separated on 0.7 % agarose gel and transferred to nylon membrane Hybond N+ (Amersham). Hybridization and washes were performed as above.

cDNA Sequence analysis: Sequence determinations were performed with vector specific and gene specific primers, using an automated DNA sequencer (Applied Biosystems, model 373A). Each nucleotide was read from at least two independent primers.

Genomic sequence analysis: Large-scale sequencing was performed by Commonwealth Biotechnology Incorporation.

Isolation of mouse *hpa*: Mouse *hpa* cDNA was amplified from either Marathon ready cDNA library of mouse embryo or from mRNA isolated from mouse melanoma cell line BL6, using the Marathon RACE kit from Clontech. Both procedures were performed according to the manufacturer's recommendation.

Primers used for PCR amplification of mouse *hpa*:

Mhpl773 5'-CCACACTGAATGTAATACTGAAGTG-3', SEQ ID NO:32

MHpl736 5'-CGAAGCTCTGGAAGCTCGGCAAG-3', SEQ ID NO:33

MHpl83 5'-GCCAGCTGCAAAGGTGTTGGAC-3', SEQ ID NO:34

Mhpl152 5'-AACACCTGCCTCATCACGACTTC-3', SEQ ID NO:35

Mhpl114 5'-GCCAGGCTGGCGTCGATGGTGA-3', SEQ ID NO:36

MHpl103 5'-GTCGATGGTGATGGACAGGAAC-3', SEQ ID NO:37

Ap1 5'-GTAATACGACTCACTATAGGGC-3', SEQ ID NO:38 -

(Genome walker)

Ap2 5'-ACTATAGGGCACGCGTGGT-3', SEQ ID NO:39 -

(Genome walker)

Ap1 5'-CCATCCTAATACGACTCACTATAGGGC-3', SEQ ID NO:40 -

(Marathon RACE)

Ap2 5'-ACTCACTATAGGGCTCGAGCGGC-3', SEQ ID NO:41 -

(Marathon RACE)

Southern analysis of genomic DNA: Genomic DNA was extracted from animal or from human blood using Blood and cell culture DNA maxi kit (Qiagene). DNA was digested with *EcoRI*, separated by gel electrophoresis and transferred to a nylon membrane Hybond N+ (Amersham). Hybridization was performed at 68 °C in 6 x SSC, 1 % SDS, 5 x Denharts, 10 % dextran sulfate, 100 µg/ml salmon sperm DNA, and ³²p labeled probe. A 1.6 kb fragment, containing the entire *hpa* cDNA was used as a probe. Following hybridization, the membrane was washed with 3 x SSC, 0.1 % SDS, at 68 °C and exposed to X-ray film for 3 days. Membranes were then washed with 1 x SSC, 0.1 % SDS, at 68 °C and were reexposed for 5 days.

Construction of *hpa* promoter-GFP expression vector: Lambda DNA of phage L3, was digested with *SacI* and *BglII*, resulting in a 1712 bp fragment which contained the *hpa* promoter (877-2688 of SEQ ID NO:42). The pEGFP-1 plasmid (Clontech) was digested with *BglII* and *SacI* and ligated with the 1712 bp fragment of the *hpa* promoter sequence. The resulting plasmid was designated phpEGL. A second *hpa* promoter-GFP plasmid was constructed containing a shorter fragment of the *hpa* promoter

region: phpEGL was digested with *Hind*III, and the resulting 1095 bp fragment (nucleotides 1593-2688 of SEQ ID NO:42) was ligated with *Hind*III digested pEGFP-1. The resulting plasmid was designated phpEGS.

Computer analysis of sequences: Homology searches were performed using several computer servers, and various databases. Blast 2.0 service, at the NCBI server was used to screen the protein database swplus and DNA databases such as GenBank, EMBL, and the EST databases. Blast 2.0 search was performed using the basic search option of the NCBI server. Sequence analysis and alignments were done using the DNA sequence analysis software package developed by the Genetic Computer Group (GCG) at the university of Wisconsin. Alignments of two sequences were performed using Bestfit (gap creation penalty - 12, gap extension penalty - 4). Protein homology search was performed with the Smith-Waterman algorithm, using the Bioaccelerator platform developed by Compugene. The protein database swplus was searched using the following parameters: gapop: 10.0, gapext: 0.5, matrix: blosum62. Blocks homology was performed using the Blocks WWW server developed at Fred Hutchinson Cancer Research Center in Seattle, Washington, USA. Secondary structure prediction was performed using the PHD server – Profile network Prediction Heidelberg. Fold recognition (threading) was performed using the UCLA-DOE structure prediction server. The method used for prediction was gonnet+predss. Alignment of three sequences was performed using the pileup application (gap creation penalty - 5, gap extension penalty - 1). Promoter analysis was performed using TSSW and TSSG programs (BCM Search Launcher Human Genome Center, Baylor College of Medicine, Houston TX).

EXAMPLE 1

Cloning of human hpa cDNA

Purified fraction of heparanase isolated from human hepatoma cells (SK-hep-1) was subjected to tryptic digestion and microsequencing. EST (Expressed Sequence Tag) databases were screened for homology to the back translated DNA sequences corresponding to the obtained peptides. Two EST sequences (accession Nos. N41349 and N45367) contained a DNA sequence encoding the peptide YGPDVGQPR (SEQ ID NO:8). These two sequences were derived from clones 257548 and 260138 (I.M.A.G.E Consortium) prepared from 8 to 9 weeks placenta cDNA library (Soares). Both clones which were found to be identical contained an insert

of 1020 bp which included an open reading frame (ORF) of 973 bp followed by a 3' untranslated region of 27 bp and a Poly A tail. No translation start site (AUG) was identified at the 5' end of these clones.

Cloning of the missing 5' end was performed by PCR amplification of DNA from a placenta Marathon RACE cDNA composite. A 900 bp fragment (designated hp3), partially overlapping with the identified 3' encoding EST clones was obtained.

The joined cDNA fragment, 1721 bp long (SEQ ID NO:9), contained an open reading frame which encodes, as shown in Figure 1 and SEQ ID NO:11, a polypeptide of 543 amino acids (SEQ ID NO:10) with a calculated molecular weight of 61,192 daltons. The 3' end of the partial cDNA inserts contained in clones 257548 and 260138 started at nucleotide G⁷²¹ of SEQ ID NO:9 and Figure 1.

As further shown in Figure 1, there was a single sequence discrepancy between the EST clones and the PCR amplified sequence, which led to an amino acid substitution from Tyr²⁴⁶ in the EST to Phe²⁴⁶ in the amplified cDNA. The nucleotide sequence of the PCR amplified cDNA fragment was verified from two independent amplification products. The new gene was designated *hpa*.

As stated above, the 3' end of the partial cDNA inserts contained in EST clones 257548 and 260138 started at nucleotide 721 of *hpa* (SEQ ID NO:9). The ability of the *hpa* cDNA to form stable secondary structures, such as stem and loop structures involving nucleotide stretches in the vicinity of position 721 was investigated using computer modeling. It was found that stable stem and loop structures are likely to be formed involving nucleotides 698-724 (SEQ ID NO:9). In addition, a high GC content, up to 70 %, characterizes the 5' end region of the *hpa* gene, as compared to about only 40 % in the 3' region. These findings may explain the immature termination and therefore lack of 5' ends in the EST clones.

To examine the ability of the *hpa* gene product to catalyze degradation of heparan sulfate in an *in vitro* assay the entire open reading frame was expressed in insect cells, using the Baculovirus expression system. Extracts of cells, infected with virus containing the *hpa* gene, demonstrated a high level of heparan sulfate degradation activity, while cells infected with a similar construct containing no *hpa* gene had no such activity, nor did non-infected cells. These results are further demonstrated in the following Examples.

EXAMPLE 2

Degradation of soluble ECM-derived HSPG

Monolayer cultures of High Five cells were infected (72 h, 28 °C) with recombinant Baculovirus containing the pFast hpa plasmid or with control virus containing an insert free plasmid. The cells were harvested and lysed in heparanase reaction buffer by three cycles of freezing and thawing. The cell lysates were then incubated (18 h, 37 °C) with sulfate labeled, ECM-derived HSPG (peak I), followed by gel filtration analysis (Sepharose 6B) of the reaction mixture.

As shown in Figure 2, the substrate alone included almost entirely high molecular weight (M_r) material eluted next to V_0 (peak I, fractions 5-20, $K_{av} < 0.35$). A similar elution pattern was obtained when the HSPG substrate was incubated with lysates of cells that were infected with control virus. In contrast, incubation of the HSPG substrate with lysates of cells infected with the hpa containing virus resulted in a complete conversion of the high M_r substrate into low M_r labeled degradation fragments (peak II, fractions 22-35, $0.5 < K_{av} < 0.75$).

Fragments eluted in peak II were shown to be degradation products of heparan sulfate, as they were (i) 5- to 6-fold smaller than intact heparan sulfate side chains (K_{av} approx. 0.33) released from ECM by treatment with either alkaline borohydride or papain; and (ii) resistant to further digestion with papain or chondroitinase ABC, and susceptible to deamination by nitrous acid (6, 11). Similar results (not shown) were obtained with Sf21 cells. Again, heparanase activity was detected in cells infected with the hpa containing virus (pF hpa), but not with control virus (pF). This result was obtained with two independently generated recombinant viruses. Lysates of control not infected High Five cells failed to degrade the HSPG substrate.

In subsequent experiments, the labeled HSPG substrate was incubated with medium conditioned by infected High Five or Sf21 cells.

As shown in Figures 3a-b, heparanase activity, reflected by the conversion of the high M_r peak I substrate into the low M_r peak II which represents HS degradation fragments, was found in the culture medium of cells infected with the pF $hpa2$ or pF $hpa4$ viruses, but not with the control pF1 or pF2 viruses. No heparanase activity was detected in the culture medium of control non-infected High Five or Sf21 cells.

The medium of cells infected with the pF $hpa4$ virus was passed through a 50 kDa cut off membrane to obtain a crude estimation of the

molecular weight of the recombinant heparanase enzyme. As demonstrated in Figure 4, all the enzymatic activity was retained in the upper compartment and there was no activity in the flow through (<50 kDa) material. This result is consistent with the expected molecular weight of the *hpa* gene product.

In order to further characterize the *hpa* product the inhibitory effect of heparin, a potent inhibitor of heparanase mediated HS degradation (40) was examined.

As demonstrated in Figures 5a-b, conversion of the peak I substrate into peak II HS degradation fragments was completely abolished in the presence of heparin.

Altogether, these results indicate that the heparanase enzyme is expressed in an active form by insect cells infected with Baculovirus containing the newly identified human *hpa* gene.

EXAMPLE 3

Degradation of HSPG in intact ECM

Next, the ability of intact infected insect cells to degrade HS in intact, naturally produced ECM was investigated. For this purpose, High Five or Sf21 cells were seeded on metabolically sulfate labeled ECM followed by infection (48 h, 28 °C) with either the pF*hpa*4 or control pF2 viruses. The pH of the medium was then adjusted to pH 6.2-6.4 and the cells further incubated with the labeled ECM for another 48 h at 28 °C or 24 h at 37 °C. Sulfate labeled material released into the incubation medium was analyzed by gel filtration on Sepharose 6B.

As shown in Figures 6a-b and 7a-b, incubation of the ECM with cells infected with the control pF2 virus resulted in a constant release of labeled material that consisted almost entirely (>90%) of high Mr fragments (peak I) eluted with or next to V_0 . It was previously shown that a proteolytic activity residing in the ECM itself and/or expressed by cells is responsible for release of the high Mr material (6). This nearly intact HSPG provides a soluble substrate for subsequent degradation by heparanase, as also indicated by the relatively large amount of peak I material accumulating when the heparanase enzyme is inhibited by heparin (6, 7, 12, Figure 9). On the other hand, incubation of the labeled ECM with cells infected with the pF*hpa*4 virus resulted in release of 60-70% of the ECM-associated radioactivity in the form of low Mr sulfate-labeled fragments (peak II, $0.5 < K_{av} < 0.75$), regardless of whether the infected cells were incubated with

the ECM at 28 °C or 37 °C. Control intact non-infected Sf21 or High Five cells failed to degrade the ECM HS side chains.

In subsequent experiments, as demonstrated in Figures 8a-b, High Five and Sf21 cells were infected (96 h, 28 °C) with pFhpa4 or control pF1 viruses and the culture medium incubated with sulfate-labeled ECM. Low Mr HS degradation fragments were released from the ECM only upon incubation with medium conditioned by pFhpa4 infected cells. As shown in Figure 9, production of these fragments was abolished in the presence of heparin. No heparanase activity was detected in the culture medium of control, non-infected cells. These results indicate that the heparanase enzyme expressed by cells infected with the pFhpa4 virus is capable of degrading HS when complexed to other macromolecular constituents (i.e. fibronectin, laminin, collagen) of a naturally produced intact ECM, in a manner similar to that reported for highly metastatic tumor cells or activated cells of the immune system (6, 7).

EXAMPLE 4

Purification of recombinant human heparanase

The recombinant heparanase was partially purified from medium of pFhpa4 infected Sf21 cells by Heparin-Sepharose chromatography (Figure 10a) followed by gel filtration of the pooled active fractions over an FPLC Superdex 75 column (Figure 11a). A ~ 63 kDa protein was observed, whose quantity, as was detected by silver stained SDS-polyacrylamide gel electrophoresis, correlated with heparanase activity in the relevant column fractions (Figures 10b and 11b, respectively). This protein was not detected in the culture medium of cells infected with the control pF1 virus and was subjected to a similar fractionation on heparin-Sepharose (not shown).

EXAMPLE 5

Expression of the human hpa cDNA in various cell types, organs and tissues

Referring now to Figures 12a-e, RT-PCR was applied to evaluate the expression of the *hpa* gene by various cell types and tissues. For this purpose, total RNA was reverse transcribed and amplified. The expected 585 bp long cDNA was clearly demonstrated in human kidney, placenta (8 and 11 weeks) and mole tissues, as well as in freshly isolated and short termed (1.5-48 h) cultured human placental cytotrophoblastic cells (Figure 12a), all known to express a high heparanase activity (41). The *hpa*

transcript was also expressed by normal human neutrophils (Figure 12b). In contrast, there was no detectable expression of the *hpa* mRNA in embryonic human muscle tissue, thymus, heart and adrenal (Figure 12b). The *hpa* gene was expressed by several, but not all, human bladder carcinoma cell lines (Figure 12c), SK hepatoma (SK-hep-1), ovarian carcinoma (OV 1063), breast carcinoma (435, 231), melanoma and megakaryocytic (DAMI, CHRF) human cell lines (Figures 12d-e).

The above described expression pattern of the *hpa* transcript was determined to be in a very good correlation with heparanase activity levels determined in various tissues and cell types (not shown).

EXAMPLE 6

Isolation of an extended 5' end of hpa cDNA from human SK-hep1 cell line

The 5' end of *hpa* cDNA was isolated from human SK-hep1 cell line by PCR amplification using the Marathon RACE (rapid amplification of cDNA ends) kit (Clontech). Total RNA was prepared from SK-hep1 cells using the TRI-Reagent (Molecular research center Inc.) according to the manufacturer instructions. Poly A+ RNA was isolated using the mRNA separator kit (Clontech).

The Marathon RACE SK-hep1 cDNA composite was constructed according to the manufacturer recommendations. First round of amplification was performed using an adaptor specific primer AP1: 5'-CCATCCTAATACG ACTCACTATAGGGC-3', SEQ ID NO:1, and a *hpa* specific antisense primer hpl-629: 5'-CCCCAGGAGCAGCAGCATCAG-3', SEQ ID NO:17, corresponding to nucleotides 119-99 of SEQ ID NO:9. The resulting PCR product was subjected to a second round of amplification using an adaptor specific nested primer AP2: 5'-ACTCACTATAGGGCTCGAGCGGC-3', SEQ ID NO:3, and a *hpa* specific antisense nested primer hpl-666 5'-AGGCTTCGAGCGCAGCAGCAT-3', SEQ ID NO:18, corresponding to nucleotides 83-63 of SEQ ID NO:9. The PCR program was as follows: a hot start of 94 °C for 1 minute, followed by 30 cycles of 90 °C - 30 seconds, 68 °C - 4 minutes. The resulting 300 bp DNA fragment was extracted from an agarose gel and cloned into the vector pGEM-T Easy (Promega). The resulting recombinant plasmid was designated pHPSK1.

The nucleotide sequence of the pHPSK1 insert was determined and it was found to contain 62 nucleotides of the 5' end of the placenta *hpa* cDNA

(SEQ ID NO:9) and additional 178 nucleotides upstream, the first 178 nucleotides of SEQ ID NOs:13 and 15.

A single nucleotide discrepancy was identified between the SK-hep1 cDNA and the placenta cDNA. The "T" derivative at position 9 of the placenta cDNA (SEQ ID NO:9), is replaced by a "C" derivative at the corresponding position 187 of the SK-hep1 cDNA (SEQ ID NO:13).

The discrepancy is likely to be due to a mutation at the 5' end of the placenta cDNA clone as confirmed by sequence analysis of several additional cDNA clones isolated from placenta, which like the SK-hep1 cDNA contained C at position 9 of SEQ ID NO:9.

The 5' extended sequence of the SK-hep1 *hpa* cDNA was assembled with the sequence of the *hpa* cDNA isolated from human placenta (SEQ ID NO:9). The assembled sequence contained an open reading frame which encodes, as shown in SEQ ID NOs:14 and 15, a polypeptide of 592 amino acids with a calculated molecular weight of 66,407 daltons. The open reading frame is flanked by 93 bp 5' untranslated region (UTR).

EXAMPLE 7

Isolation of the upstream genomic region of the hpa gene

The upstream region of the *hpa* gene was isolated using the Genome Walker kit (Clontech) according to the manufacturer recommendations. The kit includes five human genomic DNA samples each digested with a different restriction endonuclease creating blunt ends: *EcoRV*, *ScaI*, *DraI*, *PvuII* and *SspI*.

The blunt ended DNA fragments are ligated to partially single stranded adaptors. The Genomic DNA samples were subjected to PCR amplification using the adaptor specific primer and a gene specific primer. Amplification was performed with Expand High Fidelity (Boehringer Mannheim).

A first round of amplification was performed using the ap1 primer: 5'-G TAATACGACTCACTATAGGGC-3', SEQ ID NO:19, and the *hpa* specific antisense primer hpl-666: 5'-AGGCTTCGAGCGCAGCAGCAT-3', SEQ ID NO:18, corresponding to nucleotides 83 - 63 of SEQ ID NO:9. The PCR program was as follows: a hot start of 94 °C - 3 minutes, followed by 36 cycles of 94 °C - 40 seconds, 67 °C - 4 minutes.

The PCR products of the first amplification were diluted 1:50. One µl of the diluted sample was used as a template for a second amplification using a nested adaptor specific primer ap2: 5'-

ACTATAGGGGCACGCGTGGT-3', SEQ ID NO:20, and a *hpa* specific antisense primer hpl-690, 5'-CTTGGGCTCACC TGGCTGCTC-3', SEQ ID NO:21, corresponding to nucleotides 62-42 of SEQ ID NO:9. The resulting amplification products were analyzed using agarose gel electrophoresis.

5 Five different PCR products were obtained from the five amplification reactions. A DNA fragment of approximately 750 bp which was obtained from the *SspI* digested DNA sample was gel extracted. The purified fragment was ligated into the plasmid vector pGEM-T Easy (Promega). The resulting recombinant plasmid was designated pGHP6905 and the

10 nucleotide sequence of the *hpa* insert was determined.

A partial sequence of 594 nucleotides is shown in SEQ ID NO:16. The last nucleotide in SEQ ID NO:13 corresponds to nucleotide 93 in SEQ ID:13. The DNA sequence in SEQ ID NO:16 contains the 5' region of the *hpa* cDNA and 501 nucleotides of the genomic upstream region which are

15 predicted to contain the promoter region of the *hpa* gene.

EXAMPLE 8

Expression of the 592 amino acids HPA polypeptide in a human 293 cell line

20 The 592 amino acids open reading frame (SEQ ID NOs:13 and 15) was constructed by ligation of the 110 bp corresponding to the 5' end of the SK-hep1 *hpa* cDNA with the placenta cDNA. More specifically the Marathon RACE - PCR amplification product of the placenta *hpa* DNA was digested with *SacI* and an approximately 1 kb fragment was ligated into a

25 *SacI*-digested pGHP6905 plasmid. The resulting plasmid was digested with *EatI* and *AatII*. The *EatI* sticky ends were blunted and an approximately 280 bp *EatI*/blunt-*AatII* fragment was isolated. This fragment was ligated with pFast*hpa* digested with *EcoRI* which was blunt ended using Klenow fragment and further digested with *AatII*. The resulting plasmid contained a

30 1827 bp insert which includes an open reading frame of 1776 bp, 31 bp of 3' UTR and 21 bp of 5' UTR. This plasmid was designated pFast*Lhpa*.

A mammalian expression vector was constructed to drive the expression of the 592 amino acids heparanase polypeptide in human cells. The *hpa* cDNA was excised from pFast*Lhpa* with *BssHII* and *NotI*. The

35 resulting 1850 bp *BssHII*-*NotI* fragment was ligated to a mammalian expression vector pSI (Promega) digested with *MluI* and *NotI*. The resulting recombinant plasmid, pSI*hpa*Met2 was transfected into a human 293 embryonic kidney cell line.

Transient expression of the 592 amino-acids heparanase was examined by western blot analysis and the enzymatic activity was tested using the gel shift assay. Both these procedures are described in length in U.S. Pat. application No. 09/071,739, filed May 1, 1998, which is incorporated by reference as if fully set forth herein. Cells were harvested 3 days following transfection. Harvested cells were re-suspended in lysis buffer containing 150 mM NaCl, 50 mM Tris pH 7.5, 1% Triton X-100, 1 mM PMSF and protease inhibitor cocktail (Boehringer Mannheim). 40 µg protein extract samples were used for separation on a SDS-PAGE. Proteins were transferred onto a PVDF Hybond-P membrane (Amersham). The membrane was incubated with an affinity purified polyclonal anti heparanase antibody, as described in U.S. Pat. application No. 09/071,739. A major band of approximately 50 kDa was observed in the transfected cells as well as a minor band of approximately 65 kDa. A similar pattern was observed in extracts of cells transfected with the pShpa as demonstrated in U.S. Pat. application No. 09/071,739. These two bands probably represent two forms of the recombinant heparanase protein produced by the transfected cells. The 65 kDa protein probably represents a heparanase precursor, while the 50 kDa protein is suggested herein to be the processed or mature form.

The catalytic activity of the recombinant protein expressed in the pShpaMet2 transfected cells was tested by gel shift assay. Cell extracts of transfected and of mock transfected cells were incubated overnight with heparin (6 µg in each reaction) at 37 °C, in the presence of 20 mM phosphate citrate buffer pH 5.4, 1 mM CaCl₂, 1 mM DTT and 50 mM NaCl. Reaction mixtures were then separated on a 10 % polyacrylamide gel. The catalytic activity of the recombinant heparanase was clearly demonstrated by a faster migration of the heparin molecules incubated with the transfected cell extract as compared to the control. Faster migration indicates the disappearance of high molecular weight heparin molecules and the generation of low molecular weight degradation products.

EXAMPLE 9

Chromosomal localization of the hpa gene

Chromosomal mapping of the *hpa* gene was performed utilizing a panel of monochromosomal human/CHO and human/mouse somatic cell hybrids, obtained from the UK HGMP Resource Center (Cambridge, England).

40 ng of each of the somatic cell hybrid DNA samples were subjected to PCR amplification using the *hpa* primers: hpu565 5'-AGCTCTGTAGATGTGC TATACAC-3', SEQ ID NO:22, corresponding to nucleotides 564-586 of SEQ ID NO:9 and an antisense primer hpl171 5'-GCATCTTAGCCGTCTTTCTTCG-3', SEQ ID NO:23, corresponding to nucleotides 897-876 of SEQ ID NO:9.

The PCR program was as follows: a hot start of 94 °C – 3 minutes, followed by 7 cycles of 94 °C – 45 seconds, 66 °C – 1 minute, 68 °C – 5 minutes, followed by 30 cycles of 94 °C – 45 seconds, 62 °C – 1 minute, 68 °C – 5 minutes, and a 10 minutes final extension at 72 °C.

The reactions were performed with Expand long PCR (Boehringer Mannheim). The resulting amplification products were analyzed using agarose gel electrophoresis. As demonstrated in Figure 14, a single band of approximately 2.8 Kb was obtained from chromosome 4, as well as from the control human genomic DNA. A 2.8 kb amplification product is expected based on amplification of the genomic *hpa* clone (data not shown). No amplification products were obtained neither in the control DNA samples of hamster and mouse nor in somatic hybrids of other human chromosome.

EXAMPLE 10

Human genomic clone encoding heparanase

Five plaques were isolated following screening of a human genomic library and were designated L3-1, L5-1, L8-1, L10-1 and L6-1. The phage DNAs were analyzed by Southern hybridization and by PCR with *hpa* specific and vector specific primers. Southern analysis was performed with three fragments of *hpa* cDNA: a *PvuII-BamHI* fragment (nucleotides 32-450, SEQ ID NO:9), a *BamHI-NdeI* fragment (nucleotides 451-1102, SEQ ID NO:9) and an *NdeI-XhoI* fragment (nucleotides 1103-1721, SEQ ID NO:9).

Following Southern analysis, phages L3, L6, L8 were selected for further analysis. A scheme of the genomic region and the relative position of the three phage clones is depicted in Figure 15. A 2 kb DNA fragment containing the gap between phages L6 and L3 was PCR amplified from human genomic DNA with two gene specific primers GHpuL3 and GHplL6. The PCR product was cloned into the plasmid vector pGEM-T-easy (Promega).

Large scale DNA sequencing of the three Lambda clones and the amplified fragment was performed with Lambda purified DNA by primer walking. A nucleotide sequence of 44,898 bp was analyzed (Figure 16, SEQ ID NO:42). Comparison of the genomic sequence with that of *hpa* cDNA revealed 12 exons separated by 11 introns (Figures 15 and 16). The genomic organization of the *hpa* gene is depicted in Figure 15 (top). The sequence includes the coding region from the first ATG to the stop codon which spans 39,113 nucleotides, 2742 nucleotides upstream of the first ATG and 3043 nucleotides downstream of the stop codon. Splice site consensus sequences were identified at exon/intron junctions.

EXAMPLE 11

Alternative splicing

Several minor RT-PCR products were obtained from various cell types, following amplification with *hpa* specific primers. Each one found to contain a deletion of one or two exons. Some of these PCR products contain ORFs, which encode potential shorter proteins.

Table 1 below summarizes the alternative spliced products isolated from various cell lines.

Fragments of similar sizes were obtained following amplification with two cell lines, placenta and platelets.

Cell type	Nucleotides deleted	Exons deleted	ORF
Platelets	1047-1267	8, 9	+
Platelets	1154-1267	9	-
Platelets	289-435, 562-735	2, 4	-
Sk-hep1, platelets, Zr75	562-735	4	+
Sk-hep1 (hepatoma)	561-904	4, 5	-
Zr75 (breast carcinoma)	96-203	1 (partial)	+

EXAMPLE 12

Mouse and rat hpa

EST databases were screened for sequences homologous to the *hpa* gene. Three mouse EST's were identified (accession No. Aa177901, from mouse spleen, Aa067997 from mouse skin, Aa47943 from mouse embryo), assembled into a 824 bp cDNA fragment which contains a partial open reading frame (lacking a 5' end) of 629 bp and a 3' untranslated region of

195 bp (SEQ ID NO:12). As shown in Figure 13, the coding region is 80 % similar to the 3' end of the *hpa* cDNA sequence. These EST's are probably cDNA fragments of the mouse *hpa* homolog that encodes for the mouse heparanase.

5 Searching for consensus protein domains revealed an amino terminal homology between the heparanase and several precursor proteins such as Procollagen Alpha 1 precursor, Tyrosine-protein kinase-RYK, Fibulin-1, Insulin-like growth factor binding protein and several others. The amino terminus is highly hydrophobic and contains a potential trans-membrane domain. The homology to known signal peptide sequences suggests that it could function as a signal peptide for protein localization.

10 The amino acid sequence of human heparanase was used to search for homologous sequences in the DNA and protein databases. Several human EST's were identified, as well as mouse sequences highly homologous to human heparanase. The following mouse EST's were identified AA177901, AA674378, AA67997, AA047943, AA690179, AI122034, all sharing an identical sequence and correspond to amino acids 336-543 of the human heparanase sequence. The entire mouse heparanase cDNA was cloned, based on the nucleotide sequence of the mouse EST's.

15 PCR primers were designed and a Marathon RACE was performed using a Marathon cDNA library from 15 days mouse embryo (Clontech) and from BL6 mouse melanoma cell line. The mouse *hpa* homologous cDNA was isolated following several amplification steps. A 1.1 kb fragment was amplified from mouse embryo Marathon cDNA library. The first cycle of amplification was performed with primers mhpl773 and Ap1 and the second cycle with primers mhpl736 and AP2. A 1.1 kb fragment was then amplified from BL6 Marathon cDNA library. The first cycle of amplification was performed with the primers mhpl152 and Ap1, and the second with mhpl83 and AP2. The combined sequence was homologous to

20 nucleotides 157 - 1702 of the human *hpa* cDNA, which encode amino acids 33-543. The 5' end of the mouse *hpa* gene was isolated from a mouse genomic DNA library using the Genome Walker kit (Clontech). An 0.9 kb fragment was amplified from a *Dra*I digested Genome walker DNA library. The first cycle of amplification was performed with primers mhpl114 and

25 Ap1 and the second with primers mhpl103 and AP2. The assembled sequence (SEQ ID NOs:43, 45) is 2396 nucleotides long. It contains an open reading frame of 1605 nucleotides, which encode a polypeptide of 535 amino acids (SEQ ID NOs:44, 45), 196 nucleotides of 3' untranslated

30

35

region (UTR), and an upstream sequence which includes the promoter region and the 5'-UTR of the mouse *hpa* cDNA.. According to two promoter predicting programs TSSW and TSSG, the transcription start site is localized to nucleotide 431 of SEQ ID NOs:43, 45, 163 nucleotides upstream of the first ATG codon. The 431 upstream genomic sequence contains the promoter region. A TATA box is predicted at position 394 of SEQ ID NOs:43, 45. The mouse and the human *hpa* genes share an average homology of 78 % between the nucleotide sequences and 81 % similarity between the deduced amino acid sequences.

Search for *hpa* homologous sequences, using the Blast 2.0 server revealed two EST's from rat: AI060284 (385 nucleotides, SEQ ID NO:46) which is homologous to the amino terminus (68 % similarity to amino acids 12-136) of human heparanase and AI237828 (541 nucleotides, SEQ ID NO:47) which is homologous to the carboxyl terminus (81 % similarity to amino acids 500-543) of human heparanase, and contains a 3'-UTR. A comparison between the human heparanase and the mouse and rat homologous sequences is demonstrated in Figure 17.

EXAMPLE 13

Prediction of heparanase active site

Homology search of heparanase amino acid sequence against the DNA and the protein databases revealed no significant homologies. The protein secondary structure as predicted by the PHD program consists of alternating alpha helices and beta sheets. The fold recognition server of UCLA predicted alpha/beta barrel structure, with under-threshold confidence.

Five of 15 proteins, which were predicted to have most similar folds, were glycosyl hydrolases from various organisms: 1xyza – xylanase from *Clostridium Thermocellum*, 1pbga – 6-phospho-beta- δ -galactosidase from *Lactococcus Lactis*, 1amy – alpha-amylase from Barley, 1ecea – endocellulase from *Acidothermus Cellulolyticus* and 1qbc – hexosaminidase alpha chain, glycosyl hydrolase.

Protein homology search using the bioaccelerator pulled out several proteins, including glycosyl hydrolases such as beta-fructofuranosidase from *Vicia faba* (broad bean) and from potato, lactase phlorizin hydrolase from human, xylanases from *Clostridium thermocellum* and from *Streptomyces halstedii* and cellulase from *Clostridium thermocellum*. Blocks 9.3 database pulled out the active site of glycosyl hydrolases family

five, which includes cellulases from various bacteria and fungi. Similar active site motif is shared by several lysosomal acid hydrolases (63) and other glycosyl hydrolases. The common mechanism shared by these enzymes involves two glutamic acid residues, a proton donor and a nucleophile.

Despite the lack of an overall homology between the heparanase and other glycosyl hydrolases, the amino acid couple Asp-Glu (NE), which is characteristic of the proton donor of glycosyl hydrolyses of the GH-A clan, was found at positions 224-225 of the human heparanase protein sequence. As in other clan members, this NE couple is located at the end of a β sheet.

Considering the relative location of the proton donor and the predicted secondary structure, the glutamic acid that functions as nucleophile is most likely located at position 343, or at position 396. Identification of the active site and the amino acids directly involved in hydrolysis opens the way for expression of the defined catalytic domain. In addition, it will provide the tools for rational design of enzyme activity either by modification of the microenvironment or catalytic site itself.

EXAMPLE 14

Expression of hpa antisense in mammalian cell lines

A mammalian expression vector Hpa2Kepcdna3 was constructed in order to express *hpa* antisense in mammalian cells. *hpa* cDNA (1.7 kb *EcoRI* fragment) was cloned into the plasmid pCDNA3 in 3'>5' (antisense) orientation. The construct was used to transfect MBT2-T50 and T24P cell lines. 2 x 10⁵ cells in 35 mm plates were transfected using the Fugene protocol (Boehringer Mannheim). 48 hours after transfection cells were trypsinized and seeded in six well plates. 24 hours later G418 was added to initiate selection. The number of colonies per 35 mm plate following 3 weeks:

	Antisense	No insert
T24P	15	60
MBT-T50	1	6

The lower number of colonies obtained after transfection with *hpa* antisense, as compared with the control plasmid suggests that the introduction of *hpa* antisense interfere with cell growth. This experiment demonstrates the use of complementary antisense *hpa* DNA sequence to

control heparanase expression in cells. This approach may be used to inhibit expression of heparanase *in vivo*, in, for example, cancer cells and in other pathological processes in which heparanase is involved.

5

EXAMPLE 15

Zoo blot

Hpa cDNA was used as a probe to detect homologous sequences in human DNA and in DNA of various animals. The autoradiogram of the Southern analysis is presented in Figure 18. Several bands were detected in human DNA, which correlated with the accepted pattern according to the genomic *hpa* sequence. Several intense bands were detected in all mammals, while faint bands were detected in chicken. This correlates with the phylogenetic relation between human and the tested animals. The intense bands indicate that *hpa* is conserved among mammals as well as in more genetically distant organisms. The multiple bands patterns suggest that in all animals, like in human, the *hpa* locus occupy large genomic region. Alternatively, the various bands could represent homologous sequences and suggest the existence of a gene family, which can be isolated based on their homology to the human *hpa* reported herein. This conservation was actually found, between the isolated human *hpa* cDNA and the mouse homologue.

10
15
20

EXAMPLE 16

Characterization of the hpa promoter

The DNA sequence upstream of the *hpa* first ATG was subjected to computational analysis in order to localize the predicted transcription start site and to identify potential transcription factors binding sites. Recognition of human PolII promoter region and start of transcription were predicted using the TSSW and TSSG programs. Both programs identified a promoter region upstream of the coding region. TSSW pointed at nucleotide 2644 and TSSG at 2635 of SEQ ID NO:42. These two predicted transcription start sites are located 4 and 13 nucleotides upstream of the longest *hpa* cDNA isolated by RACE.

25
30

A *hpa* promoter-GFP reporter vector was constructed in order to investigate the regulation of *hpa* transcription. Two constructs were made, containing 1.8 kb and 1.1 kb of the *hpa* promoter region. The reporter vector was transfected into T50-mouse bladder carcinoma cells. Cells transfected with both constructs exhibited green fluorescence, which

35

indicated the promoter activity of the genomic sequence upstream of the *hpa*-coding region. This reporter vector, enables the monitoring of *hpa* promoter activity, at various conditions and in different cell types and to characterize the factors involved regulation of *hpa* expression.

5

Although the invention has been described in conjunction with specific embodiments thereof, it is evident that many alternatives, modifications and variations will be apparent to those skilled in the art. Accordingly, it is intended to embrace all such alternatives, modifications and variations that fall within the spirit and broad scope of the appended claims.

10

LIST OF REFERENCES

1. Wight, T.N., Kinsella, M.G., and Qwarnstromn, E.E. (1992). The role of proteoglycans in cell adhesion, migration and proliferation. *Curr. Opin. Cell Biol.*, 4, 793-801.
2. Jackson, R.L., Busch, S.J., and Cardin, A.L. (1991). Glycosaminoglycans: Molecular properties, protein interactions and role in physiological processes. *Physiol. Rev.*, 71, 481-539.
3. Wight, T.N. (1989). Cell biology of arterial proteoglycans. *Arteriosclerosis*, 9, 1-20.
4. Kjellen, L., and Lindahl, U. (1991). Proteoglycans: structures and interactions. *Annu. Rev. Biochem.*, 60, 443-475.
5. Ruoslahti, E., and Yamaguchi, Y. (1991). Proteoglycans as modulators of growth factor activities. *Cell*, 64, 867-869.
6. Vlodavsky, I., Eldor, A., Haimovitz-Friedman, A., Matzner, Y., Ishai-Michaeli, R., Levi, E., Bashkin, P., Lider, O., Naparstek, Y., Cohen, I.R., and Fuks, Z. (1992). Expression of heparanase by platelets and circulating cells of the immune system: Possible involvement in diapedesis and extravasation. *Invasion & Metastasis*, 12, 112-127.
7. Vlodavsky, I., Mohsen, M., Lider, O., Ishai-Michaeli, R., Ekre, H.-P., Svahn, C.M., Vigoda, M., and Peretz, T. (1995). Inhibition of tumor metastasis by heparanase inhibiting species of heparin. *Invasion & Metastasis*, 14, 290-302.
8. Nakajima, M., Irimura, T., and Nicolson, G.L. (1988). Heparanase and tumor metastasis. *J. Cell. Biochem.*, 36, 157-167.
9. Nicolson, G.L. (1988). Organ specificity of tumor metastasis: Role of preferential adhesion, invasion and growth of malignant cells at specific secondary sites. *Cancer Met. Rev.*, 7, 143-188.

10. Liotta, L.A., Rao, C.N., and Barsky, S.H. (1983). Tumor invasion and the extracellular matrix. *Lab. Invest.*, 49, 639-649.
11. Vlodavsky, I., Fuks, Z., Bar-Ner, M., Ariav, Y., and Schirmacher, V. (1983). Lymphoma cell mediated degradation of sulfated proteoglycans in the subendothelial extracellular matrix: Relationship to tumor cell metastasis. *Cancer Res.*, 43, 2704-2711.
12. Vlodavsky, I., Ishai-Michaeli, R., Bar-Ner, M., Fridman, R., Horowitz, A.T., Fuks, Z. and Biran, S. (1988). Involvement of heparanase in tumor metastasis and angiogenesis. *Is. J. Med.*, 24, 464-470.
13. Vlodavsky, I., Liu, G.M., and Gospodarowicz, D. (1980). Morphological appearance, growth behavior and migratory activity of human tumor cells maintained on extracellular matrix vs. plastic. *Cell*, 19, 607-616.
14. Gospodarowicz, D., Delgado, D., and Vlodavsky, I. (1980). Permissive effect of the extracellular matrix on cell proliferation in-vitro. *Proc. Natl. Acad. Sci. USA.*, 77, 4094-4098.
15. Bashkin, P., Doctrow, S., Klagsbrun, M., Svahn, C.M., Folkman, J., and Vlodavsky, I. (1989). Basic fibroblast growth factor binds to subendothelial extracellular matrix and is released by heparitinase and heparin-like molecules. *Biochemistry*, 28, 1737-1743.
16. Parish, C.R., Coombe, D.R., Jakobsen, K.B., and Underwood, P.A. (1987). Evidence that sulphated polysaccharides inhibit tumor metastasis by blocking tumor cell-derived heparanase. *Int. J. Cancer*, 40, 511-517.
- 16a. Vlodavsky, I., Hua-Quan Miao, Benezra, M., Lider, O., Bar-Shavit, R., Schmidt, A., and Peretz, T. (1997). Involvement of the extracellular matrix, heparan sulfate proteoglycans and heparan sulfate degrading enzymes in angiogenesis and metastasis. In: *Tumor Angiogenesis*. Eds. C.E. Lewis, R. Bicknell & N. Ferrara. Oxford University Press, Oxford UK, pp. 125-140.

17. Burgess, W.H., and Maciag, T. (1989). The heparin-binding (fibroblast) growth factor family of proteins. *Annu. Rev. Biochem.*, 58, 575-606.
18. Folkman, J., and Klagsbrun, M. (1987). Angiogenic factors. *Science*, 235, 442-447.
19. Vlodavsky, I., Folkman, J., Sullivan, R., Fridman, R., Ishai-Michaeli, R., Sasse, J., and Klagsbrun, M. (1987). Endothelial cell-derived basic fibroblast growth factor: Synthesis and deposition into subendothelial extracellular matrix. *Proc. Natl. Acad. Sci. USA*, 84, 2292-2296.
20. Folkman, J., Klagsbrun, M., Sasse, J., Wadzinski, M., Ingber, D., and Vlodavsky, I. (1980). A heparin-binding angiogenic protein - basic fibroblast growth factor - is stored within basement membrane. *Am. J. Pathol.*, 130, 393-400.
21. Cardon-Cardo, C., Vlodavsky, I., Haimovitz-Friedman, A., Hicklin, D., and Fuks, Z. (1990). Expression of basic fibroblast growth factor in normal human tissues. *Lab. Invest.*, 63, 832-840.
22. Ishai-Michaeli, R., Svahn, C.-M., Chajek-Shaul, T., Korner, G., Ekre, H.-P., and Vlodavsky, I. (1992). Importance of size and sulfation of heparin in release of basic fibroblast factor from the vascular endothelium and extracellular matrix. *Biochemistry*, 31, 2080-2088.
23. Ishai-Michaeli, R., Eldor, A., and Vlodavsky, I. (1990). Heparanase activity expressed by platelets, neutrophils and lymphoma cells releases active fibroblast growth factor from extracellular matrix. *Cell Reg.*, 1, 833-842.
24. Vlodavsky, I., Bar-Shavit, R., Ishai-Michaeli, R., Bashkin, P., and Fuks, Z. (1991). Extracellular sequestration and release of fibroblast growth factor: a regulatory mechanism? *Trends Biochem. Sci.*, 16, 268-271.
25. Vlodavsky, I., Bar-Shavit, R., Korner, G., and Fuks, Z. (1993). Extracellular matrix-bound growth factors, enzymes and plasma proteins. In *Basement membranes: Cellular and molecular aspects* (eds.

D.H. Rohrbach and R. Timpl), pp327-343. Academic press Inc., Orlando, Fl.

26. Yayon, A., Klagsbrun, M., Esko, J.D., Leder, P., and Ornitz, D.M. (1991). Cell surface, heparin-like molecules are required for binding of basic fibroblast growth factor to its high affinity receptor. *Cell*, 64, 841-848.
27. Spivak-Kroizman, T., Lemmon, M.A., Dikic, I., Ladbury, J.E., Pinchasi, D., Huang, J., Jaye, M., Crumley, G., Schlessinger, J., and Lax, I. (1994). Heparin-induced oligomerization of FGF molecules is responsible for FGF receptor dimerization, activation, and cell proliferation. *Cell*, 79, 1015-1024.
28. Ornitz, D.M., Herr, A.B., Nilsson, M., West, a., J., Svahn, C.-M., and Waksman, G. (1995). FGF binding and FGF receptor activation by synthetic heparan-derived di- and trisaccharides. *Science*, 268, 432-436.
29. Gitay-Goren, H., Soker, S., Vlodavsky, I., and Neufeld, G. (1992). Cell surface associated heparin-like molecules are required for the binding of vascular endothelial growth factor (VEGF) to its cell surface receptors. *J. Biol. Chem.*, 267, 6093-6098.
30. Lider, O., Baharav, E., Mekori, Y., Miller, T., Naparstek, Y., Vlodavsky, I., and Cohen, I.R. (1989). Suppression of experimental autoimmune diseases and prolongation of allograft survival by treatment of animals with heparinoid inhibitors of T lymphocyte heparanase. *J. Clin. Invest.*, 83, 752-756.
31. Lider, O., Cahalon, L., Gilat, D., HersHKovitz, R., Siegel, D., Margalit, R., Shoseyov, O., and Cohn, I.R. (1995). A disaccharide that inhibits tumor necrosis factor α is formed from the extracellular matrix by the enzyme heparanase. *Proc. Natl. Acad. Sci. USA.*, 92, 5037-5041.
- 31a. Rapraeger, A., Krufka, A., and Olwin, B.R. (1991). Requirement of heparan sulfate for bFGF-mediated fibroblast growth and myoblast differentiation. *Science*, 252, 1705-1708.

32. Eisenberg, S., Sehayek, E., Olivecrona, T., and Vlodavsky, I. (1992). Lipoprotein lipase enhances binding of lipoproteins to heparan sulfate on cell surfaces and extracellular matrix. *J. Clin. Invest.*, 90, 2013-2021.
33. Shieh, M-T., Wundunn, D., Montgomery, R.I., Esko, J.D., and Spear, P.G. J. (1992). Cell surface receptors for herpes simplex virus are heparan sulfate proteoglycans. *J Cell Biol.*, 116, 1273-1281.
- 33a. Chen, Y., Maguire, T., Hileman, R.E., Fromm, J.R., Esko, J.D., Linhardt, R.J., and Marks, R.M. (1997). Dengue virus infectivity depends on envelope protein binding to target cell heparan sulfate. *Nature Medicine* 3, 866-871.
- 33b. Putnak, J.R., Kanesa-Thasan, N., and Innis, B.L. (1997). A putative cellular receptor for dengue viruses. *Nature Medicine* 3, 828-829.
34. Narindrasorasak, S., Lowery, D., Gonzalez-DeWhitt, P., Poorman, R.A., Greenberg, B., Kisilevsky, R. (1991). High affinity interactions between the Alzheimer's beta-amyloid precursor protein and the basement membrane form of theparan sulfate proteoglycan. *J. Biol. Chem.*, 266, 12878-83.
35. Ross, R. (1993). The pathogenesis of atherosclerosis: a perspective for the 1990s. *Nature (Lond.)*, 362:801-809.
36. Zhong-Sheng, J., Walter, J., Brecht, R., Miranda, D., Mahmood Hussain, M., Innerarity, T.L. and Mahley, W.R. (1993). Role of heparan sulfate proteoglycans in the binding and uptake of apolipoprotein E-enriched remnant lipoproteins by cultured cells. *J. Biol. Chem.*, 268, 10160-10167.
37. Ernst, S., Langer, R., Cooney, Ch.L., and Sasisekharan, R. (1995). Enzymatic degradation of glycosaminoglycans. *Critical Reviews in Biochemistry and Molecular Biology*, 30(5), 387-444.

38. Gospodarowicz, D., Mescher, AL., Birdwell, CR. (1977). Stimulation of corneal endothelial cell proliferation in vitro by fibroblast and epidermal growth factors. *Exp Eye Res* 25, 75-89.
39. Haimovitz-Friedman, A., Falcone, D.J., Eldor, A., Schirmacher, V., Vlodavsky, I., and Fuks, Z. (1991) Activation of platelet heparitinase by tumor cell-derived factors. *Blood*, 78, 789-796.
- 39a. Savitsky, K., Platzer, M., Uziel, T., Gilad, S., Sartiel, A., Rosental, A., Elroy-Stein, O., Siloh, Y. and Rotman, G. (1997). Ataxia-telangiectasia: structural diversity of untranslated sequences suggests complex post-translational regulation of ATM gene expression. *Nucleic Acids Res.* 25(9), 1678-1684.
40. Bar-Ner, M., Eldor, A., Wasserman, L., Matzner, Y., and Vlodavsky, I. (1987). Inhibition of heparanase mediated degradation of extracellular matrix heparan sulfate by modified and non-anticoagulant heparin species. *Blood*, 70, 551-557.
41. Goshen, R., Hochberg, A., Korner, G., Levi, E., Ishai-Michaeli, R., Elkin, M., de Grot, N., and Vlodavsky, I. (1996). Purification and characterization of placental heparanase and its expression by cultured cytotrophoblasts. *Mol. Human Reprod.*, 2, 679-684.
42. Korb M., Ke Y. and Johnson L.F. (1993) Stimulation of gene expression by introns: conversion of an inhibitory intron to a stimulatory intron by alteration of the splice donor sequence. *Nucleic Acids Res.*, 21(25):5901-8.
43. Zheng B., Qiu X.Y., Tan M., Xing Y.N., Lo D., Xue J.L. and Qiu X.F. (1997) Increment of hFIX expression with endogenous intron 1 in vitro. *Cell Res.*, 7(1):21-29.
44. Kurachi S., Hitomi Y., Furukawa M. and Kurachi K. (1995) Role of intron I in expression of the human factor IX gene. *J. Biol. Chem.* 10, 270(10):5276-5281.

45. Shekhar P.V. and Miller F.R. (1994-5) Correlation of differences in modulation of ras expression with metastatic competence of mouse mammary tumor subpopulations. *Invasion Metastasis*, 14(1-6):27-37.
46. Zhou G., Garofalo S., Mukhopadhyay K., Lefebvre V., Smith C.N., Eberspaecher H. and de Crombrughe B. (1995) A 182 bp fragment of the mouse pro alpha 1(II) collagen gene is sufficient to direct chondrocyte expression in transgenic mice. *J. Cell Sci.*, 108 (Pt 12):3677-3684.
47. Hormuzdi S.G., Penttinen R., Jaenisch R. and Bornstein P. (1998) A gene-targeting approach identifies a function for the first intron in expression of the alpha1(I) collagen gene. *Mol. Cell*, 18(6):3368-3375.
48. Kang Y.K., Lee C.S., Chung A.S. and Lee K.K. (1998) Prolactin-inducible enhancer activity of the first intron of the bovine beta-casein gene. *Mol. Cells*, 30;8(3):259-265.
49. Chow Y.H., O'Brodovich H., Plumb J., Wen Y., Sohn K.J., Lu Z., Zhang F., Lukacs G.L., Tanswell A.K., Hui C.C., Buchwald M. and Hu J. (1997) Development of an epithelium-specific expression cassette with human DNA regulatory elements for transgene expression in lung airways. *Proc. Natl. Acad. Sci. USA*, 23;94(26):14695-14700.
50. Gottschalk U. and Chan S. (1998) Somatic gene therapy. Present situation and future perspective. *Arzneimittelforschung*, 48(11):1111-1120.
51. Ye S., Cole-Strauss A.C., Frank B. and Kmiec E.B. (1998) Targeted gene correction: a new strategy for molecular medicine. *Mol. Med. Today*, 4(10):431-437.
52. Lai L., and Lien Y. (1999) Homologous recombination based gene therapy. *Exp. Nephrol.*, 7(1):11-14.

53. Yazaki N., Fujita H., Ohta M., Kawasaki T. and Itoh N. (1993) The structure and expression of the FGF receptor-1 mRNA isoforms in rat tissues. *Biochim. Biophys. Acta.*, 20;1172(1-2):37-42.
54. Le Fur N., Kelsall S.R., Silvers W.K. and Mintz B. (1997) Selective increase in specific alternative splice variants of tyrosinase in murine melanomas: a projected basis for immunotherapy. *Proc. Natl. Acad. Sci. USA*, 13;94(10):5332-5337.
55. Miyake H., Okamoto I., Hara I., Gohji K., Yamanaka K., Arakawa S., Kamidono S. and Saya H. (1998) Highly specific and sensitive detection of malignancy in urine samples from patients with urothelial cancer by CD44v8-10/CD44v10 competitive RT-PCR. *Int. J. Cancer*, 18;79(6):560-564.
56. Guriec N., Marcellin L., Gairard B., Calderoli H., Wilk A., Renaud R., Bergerat J.P. and Oberling F. (1996) CD44 exon 6 expression as a possible early prognostic factor in primary node negative breast carcinoma. *Clin. Exp. Metastasis*, 14(5):434-439.
57. Gewirtz A.M., Sokol D.L. and Ratajczak M.Z. (1998) Nucleic acid therapeutics: state of the art and future prospects. *Blood*, 1;92(3):712-736.
58. Hida K., Shindoh M., Yasuda M., Hanzawa M., Funaoka K., Kohgo T., Amemiya A., Totsuka Y., Yoshida K. and Fujinaga K (1997) Antisense E1AF transfection restrains oral cancer invasion by reducing matrix metalloproteinase activities. *Am. J. Pathol.* 150(6):2125-2132.
59. Shastry B.S. (1998) Gene disruption in mice: models of development and disease. *Mol. Cell. Biochem.* 1998 Apr;181(1-2):163-179.
60. Carpentier A.F., Rosenfeld M.R., Delattre J.Y., Whalen R.G., Posner J.B. and Dalmau J. (1998) DNA vaccination with HuD inhibits growth of a neuroblastoma in mice. *Clin. Cancer Res.*, 4(11):2819-2824.
61. Lai W.C. and Bennett M. (1998) DNA vaccines. *Crit. Rev. Immunol.*, 18(5):449-484.

62. Welch P.J., Barber J.R., and Wong-Staal F. (1998) Expression of ribozymes in gene transfer systems to modulate target RNA levels. *Curr. Opin. Biotechnol.*, 9(5):486-496.
63. Durand P., Lehn P., Callebaut I., Fabrega S., Henrissat B. and Moron J.P. (1997) Active-site motifs of lysosomal acid hydrolases: invariant features of clan GH-A glycosyl hydrolases deduced from hydrophobic cluster analysis. *Glycobiology*, 7(2):277-284.
64. Thuong and Helene (1993) Sequence specific recognition and modification of double helical DNA by oligonucleotides *Angev. Chem. Int. Ed. Engl.* 32:666
65. Dash P., Lotan I., Knapp M., Kandel E.R. and Goelet P. (1987) Selective elimination of mRNAs in vivo: complementary oligodeoxynucleotides promote RNA degradation by an RNase H-like activity. *Proc. Natl. Acad. Sci. USA*, 84:7896.
66. Chiang M.Y., Chan H., Zounes M.A., Freier S.M., Lima W.F. and Bennett C.F. (1991) Antisense oligonucleotides inhibit intercellular adhesion molecule 1 expression by two distinct mechanisms. *J. Biol. Chem.* 266:18162-71.
- 5 67. Paterson Paterson B.M, Roberts B.E and Kuff EL . (1977) Structural gene identification and mapping by DNA-mRNA hybrid-arrested cell-free translation. *Proc. Natl. Acad. Sci. USA*, 74:4370.
68. Cohen (1992) Oligonucleotide therapeutics. *Trends in Biotechnology*, 10:87.
69. Szczylik et al (1991) Selective inhibition of leukemia cell proliferation by BCR-ABL antisense oligodeoxynucleotides. *Science* 253:562.
70. Calabretta et al. (1991) Normal and leukemic hematopoietic cell manifest differential sensitivity to inhibitory effects of c-myc antisense

oligodeoxynucleotides: an in vitro study relevant to bone marrow purging. Proc. Natl. Acad. Sci. USA 88:2351.

71. Heikhila et al. (1987) A c-myc antisense oligodeoxynucleotide inhibits entry into S phase but not progress from G(0) to G(1). Nature, 328:445.

72. Reed et al. (1990) Antisense mediated inhibition of BCL2 protooncogene expression and leukemic cell growth and survival: comparison of phosphodiester and phosphorothioate oligodeoxynucleotides. Cancer Res. 50:6565.

73. Burch and Mahan (1991) Oligodeoxynucleotides antisense to the interleukin I receptor mRNA block the effects of interleukin I in cultured murine and human fibroblasts and in mice. J. Clin. Invest. 88:1190.

74. Agrawal (1992) Antisense oligonucleotides as antiviral agents. TIBTECH 10:152.

75. Uhlmann et al. (1990) Chem. Rev. 90:544.

76. Cook (1991) Medicinal chemistry of antisense oligonucleotides - future opportunities. Anti-Cancer Drug Design 6:585.

77. Biotechnology research news (1993) Can DNA mimics improve on the real thing? Science 262:1647.

WHAT IS CLAIMED IS:

1. An isolated nucleic acid comprising a genomic, complementary or composite polynucleotide sequence encoding a polypeptide having heparanase catalytic activity.
2. The isolated nucleic acid of claim 1, wherein said polynucleotide or a portion thereof is hybridizable with SEQ ID NOs: 9, 13, 42, 43 or a portion thereof at 68 °C in 6 x SSC, 1 % SDS, 5 x Denharts, 10 % dextran sulfate, 100 µg/ml salmon sperm DNA, and ³²p labeled probe and wash at 68 °C with 3 x SSC and 0.1 % SDS.
3. The isolated nucleic acid of claim 1, wherein said polynucleotide or a portion thereof is at least 60 % identical with SEQ ID NOs: 9, 13, 42, 43 or portions thereof as determined using the Bestfit procedure of the DNA sequence analysis software package developed by the Genetic Computer Group (GCG) at the university of Wisconsin (gap creation penalty - 12, gap extension penalty - 4).
4. The isolated nucleic acid of claim 1, wherein said polypeptide is as set forth in SEQ ID NOs:10, 14, 44 or portions thereof.
5. The isolated nucleic acid of claim 1, wherein said polypeptide is at least 60 % homologous to SEQ ID NOs:10, 14, 44 or portions thereof as determined with the Smith-Waterman algorithm, using the Bioaccelerator platform developed by Compugene (gapop: 10.0, gapext: 0.5, matrix: blosum62).
6. A nucleic acid construct comprising the isolated nucleic acid of claim 1.
7. A host cell comprising the nucleic acid construct of claim 6.
8. An antisense oligonucleotide comprising a polynucleotide or a polynucleotide analog of at least 10 bases being hybridizable *in vivo*, under physiological conditions, with a portion of a polynucleotide strand encoding a polypeptide having heparanase catalytic activity.

9. The antisense oligonucleotide of claim 8, wherein said polynucleotide strand encoding said polypeptide having heparanase catalytic activity is as set forth in SEQ ID NOs: 9, 13, 42, or 43.

10. The antisense oligonucleotide of claim 8, wherein said polypeptide having heparanase catalytic activity is as set forth in SEQ ID NOs: 10, 14 and 44.

11. A method of *in vivo* downregulating heparanase activity comprising the step of *in vivo* administering the antisense oligonucleotide of claim 8.

12. A pharmaceutical composition comprising the antisense oligonucleotide of claim 8 and a pharmaceutically acceptable carrier.

13. A ribozyme comprising the antisense oligonucleotide of claim 8 and a ribozyme sequence.

14. An antisense nucleic acid construct comprising a promoter sequence and a polynucleotide sequence directing the synthesis of an antisense RNA sequence of at least 10 bases being hybridizable *in vivo*, under physiological conditions, with a portion of a polynucleotide strand encoding a polypeptide having heparanase catalytic activity.

15. The antisense nucleic acid construct of claim 14, wherein said polynucleotide strand encoding said polypeptide having heparanase catalytic activity is as set forth in SEQ ID NOs: 9, 13, 42 or 43.

16. The antisense nucleic acid construct of claim 14, wherein said polypeptide having heparanase catalytic activity is as set forth in SEQ ID NOs: 10, 14 or 44.

17. A method of *in vivo* downregulating heparanase activity comprising the step of *in vivo* administering the antisense nucleic acid construct of claim 14.

18. A pharmaceutical composition comprising the antisense nucleic acid construct of claim 14 and a pharmaceutically acceptable carrier.

19. A nucleic acid construct comprising a polynucleotide sequence functioning as a promoter, said polynucleotide sequence is derived from SEQ ID NO:42 and includes at least nucleotides 2535-2635 thereof or from SEQ ID NO:43 and includes at least nucleotides 320-420.

20. A method of expressing a polynucleotide sequence comprising the step of ligating the polynucleotide sequence into the nucleic acid construct of claim 19, downstream of said polynucleotide sequence derived from SEQ ID NOs:42 or 43.

21. A recombinant protein comprising a polypeptide having heparanase catalytic activity.

22. The recombinant protein of claim 21, wherein said polypeptide includes at least a portion of SEQ ID NOs:10, 14 or 44.

23. The recombinant protein of claim 21, wherein the protein is encoded by a polynucleotide hybridizable with SEQ ID NOs: 9, 13, 42, 43 or a portion thereof at 68 °C in 6 x SSC, 1 % SDS, 5 x Denharts, 10 % dextran sulfate, 100 µg/ml salmon sperm DNA, and ³²p labeled probe and wash at 68 °C with 3 x SSC and 0.1 % SDS.

24. The recombinant protein of claim 21, wherein the protein is encoded by a polynucleotide at least 60 % identical with SEQ ID NOs: 9, 13, 42, 43 or portions thereof as determined using the Bestfit procedure of the DNA sequence analysis software package developed by the Genetic Computer Group (GCG) at the university of Wisconsin (gap creation penalty - 12, gap extension penalty - 4).

25. A pharmaceutical composition comprising, as an active ingredient, the recombinant protein of claim 21.

26. A method of identifying a chromosome region harboring a heparanase gene in a chromosome spread comprising the steps of:

- (a) hybridizing the chromosome spread with a tagged polynucleotide probe encoding heparanase;
- (b) washing the chromosome spread, thereby removing excess of non-hybridized probe; and
- (c) searching for signals associated with said hybridized tagged polynucleotide probe, wherein detected signals being indicative of a chromosome region harboring a heparanase gene.

27. A method of *in vivo* eliciting anti-heparanase antibodies comprising the steps of administering a nucleic acid construct including a polynucleotide segment corresponding to at least a portion of SEQ ID NOs:9, 13 or 43 and a promoter for directing the expression of said polynucleotide segment *in vivo*.

28. A DNA vaccine for *in vivo* eliciting anti-heparanase antibodies comprising a nucleic acid construct including a polynucleotide segment corresponding to at least a portion of SEQ ID NOs:9, 13 or 43 and a promoter for directing the expression of said polynucleotide segment *in vivo*.


1/34

Fig 1

1 CTAGAGCTTTGCACTCTCCGCTGCGCGGAGCTGGGGGGGAGCAGCCAGGTGAGCCCA
51 AGATGCTGCTGCGCTCGAAGCTGCGCTGCGCGCGCGCTGATGCTGCTGCTGCTGCGGC
M L L R S K P A L P P P L M L T L L G P
121 CGCTGGGTCCCTCTCCCTGCGCGCTGCGCGAGCTGCGCAAGCAGGACGTGTGTG
L G P L S P G A L P R P A Q A Q D V V D
181 ACCTGGACTTCTTCACCAGGAGCGCTGCACTGCTGAGCCCTGCTTCTGCTGCTGCA
L D F F T Q E P L H L V S P S F L S V T
241 CCATTGACGCCAACCTGGCCACGGACCGCGGTTCCTCATCTCTGCGTTCTCCAAAGC
I D A N L A T D P R F L I L L G S P K L
301 TTGTAACCTTGGCCAGAGGCTTGTCTCTGCTACCTGAGGTTTGGTGGCACCAGACAG
R T L A R G L S P A Y L R F G G T K T D
361 ACTTCTTAATTTTCGATCCCAAGAGGAATCAACCTTTGAAGAGAGAGTTACTGCGAAT
P L I F D P K K E S T P E E R S Y W Q S
421 CTCAGTCAACCAGGATATTTCRAATATGGATCCATCCCTCTGATGTGGAGGAGAGT
Q V N Q D I C K Y G S I P P D V E E K L
481 TACGGTTGGAATGGCCCTACCAGGAGCAATTGCTACTCCGAGAACACTACCAGAAAAGT
R L E W P Y Q E Q L L L R E H Y Q K K F
541 TCAAGAACAGCACCTACTCAAGAGCTCTGTAGATGTGCTATACACTTTTGCAAACTGCT
K N S T Y S R S S V D V L Y T F A N C S
601 CAGGACTGGACTTGATCTTTGGCCTAAATGCGTTATTAAGAACAGCAGATTTCAGTGGAA
G L D L I F G L N A L L R T A D L Q W N
661 ACAGTTCTAATGCTCAGTTGCTCTGCTGACTACTGCTCTTCCAAGGGGTATAACATTCTT
S S N A Q L L L D Y C S S K G Y N I S W
721 GGGAACTAGGCAATGAACCTAACAGTTTCCTTAAGAAGGCTGATATTTTCATCAATGGGT
E L G N E P N S P L K K A D I F I N G S
(T)
781 CSCAGTTAGGAGAAGATTATATTCATTTGCATAAACTTCTAAGAAAGTCCACCTTCAAAA
Q L G E D Y I Q L H K L L R K S T P K N
(F)
841 ATGCAAACTCTATGGTCTGATGTTGGTCAGGCTCGAAGAAAGACGGCTAAGATGCTGA
A K L Y G P D V G Q P R R K T A K M L K
901 AGAGCTTCTGAAGGCTGCTGGAGAAGGATGATTGATTGATTGATTGATTGATTGATTGATT
S F L K A G G E V I D S V T W H H Y Y L
961 TGAATGGACGGACTGCTACCAGGGAAGATTTCCTAAACCTGATGATTGAGACATTTTAA
N G R T A T R E D F L N P D V L D I F I
1021 TTTTCATCTGTGCAAAAAGTTTCCAGGTGGTTGAGAGCACCAGGCTGGCAAGAGGCTCT
S S V Q K V F Q V V E S T R P G K K V W
1081 GGTAGGAGAAACAAGCTCTGCATATGGAGGCGGAGCGCCCTTGCTATCCGACACCTTTG
L G E T S S A Y G G G A P L L S D T F A
1141 CAGCTGGCTTTATGTGGCTGGATAAATTGGGCTGTGAGCCCGAATGGGAATAGAGTGG
A G F M W L D K L G L S A R M G I E V V
1201 TGATGAGGCAAGTATTCTTTGGAGCAGGAACTACCATTTAGTGGATGAAACTTCGATC
M R Q V F F G A G N Y H L V D E N P D P
1261 CTTTACCTGATTATTGGCTATCTCTCTGTTCAAGAAATGGTGGGCACCAAGGTGTTAA
L P D Y N L S L L F K K L V G T K V L M
1321 TGGCAAGCGTGCAAGGTTCAAAGAGAAGGAAGCTTCGAGTATACCTTCATTGCACAAACA
A S V Q G S K R R K L R V Y L H C T N T
1381 CTGACAATCCAAGGTATAAAGAAGGAGATTAACTCTGTATGCCATAAACCTCCATAAGC
D N P R Y K E G D L T L Y A I N L H N V
1441 TCACCAAGTACTTGGCGTTACCTATCTTTTCTAACAAGCAAGTGGATAAATACCTTC
T K Y L R L P Y P F S N K Q V D K Y L L
1501 TAAGACCTTTGGGACCTCATGGATTACTTTCCAAATCTGTCCAACTCAATGGTCTAACTC
R P L G P H G L L S K S V Q L N G L T L
1561 TAAAGATGGTGGATGATCAAACTTGGCACCTTTAATGAAAAAAGCTCTCGGCGCAGGAA
K M V D D Q T L P P L M E K P L R P G S
1621 GTTCACTGGGCTTGGCAGCTTCTCATATAGTTTTTTTGTGATAAGAAATGCCAAGTTG
S L G L P A F S Y S P F V I R N A K V A
1681 CTGCTTGCATCTGAAAAATAAATATACTAGTCTGACACTG
A C I

2/34

FIG. 2



3/34

FIG. 3A






FIG. 3B



4/34

FIG. 4



5/34

FIG. 5A





FIG. 5B



6/34

FIG. 6A





FIG. 6B



7/34

FIG. 7A





FIG. 7B



8/34

FIG. 8A





FIG. 8B



9/34

FIG. 9A





FIG. 9B



10/34

FIG. 10A





FIG. 10B

Heparin-avidogel



11/34

FIG. 11A





FIG. 11B



12/34

FIG. 12A




FIG. 12B




FIG. 12C




FIG. 12D





FIG. 12E



mouse	CTGGCAAGAAGGCTCTGGTTGGGAGAGACGAGCTTACGGTGGCGGT	50
human	CTGGCAAGAAGGCTCTGGTTAGGAGAAACAAGCTCTGCATATGGAGCGGA	1115
mouse	GCACCCTTGCTGTCCAACACCTTTGCAGCTGGCTTTATGTGGCTGGATAA	100
human	GCGCCCTTGCTATCCGACACCTTTGCAGCTGGCTTTATGTGGCTGGATAA	1165
mouse	ATTGGGCCTGTCAGCCCAGATGGGGCATAGAAGTCGTGATGAGGCAGGTGT	150
human	ATTGGGCCTGTCAGCCCGAATGGGAATAGAAGTGGTGATGAGGCAAGTAT	1215
mouse	TCTTCGGAGCAGGCAACTACCACTTAGTGATGAAAACCTTTGAGCCTTTA	200
human	TCTTTGGAGCAGGAAACTACCATTTAGTGATGAAAACCTTCGATCCTTTA	1265
mouse	CCTGATTACTGGCTCTCTCTTCTGTTCAAGAACTGGTAGGTCCCAGGGT	250
human	CCTGATTATTGGCTATCTCTTCTGTTCAAGAAATTGGTGGGCACCAAGGT	1315
mouse	GTTACTGTCAAGAGTGAAAGGGCCAGACAGGAGCAAACTCCGAGTGTATC	300
human	GTTAATGGCAAGCGTGCAAGGTTCAAAGAGAAGGAAGCTTCGAGTATACC	1365
mouse	TCCACTGCACTAACGTCTATCACCCACGATATCAGGAAGGAGATCTAACT	350
human	TTCATTGCACAAACACTGACAATCCAAGGTATAAAGAAGGAGATTAACT	1415
mouse	CTGTATGTCCTGAACCTCCATAATGTACCAAGCACTTGAAGGTACCGCC	400
human	CTGTATGCCATAAACCTCCATAACGTACCAAGTACTTGCGGTTACCCCTA	1465
mouse	TCCGTTGTTTCAGGAAACCAGTGGATACGTACCTTCTGAAGCCTTCGGGGC	450
human	TCCTTTTTTCTAACAAGCAAGTGGATAAATACCTTCTAAGACCTTTGGGAC	1515
mouse	CGGATGGATTACTTTCCAAATCTGTCCAACCTGAACGGTCAAATTCTGAAG	500
human	CTCATGGATTACTTTCCAAATCTGTCCAACCTCAATGGTCTAACTCTAAAG	1565
mouse	ATGGTGGATGAGCAGACCCTGCCAGCTTTGACAGAAAAACCTCTCCCCGC	550
human	ATGGTGGATGATCAAACCTTGCCACCTTTAATGGAAAAACCTCTCCGGCC	1615
mouse	AGGAAGTGCCTAAGCCTGCCTGCCTTTTCTATGGTTTTTTTGTGATAA	600
human	AGGAAGTTCCTGAGGCTTGCCAGCTTTCTCATATAGTTTTTTGTGATAA	1665
mouse	GAAATGCCAAAATCGCTGCTTGTATATGAAAATAAAA	637
human	GAAATGCCAAAGTTGCTGCTTGCATCTGAAAATAAAA	1702

FIG. 14





Figure 15



16/34

Figure 16

```

ggatcttgggtcactgcaatctctgcctcccatgcaattcttatgcatca      50
gcctcctgagtagcttggattataggtctgcgccaccactcctgggtaca      100
ccatgttggccaggctgggtcttgaactcttgggctctagtgatccacccg      150
ccttggcctcccaaagtgtggtgattacaggtgtgagccatcacacccgg      200
cccccggtttccatatagtaactcacatgtagaccacaaggatgcacta      250
tttagaaaacttgcaatggtccacttttcaaatacaccacaacatgttaaa      300
gaaattgggtatgactgggcatggcacagtggctcatgcctgcaatcctag      350
cattttgtgaggctgagacgggcagatcacgaggtcaggagattgagacc      400
atcctgacagacatgggtgaaatcccatctctactaaaaatacaaaacaat      450
tagccgggggtgtatggcaggccctgtagtcccagctactcgggagggtg      500
aggcaggagaatggcgtgaatccaggaggcagagcttgacgtgagccgag      550
atggtgccactgcactccagcctgggcgacagagcgagactccgtctcaa      600
aaaaaaaaaaaaaagaaagaaattgggtatgactgttgactcacaacaggag      650
tcaggggcatggggtggggtgttaagattaatgtcatgacaaatgtgaaa      700
agaaacttctgtttttccaactccacgtctgctacatattattacactc      750
ttctggtagtgtggtgtttatgtgtgaatttttttccatatgtatacagt      800
aattgtaggatgtaacctgtattctagtgtgcaaaactcactatgagctta      850
gcttttaagttgcttaagaataggtagatctatgcaaaataatgataatta      900
ttattattatttttaagagagggtctcactttgtcaccagggtggagtg      950
agtgggtgtgattaaagggtcactgcaacctccacctccagggtcaaataa      1000
acctccacactcagcctcccaagtagctggaaaccaaggcacgggccacc      1050
acgcctggctaattttttgtattttttgtagagatgggggtttcatcatgt      1100
tgccagggtgttcttgaattcctcggtcaagcaatcctcccaaccttg      1150
cctcccaaaatgctggcatcacaggcatgatggcatcactggcatcacat      1200
accatgctggcctgattttatgcaattagatatgcatttcaaaataatc      1250
tatttttatttgttgcttatttgggtgtacaatctcaagtggaaaaatct      1300
aagggttttgggtgttatttgccttactcaaccaatattttattagactctta      1350
ctaagcaccaacatgatcacatgcctgagctatggctagcatagcgtgtg      1400
agacaaacttaactctctctgttttgggtggagcatataatctagtagatgaag      1450
ccaatgttgagcaacatcacataactaacaatttgaggatgctacgagag      1500
tgtctaacaatttgaggatgctacgagagtgctacaacaatttgaggatgc      1550
tatgagagtggtgcatggagagctgcctggagattgagagaaagcttcct      1600
tgagggaagtacatttcagctgaaacacactgccatctgctcgagggtt      1650
tgtaactgcattcacatcccgattctgacacttcacatcccgattctgac      1700
acttcacccagttactgtctcagagcttgggtccgcatgtgtaaaacaag      1750
gacagtatgcacttggcagggttgtgagaagggaagagaacacaagtaaa      1800
gcacctgtatcaggcatcacataggaactaagcgtgcgatgcttgctatg      1850
attatacatcagtgttaagcatcaaggaaaagctgaagaaaagctgacca      1900
acagcgaaagataaatgcgcagaggagaaaatttggcacaaggctccaaatt      1950
caggggagctccgtactctacactttgtatgggggcttcagggtcctgagt      2000
tccagacattggagcaactaacctttaaagattgctaaatattgtcttaa      2050
tgagaagttgataaagaattttgggtggttgatctcttccagctgcagt      2100
ttagcgtatgctgaggccagattttttcaagcaaaagttaaaatcctgag      2150
aaactgcctggccaggagacaatcagattttgggtggtcgaagtgaacag      2200
caagtgtttataagctagatgggagagggaaggatgaatactccattgga      2250
gggttttactcgagggtcagagggatacccgggcgccatcagaatgggatct      2300
gggagtcggaaacgctgggttcccacgagagcgcgacagaaacagtgctc      2350
aggaagcctgggtccgggatgcccgagcgtgctccccgggctcctcccc      2400
ggcgctcctccccagcctcccgggcgcttggtatcccgccatctccgc      2450
acccttcaagtgggtggtgatttctgtaagtgaacgtgacccgccaccg      2500
aggggaaagcgagcaagggaagttaggagagagccgggcaggcggggcgggg      2550
ttggattgggagcagtgaggaggatgcagaagaggagtgaggaggatgga      2600
ggcgagctgggaggggtgaggagcgtaacgggGCGGAGGAAAGGAGAA      2650
AAGGGCGCTGGGGCTGCGGGGAGGAAGTGCTAGAGCTCTCGACTCTCCG      2700
CTGCGCGGAGCTGGCGGGGGGAGCAGCCAGGTGAGCCCAAGATGCTGCT      2750
M L L
GCGCTCGAAGCCTGCGCTGCGCGCGCGCTGATGCTGCTGCTCCTGGGGC      2800
R S K P A L P P P L M L L L L G
CGCTGGGTCCCTCTCCCTGGCGCCCTGCCCCGACCTGCGCAAGCACAG      2850

```

Fig. 16 (continued)

17/34

P L G P L S P G A L P R P A Q A Q	
GACGTCGTGGACCTGGACTTCTTCACCCAGGAGCCGCTGCACCTGGTGAG	2900
D V V D L D F F T Q E P L H L V S	
CCCCCTGTTCTCTGTCACCATTTGACGCCAACCTGGCCACGGACCCGC	2950
P S F L S V T I D A N L A T D P	
GGTTCCTCATCTCTCTGGGGtaagcgccagcctcctgggtcctgtccctt	3000
R F L I L L G	
tcctgtcctcctgacacctatgtctgccccgccagcggtctccttcttt	3050
tgcgcgggaaacaacttcacaccggaacctccccgcctgtctctccccacc	3100
ccacttcocgcctctcattctccctctcctcccttactctcagacccca	3150
aaccgctttttggggggtatcatttaaaaaatagatttaggggttacaag	3200
tgcagttctgttccatgggtatatttgcatgtgtggcatctgggtcctt	3250
agtgttaactgtcaccggaatgtgtacattgtatctaataaggtaatttct	3300
catccctcatccctctccaccctccaccttttgaggtctccagtgctt	3350
actattccaactaagtccatgtgtacacattgttttagcgccactctaaat	3400
gagcctttttgtttcattctctgtaagtgttgataggcaccacctaa	3450
ggtcaggtataagtggaaatttgaaaaagaaactgcccacttgccccagt	3500
acttccttagccaagaggagggaaccaggcaggtgcacctgaaggcctg	3550
tgagtgcttgattgtctgtgcagtgtaggacaagtaagattgtgcatagc	3600
cttctgtattttaagactgtgttaggaagatttctctttcttttctttct	3650
ttttctttttctttttcttttttttttaggcagatgaaaaggcgctca	3700
cagaacaggaataaaaaatctaaatattcaataaatgagacctaggagact	3750
actgcagtgacttacaaagtccataaaaaagatgtctctccaaaatggg	3800
gctgcataaatgtggtgctgcttatcagctctaagttttttccttacctg	3850
agaaagaaggaacctgatgcaggttcagggtcctgccccatgaatgcag	3900
gctgactccaagatggggagctacagggaacaatcccaggtcttctaggcc	3950
tcttattttaggccctgggagcctccagagatgggccacatcttgaccagcc	4000
cagatagagggaagatcaccattatctcacctctgtgtcaataacctag	4050
atgtgtcctcctgagcccacactatagttgccagcgctaattttaatgg	4100
gtagtgtactggttaagagatggacagacatcctggcttgactctcagc	4150
tctggcaagatgagtgactgtgttttccatatctctggccacaccaa	4200
ccttgatttcttctcagctgtgaatttctcaagcttgccctcaagga	4250
ttattgcccaggaatttgatgatattggtaagagcttctcagtggttgacc	4300
catagtaagtgtttgacgtttcaaacgaattgtttcttctaggacatgg	4350
tgagcatttggttagccattcaccggttttctgtttcttggatcatagtt	4400
aacctctccttttcttctggtgactacaatttctggtggggaagaatcc	4450
ttactttctgcccctccctttaaaggataggaagctgatactaggcagca	4500
ctagttgggggataggaagattgttccagagaaatgctgaaccatagggc	4550
tccagatcacaggaccccagctcttagcttgctggggtgtggggtggggg	4600
gggcggttactgaacatgggtatgaagtagatgtccatttactgaaatgt	4650
gaggacctgaggcctcttctattgtgtgagccagcatattccccaaacctc	4700
tccccaaagaaaggacagatgggggttccccctggagtaacaggtccaaa	4750
agaaaaaacatacagtggtgacttccaggatctgggcctgatccccagca	4800
gtcaagctccccgaattgactaacacccccctaacagtagaaattcca	4850
atctgcaatttagtgaggatgatacctttattcttcttaaatacatctct	4900
tcatttcccagagcaccctttttccccctcctctgcacctttttgttaaa	4950
gactggagtataatgaaataccaagagagcataacatgtgatacataaaa	5000
ctttttttctggtttacaaaacagttcattctgtccatacgtgcttctc	5050
tccaaggctggctgctgtctgttccagcccgcttcgcttgagaggccat	5100
ctgccatacctgctccccagacgcatcgacaagcaccccagagtggtat	5150
ctgctaagacctaaaagaggagggaaccccctctcctcatctaagacctc	5200
gcttctaaattagagtgtaggggtccatctccccaggagggggcaagggc	5250
ccaaacagcccagccatctcagaagacaacaactaagctttgtagggtcc	5300
acagtagaggagagtaagacgcctgtgttttaatttattacagttcctca	5350
aaagtgaagatgtgtgggcgggatggcaagagctgagcagacgaaagctg	5400
aaggaataaggaaagagaggaggacaaaacagctgacacttctcagtt	5450
cttgtcatttgcctggcctgttctaaagcaccttctaggtattaatccat	5500
ttagtcttggctacaacactgtgagtaactagttttgtcacccccatttt	5550
aaaaatgaagaaagtgaaggctcaggaggttaagtaacttggccacagtt	5600
tgaaactagactctgatcatgagataatagtgcccataaaaaaggga	5650
gcagattatatttttaaggaaagagagtaggataatggtagaaaaagat	5700

Fig. 16 (continued)

18/34

tgttttgaaaggaattgagagattgatataatgaaaagaagcattcacat 5750
gagagtacaggtatcagggcccaaaccttcatctaaggtaacttcaaagag 5800
gcctaagcaaaacttagtcactggcgtggttctagtctccatgatggcaaa 5850
tacatttgtgtacagcccaactccacacaaaacttaaataccaatgataga 5900
gcaatctaaaaatttgaagaaaaaatctttcaatttgtcgtcttcccaga 5950
gggacttaatacaagaaaccaatcaaaaatacttccctaagcctaactgtgtg 6000
cagaactccaaagagagcccagccctaaatcaacactgtccaatggaaat 6050
ataatataatgtgggcctcatatgcaaggctcatatgtaattttaaatttt 6100
ctagtagccatattaaaaaggtaaaaagaacaagtgaatttaattttaa 6150
taattttatttagttcaatagatccaaaatgttttctcagcatgtaatca 6200
atataaaaaatattaatgaggtatttatttctcttctcaaaccaagtc 6250
tattctataatctggcgtgtattatttacagcacttctcagactatattt 6300
ctttctttctttttttttccgagacaattttgctcttgtcacccaagct 6350
agagtacaatggcggttacctcggtcactgcaacctccgcctcccggtt 6400
caagttatttctcctgctcagtcctccaagtagctgggactagaggcatg 6450
caccaccaagcctggctaattgtgtatttttagtagagacagggtttcac 6500
catgttggcaggctaattctcaaacctcctgagctcaggtgatatgccac 6550
ctcgccctcccaagtggtgggattacaggcgtgagccactgcaaccggc 6600
ctcagattaaactatatttcaagcgttcagtagccaatgtagctagtgtct 6650
atggtagtggacagtagacagatctgcatttcaattaagacacgtatacaag 6700
catagttcaataatgcacggtaaaaaaaagtatagtgtgagtcgggtggt 6750
agaaatccataaactactgcagagcaaaaagtgttacgaacagcaatctcagt 6800
gataatgcaaccatgcttgccttttctcattgcaatttgcttattttcctca 6850
gcaaagttcatccatttttgccaattcaataaataatttactgataaaaaac 6900
tttcaatattagattcttgcattctcatagacagagttgcttttcacatt 6950
tagaaaaattacttcaattgttaaacacacggttttgataaccagtggttg 7000
aaagaggtgcagactccccatgtgcctattgatggcagaaaatttcacag 7050
ccaaagggaaaacaaagggctggggacaatcacacacctcatgtctcctaa 7100
ctcctgggaagtgtgtcctctctgattgagctcttattattgccttccc 7150
actaaccttactcactgttggagccctttgcagggttaacctgtctct 7200
gtcctctcacagaataatctcctctacctccttgtccaagctacaacttg 7250
gctattctctgatgacactgtcttccctgtagcccttttgagtaattggct 7300
gcatatttctcccatagtcaggttcttttctgttctccagtcctggcttct 7350
ggatgacagcccaactagtttgaactccatactgctatagttcaagtcctt 7400
tttgacttgttaccttgggcaaaattacctccttttgttcagggtccttgt 7450
ttgtaaaatgacgataataatgccatttgcctcagtggttattttgaaa 7500
ttgagtgaagaaggcgggtagcttccctacacgctcagtgtagactagc 7550
ctgatgtgcattacgggtgatgccatgactcagtggttttctcctcatctc 7600
cacatctggctctcatccagtgctcctgcttacggcactctgtccccctc 7650
ttacttactcccccttattaactgaagactggcactgatctcacagtttc 7700
ctctccacttctagctctcaccatcatccttagatgacttcaagtcacctta 7750
gataaacctgtctcagtttcttcaactcacattttttataacagataatgt 7800
tacactcaagttgttaacagaaccagcttatccagctcatgaaatgtatgc 7850
atttcatctcaactctgtattcagtgacatcctgtgggtatctggaaatc 7900
agccatgggtgagaatatttaccatggaaattggcaataactaaaaagcag 7950
agcaccttttttctgagagccagaccatagctcttctactccatagcac 8000
ccatcataaataatttttaaatacctccactgaacagcttcttctctctc 8050
tacttcttccatctctgatttgagcttcttaatttatcatgtgaaccact 8100
cttgtaataataaccccaaatccctgttccattgttcttctgctaaaaat 8150
actaaacctgggttagtccaaaccatattttctctcttggaaatctacagg 8200
gtggcccaaaaacctggaaatggaaaaatattacttatttaatttaattgt 8250
atattaataagccattttaatgcttcatctccagtcctcagtgggcaccct 8300
gtatagctgggctattgagctcttgcgggaggaggagtggaacagtcctcc 8350
cagccacacagactgatgttgacccaaaacatttttagcttccagacttc 8400
cctggcccttagtgttacccttaactctccatttctctgcttcttcacatt 8450
ctctacttttttaaaaatctctgactccaccttcaccttatcattcttagc 8500
acatgaccatacttctgcttcccaagaaaatgagcaattacttccctttt 8550
ccttttctcctgtcatcaaatctgcagacatgtcatgcctaagtccagc 8600
tttctctcttctctgatctcagtcgtcttcttccatttctgcccctgaat 8650
cccgctccctcccaaccccccaaggacttgcctctatcagtcacctcttc 8700
cctctctgtatcttcaactcctccattttactggcttcttctcctcaagc 8750

19/34

Fig. 16 (continued)

ctttcccccaagcctttcccatctcaattacctcctcgacatgcctctgc	8800
agaaaccaccccggtttctccctcccctcggcagcctgttcttccctgttc	8850
tgccctcatgatggcaccatcattgtgtcactaaaatcaatctctccgac	8900
atcatcaatggccttctcttgttgggaaacctaataaacactttatctta	8950
tttggctcttttattgggttgaatgaggttaccctcgaaatccatattaga	9000
agtccctaacccttgagtagctcagaatgtgactttatttgggaatagggtc	9050
attgcagacgtttatttagttaggatgaggtcatactggaatgtgatgggct	9100
gcttatctaataatgactgatgtccttataacaaggagaaatttggagaca	9150
gacacgcacataggagaaataccatgtgatgacaggagttatggagttgg	9200
agtcaaaaagctatgggaacttaggagaaagacctggaacaaatcctttc	9250
ctgcgcctagagagggagtagtggccctgccactaccttgaattcaacgtt	9300
tcgggttttcaaaactgtaagacaatacatttctgttgttcaaaccaatt	9350
agtttgcagtaactctgcgactgcagccctaacaactaatacagtctctt	9400
ggaggcatttggcaaggttgacaatggaagcactttcttacccttttagg	9450
tctgtcgcctttcttgttgggggtgttttctaacaattcctctccatct	9500
ctctctctctagtttgtcttaaacatttgggtgttcttccagacttctgacct	9550
aggccttcttttcaacttcacatattccctgggtggtctcaccacttcc	9600
agaaattactttaattactgtcactgcagtaactgtgctggaactgttta	9650
acaactggctctctgggaagaggggagactggttgatggtttttgctgat	9700
ttctgtggtgttaataactccctccatggccaatttccaaactgccaacagt	9750
ttaacaactggctcacaatttttctccaaatttaacatttggctttcaca	9800
ggccaacaacgtggttacagccaactccagcacacctctgcttttgtgtca	9850
gagagaagtaacttattttgtacaaaaggtaaaataaaaaacacctgcag	9900
gcccccttttttcttaacaaactgctctagaaatagaatagctgaagc	9950
ttcttttatgcattcatctgttatttccatgtcactgtggtgggtgggatt	10000
atttttccctttatttttcttctgtatatggttgaaatactgtacctttgatc	10050
agtttttagttttatggcatgttttgcacccatattaaatctagtttttgt	10100
cagagggcgctcaatatatttttctcaaaaacagaaaatatttcattgcaa	10150
aggagacaaaacaaaaggctccttaataccaaaactttgaaatgtgatttc	10200
ttgtacttggcagtggtccaagtggttaaacccaaacagttatgggttttca	10250
ttttgttcaggaaagtctttgtctggcagcgacttacccttacatcaggc	10300
gggccttgcctcattcattcacttaagtattttattaaacaccagcggtgtg	10350
ccaagtacttatctaggtatcggttagattctgataagtcagtcaggtcc	10400
ctgctctcagggtgcttgcagcagagatgggggctgcaatagagagtaag	10450
ccaaggaaatgaaaaaggaagttgatttcagagagtgatgaatgtctatga	10500
agaaaatgaaggcagcgagtgatggagagtgacccaagggtggtacag	10550
tttgtacctctaaggaccagactgtgacccaggtcactcacagatgccccg	10600
tcagtgtatggtccacagcaactttccaggtgctcgtttctcctccacttcc	10650
cagtctcttggccagcgcgactgcttacaataacagctagagggaatcta	10700
aatgaggttctctatcatcaaacccaatcaaaatgccaaaggaacagaat	10750
cagtgcctggctgaaggcagtggaacagggccagcctggagtggttctct	10800
ctgaggaagtctctcatcttgggttttagggccataccttgtgacctgtga	10850
gctaggggttggcagtccttgacatttctactgagggaactgcctgtctat	10900
attcccgccctgtatgtgtctcctgagttccagacacacagggcggaagcg	10950
cctgatggatggaagtatgtttttgggtgttccattggtatctcaaatc	11000
tacaaaacttagtgcccccttctcctcctgttcttccccatcttcagttc	11050
atcacctgttctctcatccagcaaatgatattaccatcttccaaggagctt	11100
cccaggagtaatccttgactcctcctcaacatccaattaataatcaaate	11150
taggccagggtacaatagctcacgcctataatcccagcactttgggaggct	11200
gaggcaggtggatcatttgaggccaggagttcaagaccagcctggccaac	11250
aagggtgaaacctgtctcatttaaaaaaagttatttttaaaaaactcaaatc	11300
attattttctacctctaagtgtgtcttgaatttatccatctctctccatct	11350
ctgagctgttaccttaacctcagtcctacagttttgtctacgttaacatg	11400
accagagctctgttcttagtctgggtgaggtcactccagctgcttcagatc	11450
cttccatggtcaccgttggccctcatataaagttggcactcctggacatg	11500
tggcttacggggccctccgtgatgtggccctatttgcctctctccattctgt	11550
tctctccagcctctctgcccccatctctagggaccaaccaacccttct	11600
gctcgtcaaagggtgcagcttctcttctatctctgtgtctttggacagact	11650
tttcccttcacctgggaatgctttcttcaatcctacccactctctttaat	11700
ctagataagggtttattcttttgaatgtctagcagtgaaaccatttcccc	11750
tgaaaaaccttctctaaccaacccccctaccctcagcccaagggtctagatt	11800

Fig. 16 (continued)

20/34

aggagtccctctgaaatgtttccatagcatttttaaaagaattgcctatttta 11850
cttgttcgtatctatcactaaactacaaattgtatgagaacagccactat 11900
ctctgcctgggttcaccattcatctccagcaactagcataatgcctggcag 11950
agtgcagcctgcaacaaatatttgggtgaataaaattaacagatggctttatc 12000
tccttaagtaaattcttgcttttttcacctattaaaaacagacgcacaggcc 12050
agggtgtgggtggcccatgcctgtaatcccagcactttggcaggctgaggtg 12100
ggcggatcacctgagggtcaggagttcaagaccagcctggccaacatgggtg 12150
aaaccccatctctaaataaaaaatacaaaaaattagctgggcatgggtggg 12200
tgcgtagtagtcccagctactaggaggctgaggcaagagaatcgcttgaa 12250
cccaggaggcagaggtggcagtgagccgagatcatgccactgtaçtccag 12300
cctggatgacagagaccctgtctcaaaacacacacacacacacacacaca 12350
ca 12400
taacgtgcttgttatggaacacttgtaaaaatacaggaaagtaataaaaaa 12450
gtctaccatctagctcaccacataatgaccattgctatcatcctggcata 12500
attctctcctgtatataaaatatattcttttattgttaaaattacacta 12550
tgagtactattttttattttattttactgtggcaaaaatgcgcaaaacataaaa 12600
cttgccatttttaagggtatgcagtttgggtgcattcaccacactcacattgt 12650
tgtgcaaatatcaccactatctatctcagaacttcttctgtcttcccaaac 12700
tgaaactctgtacccattaaacaatagtgcatcctctgttttccccctccc 12750
tacaattttattttttatttgggtttgtaccaaactgaaaatagctgcttct 12800
tccttacttagttcagatttagcatttccattttatttagccgtgggtttga 12850
ggatgccatgacagatgccatccttccctagagctcttggggctgtcagg 12900
tatttcagtcagggtgaattcgggttgataacattttaaaatctcacttt 12950
attctgaggttccctagtgctcagagcccaccgtatttttagggactcccaa 13000
gttacaacacaaaaatattggtgagggaatcactgaagttttaacacaag 13050
agacttacatttttgttcaatttctatcttttagtttatttccctaagcata 13100
aagaaatactttgaaaattttacatagcattatacatatttaattaaagca 13150
tgagcacatctttaaactttaaattttagatcagatctttaaattcctagg 13200
atattaagagggtactggcaatttggccagggtgtgggtgggttcacgcctata 13250
atcccaacacttttgggagggtgaagtgggcgaaattgctagagcccaggag 13300
gtggagggtgcaatggcctgagatcacgccatcgtaactccagcctggatg 13350
atgagaatgaaatccctgtctcaaaaaaaacacacacacacacacacacac 13400
gaagaagtattggcaatcagtgctccagggaataatttccctgacttgaaat 13450
aaacctacatgtagacaaactaattaggccattccaagagttgctagcct 13500
tggtttaatatgttttcagagcattccagggaagcagtggtggccagcattg 13550
catgtttgatacttcagaaatgtatgacagggtgttctcttaccaggtc 13600
ttctgttttcttagttttgtctatgttaataatttatgaaatcctcatct 13650
ttttgagggaagggtattatagatcatttctaattccattttctagcatttg 13700
gtaccatttctaagcacatgataggcaccatttggagcatttttggcttg 13750
acagaatatgcatcttagaattgttcaaattagagggtgtcagtgatgggaa 13800
ttagaatactatataatttctaagtcatttgacttaatacaaaaagaatga 13850
ttttccttgggtggggaatgggtgaaggaggaggaggagtttaagaaggaggaga 13900
agagatcctaagtcatttataaaacttctctggaagacaggtgtgtgaag 13950
acttttttaaaagtcatcaccacaaattgtgtgtgtgtgtgtgtgtgtgt 14000
ttaaatagacttttatttttagagcagtttttaggttcacagcaaaattga 14050
atgcaaggacagagatttcccataaaacccctgcccacacacatgcatag 14100
cctccctcattatcaacatccccaccagagaggtgtttgttctagttgat 14150
gaacctacactgacacatcattatcacccaaagtccatagttcacggcag 14200
gggtcactgtcggtgtacattctatgggtttgagcaaatgtataatgaca 14250
tgtatccaccattatagtaacatacagagttttttagtgccctgcaaat 14300
ccccgtttctccacctattcatccctccctctctgcatttccacccccag 14350
ccccgttaaccgctgatctttttactgtcccatagtttctggacgatcta 14400
tttttcagacagacacagagctgtctttcccttagtttctattctatcat 14450
ttcttttctccccatccatcataaaaggctatgagtttttttaagtgttg 14500
aacaccatcctacttgtcaagttaaaacataagctcctggctgggtacag 14550
tggtcatgcctgtaattctcagcattttgggaggctgtggcagaagcatc 14600
acttgaaggcagaagtttgagaccagcctgggcaacatagcaagacccca 14650
tccctcca 14700
ca 14750
ccctcagggttctagaagatcagtccttcaattagattcagattgagatg 14800
cttctcttttaaaccaatgattccctttctatcatgcccacataagaaaac 14850

Fig. 16(continued)

21/34

aaataaaaattaaacaatactgacctgtaatctcagctacccaggaggcag 14900
 aagcagaactgcttcaacccggcaagcagaagttgcagtgaagtgcagatc 14950
 gcgccactgcaactccagcctgggaaacagagcaagattctgtctcaaaaa 15000
 caaaaacaatgtgatttctcctctaagtcctgcacagggaatgttaaga 15050
 aataggtccaccaggaaagaaggaagtaagaatgtttgactagattgtct 15100
 tggaaaaaatagttatactttctgtgtcttcttaacagTCTCCAAA 15150
 S P K
 GCTTCGTACCTTGGCCAGAGGCTTGTCTCCTGCGTACCTGAGGTTTGGTG 15200
 L R T L A R G L S P A Y L R F G
 GCACCAAGACAGACTTCTAATTTTCGATCCCAAGAAGGAATCAACCTTT. 15250
 G T K T D F L I F D P K K E S T F
 GAAGAGAGAAGTTACTGGCAATCTCAAGTCAACCAGGgtgaaaattttta 15300
 E E R S Y W Q S Q V N Q
 aagattcactctatattttaattaacgtcagtcctgcatgagaatgcttt 15350
 gagaaaaactgtttttctcacacctaacaattaatgagattaacttctctc 15400
 tcccctcatctgacctgtggaggaatctgaacaagaggaggaggcagtg 15450
 gcagggttctcttatcatgatgtttgtcatgttcagtgtaggcctcaaa 15500
 aaaaaaaaaaaaaaaaaaaggcgtcctggatataactgagagctcattg 15550
 tacagtaaatattaataaaacagtgattgttagctgaaggatagaactgct 15600
 tggaggagcagatgggttagaatcgcgtaaaactaaagagcatttctagc 15650
 caaagacacaatgatagattgaaggataatttattctaaatatagaatatg 15700
 ggtgaacgagatctgtggacttctgggctccaacgttagattctgatttt 15750
 agcaagcttgtcaggggattctgatattgaaaggctgtggccttcacctg 15800
 agaaaacctgcccattggggccatgaaaatttgcctgtctttcagaagt 15850
 ctatcagacatcaaatggaagttaaactcgatatcttaacaattactaggat 15900
 gggcgagtgactcacacctgtaatcccaacactttgggaggctgaggca 15950
 ggaggatcacttgagcccaggagttcgggaccagcctgggcaacatagag 16000
 agacgttgtctctatttttttaataatttaaagagaaaaaataactgaaa 16050
 tattgtatacaccactgaattataataatgtgtatataatgtatatattc 16100
 attatgaggaatatttgattatttcatatattatattctttctctgtt 16150
 tattttatccagttatgaagtatttagaacaattcatcagtaattggggc 16200
 taaattgacagaatagtaatacagagaaaaatagaaaaagacagatgggtta 16250
 tctttgaataaccaggttggagttgtttatgggtttgtttttgttttggg 16300
 ggcggttttttagacagagtcccactctgttgcccaggctggagtgagct 16350
 ggcacaagcatggcccactgcactccttgacctcttgggctcaagcaatct 16400
 tcccaccttagcctcctgagtagctgggaccacaggtgcatgtcaccaca 16450
 cccagctaatttttttttttttggtagagacagctctttctgttatcca 16500
 ggctgatctcaaaactcctgcactcaagtgtacccctgccttggcgctccc 16550
 aaagtattgggattataggcatagccaccaaccccaacctagtttctatt 16600
 tagacttggccctttcccaaccagctcatttgtgtccaaaagatctcataaa 16650
 tgtagacaggaaactgtcctttgtctcatcagttttcttcatcctgtgtct 16700
 agggggatgggtcggtgggggaaactggggttatgcaagttcctctgaaac 16750
 atcctctgtgagcccagggtggtgaggaaccagccgagcagagtcag 16800
 tgtgcagctttccagaaaggaagtcacagccagtcagccggccctggca 16850
 gccagcaccgggcaaccctgctgtctgtgtgataaagaaatggtctgcctg 16900
 acaggatgggtgtggatttttctttttcttttttttttttttgagacagg 16950
 gtctggctctgtcggccaggctggagtgcaatggcgggatcttggctcac 17000
 tgcagcctctgcctcccaggctcaaggcatcctcccacctcggctctccc 17050
 agtagctgggaccagggcacacacccacacgcccactaagttttcgta 17100
 ttttttagtagaggcagggttttactatgttgtccaggctagtctcaaact 17150
 cctgagctcaagctatccatctgccttggcctcccaaagagctggaatta 17200
 caagcgtgagccaactgtgcctgaccagggtggatttttcaagtgcacat 17250
 gttgtgtgtcccaagagctctgatggtaacaaattccaagcgaaaaaagt 17300
 caatggttcccacccatcctacctcccatgatggcaagaggaaatcacca 17350
 cactgcagatacagtcctatgtaaaacaaattgctatggattttgaaagt 17400
 aaccttaagagaactgcactatgttttcttcattagagttctctggtaat 17450
 ttccagcttttttttttttttttttagacagtgctcgctttgtcgccc 17500
 agtgtcaccaggctggagtgagtgacgtgatctcggtcactgcaacc 17550
 tccgctcgtgggttgaagtgtattctcctgcctcagcctcctgagtagct 17600
 gtatttttagtagagacgaggtttcaccatttggccaggctggtctcgaa 17650
 tcttgacctcaagtgtatcgcccatctcagcctccaaagtgcgtggatt 17700

Fig. 16 (continued)

22/34

acaggtgtgagccactgcacccggccagtaatttcaagcttctgaggagc 17750
cctttgaaattgttaataacttgtagctatgtccaacatatccatgttca 17800
gtgtatgttcgatatttcttaggaaacctgcccttggtgttttctttgt 17850
ggtaattcatgagccggcaaatgtgacatgtgttacagaatatacctttt 17900
ctctgtctctcctacctcataaccagaacttaattatcctgtcttttagtcac 17950
ataaatagctaactaaataaatatatgagatttcagtcctgctcactgtga 18000
aaatagaccttctaaatgatctcttccacttgagATATTGCAAATATG 18050
D I C K Y
GATCCATCCCTCCTGATGTGGAGGAGAAGTTACGGTTGGAATGGCCCTAC 18100
G S I P P D V E E K L R L E W P Y
CAGGAGCAATTGCTACTCCGAGAACACTACCAGAAAAAGTTCAAGAACAG 18150
Q E Q L L L R E H Y Q K K F K N S
CACCTACTCAAGtaagaataagaaggcaccctagagatgttccagcccca 18200
T Y S
aagatatttgaataggttggactcgggcaccaatctagcaagtcctacgg 18250
aagttgtataaagctgaaaaatactgaagcatttcccaaatgggaaatcct 18300
aaactcaaaacttgcttttttgggttttttgggtttgtttgttttcttcat 18350
ctgacattgtcttagtagtcacagaatgaaagataaatcaatcattcatga 18400
tctaacaatgaccttcagtgctctaaaaaactacggagtcaggaaacaa 18450
tgaatatattcctcatgtaaaaataaaatacagacataaaagggcaaaa 18500
catgaacatcatctcagctcttcaaataggatgacattgtcacatggggct 18550
ccagtatgccttgggttagagcatttaagcaggaggccctgagtcactcc 18600
agacagtccttgaccaccaagcagcattctcttttggttcctctgtggct 18650
tttgcaaacacagggttagctcagctacccattagtagttttcagtcac 18700
taaaacagctctccagctcttcaaataggatgacattgtcacatggggct 18750
ttaaagcaagtgaacaagggaaccccccttttttttttttttgagatgga 18800
atctcactcttgtcgccagcctggagtgcaatggcgcaatcttggtcga 18850
ctgcaacctccacctccaggttcaagagattctcctgccttagcctcct 18900
attcattatgaggaatatttgattattcagttcctgtagggttaaagatat 18950
tcccccgatcatattattgattattgagtagctgagattacagggtgcct 19000
gccaccacgaccggctaatttttggatttttttagtagagacagggtttc 19050
accattgttgccagggtccaggctcgtctcgaactcctgacctcagggtga 19100
tccaccacctccagcctcccaagttctgggattacaggcgtgagccacc 19150
actcctggccacaatccttttttaactatgaaataatatttttatctgaag 19200
tttgatgtttatacccaactgagggatgatgttcccatatctcagttaaa 19250
gaaataacctgctcagatacttcaagctcttcttttgacttttgaaaaa 19300
aatgatcttgagattactatactttgttgggttagttaacattatttaa 19350
agtatattatttttaattaatattctttgtaagattttactgtatactacc 19400
tggagttcaatgtatcagatggatttcaaatttatgtacattttttatgt 19450
atatgggtacagaaaaaatgtgatccataagaaatcagaaaaatagcgcat 19500
atgctaataagctaatgttgcctctaaaaaacttatttttgcatttttta 19550
gagggggatatactctgacactttaataagtgttaattaatattgactgg 19600
aatttggcatgaggcaggggccatttcagatcccattaaaggaatgacaca 19650
taccagagaaccacagaagtaaggccacatttgtaataaatcattatagc 19700
tctgctaggagaagaccagttgtattaggttaattaatggatttgctctt 19750
aaaacacatgtcccggaagatataggtgagtcctggggggccgcattaaa 19800
cattataccaatgtatcttacatttctaagaaagttttactactttacag 19850
gatctttctgtttacaaaaatggaaggtttccaaactccaggacttggcttt 19900
catagttcctacaccaggggaaatgccttcctttgtctaactatgcaacca 19950
ggttagttagtgtaagtcaggccaccctgttggaatgctaaaagggtaca 20000
acaaacacagaattttatttgcatttgtaaaacatttgatttctggctcga 20050
aattttcagttttcatgggcacgtcatggaaacagaaatcttctgtgttt 20100
agtttgggcacctactctttagtgtagcaaatatttcagaagccaatagg 20150
ggattccacaaattgttctgaacctgtggctgagactggtaattggctgag 20200
tgacatggggacataccacaaaagaagggttagcaaaaggctgctgagat 20250
aaggacatgttcatgttagctagtggtgcaccttaaaacacatgt 20300
cccaggctgggtgtgtgtgctcacgcctgtaatcccagcacttggggagg 20350
ctgaggcggtggattacctgaggtcaggagttcgagaccaacctggcca 20400
acatagtgaaacctcatttctactaaaaatacaaaaattagccaggcatg 20450
gtggcgggcgctgtagtcaccagctactcaggaggcaggcaggagaatta 20500
cttgaatctgggaggcagaggttgtggtgagccgagattgcgccaccgca 20550

Fig. 16 (continued)

23/34

cgctagcctggcgacaaagtgagactctgtctcaaaaaacaaaaacaa 20600
 aaaacaaacaaacaaaaacaaacaaacaaaaaacgggtatcccagaa 20650
 gatacaggtaagtttttctaacacaggtcctctgtatggtgcgttccact 20700
 taagtagaagatgacaaaaacatttgtcatgagaatatagactcacattt 20750
 taaacctgtttgagcaggaaaaggaagcaatgttacagatgtaattctgg 20800
 gtgtgactgcgaaaaggatgactcccttattaaagtagtcatcctgagt 20850
 agctaactctttgtacttctcttctcctcctgttccctcatcaccaca 20900
 ttcttccgttgctacacccaggccacattggatgctgacatagactta 20950
 catggtacagtccaagggaagatctgccatttttttcaatgtgtcatct 21000
 tggttatcttcattccaaggatctctccactttttatcacagtaagagatg 21050
 agagtctggaaaaggattgggaataagataatgaattgtaagttttaatt 21100
 gttcttcgtatttttggggaaggagtaggctagggtggtccttctgttttt 21150
 ttttgttttttttttaaaagtagatgtggccagacgtgggtggctcacgcc 21200
 tgtaatcccagcactttgagaggctgaggcaggtggatcacttgatgtca 21250
 ggagttcaagaccagcctggccaacacagtgaaaccccgctctttactaaa 21300
 aatacaaaaactagccgggcttgggtggcggtccacctgtagtcccagctac 21350
 tgcagaggtggaggcaggagaatcacttgaacccgggaggtggaggttgc 21400
 agtgagccaagatcatgccattgtactccagcctggcgacagaaacata 21450
 ctctgtctcaaaaaaaaagagaaaaagaaaaagaaaaagaatggatttga 21500
 actcagtcgtcaatagccctctattccaggagatgttacagttgattatgt 21550
 tatagggggtgtataatagaatttcgagctatgtaaattccaagtgcatt 21600
 tggaaagaatgaagaaatggaggaagggttaaagtatgagtgcagcatcc 21650
 aggttttttgaaaaatgctataatctttgttcagggttagtacaaagtgt 21700
 atttagctgtaagggttttttgggtgatttacagacagttttcacatgtgtc 21750
 atttcaaccttggttttatggcgaaaggcatgtgatggtgcttgtcccagg 21800
 actttagatccatactctgaggttctgtcgggcaaagatattaccctga 21850
 tcatattatagctctataagtgggagagttgtgcctggagctcaagtctta 21900
 tgattctgtatccagggtcctctcaacatgattttgcaatataaaag 21950
 cctataatgtgtgactaaagcaggtcactcacccttgtaacagactcta 22000
 gtaatggtactgccacaaacggctgcgtgatattgggcaaagacttacc 22050
 ttatttgaatctcagtttctcctcctagaaaaatgaggggtggaggttaagca 22100
 taggctgatgtaagggttaagcctcactgcccctaaactgtggctctaag 22150
 atccagtagaatgtcgggtcacaggactctaggagcttttcaaacccaa 22200
 atgtctgtcattccttgatggttaggcagcagtttatggaagtgggcgaca 22250
 cagcaaatatcaaaatacctaaagcagcttgcaagagttgtttctgccta 22300
 gtggtctttatagtttaatttaaatagtttaatttttttttttttgagac 22350
 agagtcttgcctctgttaccaggctgcagtgagtgccacaatctcggt 22400
 cactgcaacctccacctccgggtttgagcaattctgtctcagcctccca 22450
 agtagctgggactacaggtgcatgccactgcaaccagctaatttttgtat 22500
 ttttagtagagcgggtttcaccatattgggcaggctggtctcgaactc 22550
 ttgacctcaggtgatccacctgcctcagcctcccaaagtgtgggattac 22600
 aggcatgagccactgcaccagcttaaatagctaattttaattatttc 22650
 tatagttattcaagtaattcaggccaaagacttagaaaaacaaacaaag 22700
 ccacttttaaggagaaagggtgtaagtttgccagatagatagagatcttt 22750
 ctttttttaactacaagagttcagggaatgaattactctttaacaaacgact 22800
 atagatatacatgaaaaattggaaggacttattatgcatatgataatcaat 22850
 ttaaagacaacacttaaaattatattgttgccactctcaaaaagtggtaa 22900
 tagaacagctaattggtttaaaaagcagagtacagaagttcccaacttat 22950
 ggcaccttaatatcgcagaaaaactttttaagcatgcctaggccacaaa 23000
 aatacctgtattttgattattaaattgtaaggtctacacaacctaatagt 23050
 aataggtccaatagtaattgtgtccaatagatgttgatgttttttccct 23100
 gcaaaccttaaaagatcctacagtgccctctgtaaatagcactgcctggtta 23150
 gagttgaatttcagataaataattttttcatgttaattatttttctttt 23200
 ctttactttttttttgtttttttgttttttttttttttgagaca 23250
 ggggtctcattctgttgcagggtgctgtgcaatggcatgatcatggctc 23300
 actcagccttgacctcctcagggtgatcctccacctcagcctc 23350
 ccaagtagctagctgggactacaggtgcttaccatcatgcccggttaatt 23400
 tttgtgtttttgtagagatgtggttttgccatgttgccagggtggtct 23450
 tgaactcctgggtcgaagtgtccgcccgcctcggcctcccaaagtgtca 23500
 ggtgacaggtgacgtacactgacactggccctggcggaagtatttctt 23550
 aatggttacataggtacatacactaaacattattttgtctatatgaagt 23600

Fig. 16 (continued)

24/34

tcaagtttaactagggtgccctgcaacttttagttgctaaatcctgtagctg 23650
 taccatgcatctcactgggtgctccccagcttgccctgacagagtttggg 23700
 aaccatagtcctataactctaggccaatttttttaattgtaaaatttgattc 23750
 attttaaatataataaataataacaggaatttttttaaaaattgttttaaa 23800
 tataattaaaattatcaaaatatttttaactgaacttgtagactagagat 23850
 attttagattatgaagagtgggtttatgctaactaatgacagctctggcta 23900
 tgcattgtggagcactgagctataaattgtggcttcccaattctcctgat 23950
 gtcaactgaacaaaacctaaagtgtcagaccagagcttctggatcttcca 24000
 tgggatttcattcaacagctggagcaaatgaagtacagattgatttttttt 24050
 aatttgtccaattttgtgtgtctcaaaaacataattataatcatttattag 24100
 aactagaattttctcagttttaacaacagaaatagttattcattatgaaaa 24150
 gcgaatctggaggccttctgtgtgggtgccaatctaaccattaaattgtga 24200
 cgtttttcttttagGAAGCTCTGTAGATGTGCTATACACTTTTGCAAAC 24250
 R S S V D V L Y T F A N
 GCTCAGGACTGGACTTGATCTTTGGCCATAATGCGTTATTAGAAGACAGCA 24300
 C S G L D L I F G L N A L L R T A
 GATTTCAGTGGAAACAGTTCTAATGCTCAGTTGCTCCTGGACTACTGCTC 24350
 D L Q W N S S N A Q L L L D Y C S
 TTCCAAGGGGTATAACATTCTTTGGGAAGTAGGCAATGgtgagtacccca 24400
 S K G Y N I S W E L G N
 gggaacaattcattaataaggagattccccactagcattatttcttttct 24450
 tttctttttcttttcttttttttttttttttgagacagagctctcgactgc 24500
 tggccaggctggagtgcagtgccgccaacctcggtcacttgaagctctgc 24550
 ctcccaaaacggcattctcctgctcagcctcccgagtagctgggactac 24600
 aggcaccgcgcaaccgcgcccggctaattttttttttttttttttttttt 24650
 tttttttgcattttttagtagagacgggtttcaccgtgttagccaggatg 24700
 gtcttgatctcctgacctcgtgatctgcccctcctcgccctcccaaagtgc 24750
 tgggattacagccgtgagccagcggcggtcagcattatttcttatga 24800
 caacttttttttttttttgagacggagtctcgctctgtcgccaggctgg 24850
 agtgagtggtggcccatctcggtcactgcaagctccacctcccagggttca 24900
 cgccattctcctgctcagcctcccgagtagctgggactacacgcacccg 24950
 ccaccacgcccggcattttttttgtatttttagtagagacgggttttca 25000
 ccgtgttagccaggatggtctctatatcctgaccccatgatctgcccgc 25050
 tcggcctcccaaagtgtgggtattacagggctgagccactgcgcccggcc 25100
 aacactctttttattattagcaaatatacttctgctgggacattcttg 25150
 caagtgtctcaacaatgcaacttttggagtgcatgtggcagaaactcctg 25200
 ctgtatttattccagaacctattattgttaattccagttttatgtttacatt 25250
 tgaagtgagaaccagttggagccagcaacgttcccagctccaaagttccc 25300
 ttgagattttcagaatcacttaaccctattatgcttggcaacctggactc 25350
 agcaaaactgggagtcagcagtttgttttattcatccctctcttctca 25400
 gtttctcaaagtgtcagttaatctcagtaacccccattgcaaccttcatt 25450
 acctgccaagcgtctcagaacttgccagtatagaatcctacgtgggtca 25500
 agctcctgactgtctccttcttcaactcttttttgcaaagaacttgtaa 25550
 ttttaactataagttatcatgattcgccacatttattcaaaacatagagt 25600
 gctttttccacatatcagccaatggaaataaggattaaatgggaaatgaa 25650
 atgtagtaataaggataagcacagctcttctcctgctcaaaacttttttt 25700
 ttttttttttcagacaagatcttgctctgttaccaggctggagtgcagt 25750
 ggcgtgttcatagtctcaatgtaacctccaactcctgggctcatgcaatct 25800
 ctcacacctcagccccctgattagctaggactacactatgcctagccaat 25850
 tttttttcttttctgtgtgtgtgtgcccaggctgtctcgatctcctggc 25900
 ctcaagtaatcctcctgctcggccttctaaaagtgtgggattataggca 25950
 tgagccaactgtgccgggtctcaaacctttttttccaaagtaaatgaagtt 26000
 attagatatggaatatagtctagttcccagatatccatatccattgggttt 26050
 attaccctcattattaaacttcaaatgttttaatagacctcatatctcag 26100
 ttatacagttaaaaattttgttttgttttctggagtatcttatttataa 26150
 ctatgagtttttactttacttttatttttattttttgagacagacgcttg 26200
 ctctgtcactcaggtgtggagtgcgggtgtcgtgatcatggctcactatggc 26250
 ctgcaccttctgggtcgaagtgtacctctccctcagcctcccaagctgag 26300
 actacaggcatgcaccaccacatctagctaatttttttttttcccatgg 26350
 aacaaggctttactatgttaccagagtggtctcaaacctcctggcctcag 26400
 gggatcctcctgtctcagcctacaaaatgctgggattacaggcatgagc 26450

Fig. 16 (continued)

25/34

catagcgccagacctgggttttacttttcttgactttgaattacaagtttt 26500
 tgtaatttggaatgttttgggttttactgtgtatgtttgct 26550
 tttaaatacaacatttctcgatataatatttgagaattgctgtctttcag 26600
 AACCTAACAGTTTCCTTAAGAAGGCTGATATTTTCATCAATGGGTCGCAG 26650
 E P N S F L K K A D I F I N G S Q
 TTAGGAGAAGATTTTATTCAATTGCATAAACTTCTAAGAAAGTCCACCTT 26700
 L G E D F I Q L H K L L R K S T F
 CAAAAATGCAAACTCTATGGTCCTGATGTTGGTCAGCCTCGAAGAAAGA 26750
 K N A K L Y G P D V G Q P R R K
 CGGCTAAGATGCTGAAGAGgttaggaactagaggatgcagaatcactttac 26800
 T A K M L K S
 ttttcttcttttttcttttggagacagagtctcactctgtcagccagactg 26850
 gagtgcagtggtacaatcatggctcactgcaacttcgacctcccaggctc 26900
 aagcaatcctcccatctcagtcctccacaaatagctgggactacaggtgcac 26950
 atcaccacacctggctactttaaaaaaattttttgtagagatggggtct 27000
 ccctgtgttggccaggctgggtctcttgaattcctgtgtcaagccatcct 27050
 tccacctcagcctcccagagtgcaggattacaggcatgagccaccacac 27100
 ccagccaccacttttcttaaaaaaaagattctctctggtagacaa 27150
 tctcaatagttccacatgttattaaacaatctgctgcctgaatacatgat 27200
 ttacaaaaaaaggaaattttgacgggttcagaatatcaagggatctgag 27250
 gcaaatgtcacctatgataaaatttgcctatcaaaattaggaagtgtgt 27300
 ttacctgatcctaaagcagtaaccagcccatttctaggggaataaaactct 27350
 catgcgtatattgtgcatatataatgtattatgactgagtgataataaa 27400
 attttttttctagCTTCTGAAGGCTGGTGGAGAAGTGATTGATTCAGTT 27450
 F L K A G G E V I D S V
 ACATGGCATCagtaagtatgtctcctattcttaatactaggaaagtaagg 27500
 T W H H
 ctagctttattttattacctagtattcaaaaagttagttcatttaactgcc 27550
 aattgactgcagttcaaataagaaacaaatagtgctcctaagtagcactgt 27600
 actccaatttttaataatataaaaaaaatttttaagttatttttaataatg 27650
 tagtggtttctataaagatcacttttatacagaagaacagtgccaattaac 27700
 ccatggaacatataagtagctaaaaccaattgcttgccaaagaaccagta 27750
 acccaggagtagatgtccttgccactgtgtttttcaagacagagtaact 27800
 gatttctagttactttgcatagaatggactcctcctcataactcccttcca 27850
 tcttggtctttccctagtagaacttctaccttttttagtaacaggtgag 27900
 tgggagaggtaagaaggagaataaggtcagcaattaacctaagaagcagaa 27950
 agtaaaatttgttatttttttctgaatattttctgtgtaatttagCTAC 28000
 Y
 TATTTGAATGGACGGACTGCTACCAGGGAAGATTTTCTAAACCTGATGT 28050
 Y L N G R T A T R E D F I N P D V
 ATTGACATTTTATTTCATCTGTGCAAAAAGTTTTCAGGtaaatagtct 28100
 L D I F I S S V Q K V F Q
 ttttaaaactttttaatgtaaaaccagaatccttattttatagtctagcta 28150
 gttctaaattctataggtatgtatattttacatgtttttctaattttagag 28200
 aacaagcactatgacttatccactgttagttttcccttagcattgggtc 28250
 ttaccccatgtacgtgattagaaatttgaaatatttccaatagccttttag 28300
 tagaattaaactcacatagatgataagaatgggttggttcaacttcatgttc 28350
 cttccacagcctaactatttcaataaaaagaagtttcccaagacctaaatg 28400
 actatgaacataattttataactatataggaggggtgggtctaggaataca 28450
 aagttttgaaatgctgttaactcttcaacaccacagttgaaaccaaggtca 28500
 gcttttttgcaattaccatggatacttttctgttctatagGTGGTTGAGA 28550
 V V E
 GCACCAGGCTGGCAAGAAGGTCTGGTTAGGAGAAACAAGCTCTGCATAT 28600
 S T R P G K K V L G E T S S A Y
 GGAGGCGGAGCGCCCTTGCTATCCGACACCTTTGCAGCTGGCTTTATgtg 28650
 G G G A P L L S D T F A A G F M
 agtgaagcagcgctggccttaggggtcagagtgcagctcttctccatcct 28700
 tctattctgtgaaatagctccccagccaaaagcagatcaaagaccgtt 28750
 tcagtggctgagccccaaattcatgccagattttgcaagaaaatgattt 28800
 actaaagcttgagggacatctttaacaagtgttccaaatataatcactata 28850
 aggatgaattgtttcagaaattttggtcctttaattatggccataaatat 28900

26/34

Fig. 16 (continued)

gtcaagtagtccttactctaagaagtaactgtaaaagaatgcatatag 28950
 ccggatatggttagtccctgtaatcccaataactttgggaggccaaggtgg 29000
 gaggattgcttgagcccaggagtttgaggctgcagtgaattatgatggtg 29050
 ccaactgcactctagactgggcaacagagtggagactgtcttttttttccc 29100
 ctctgtcaccagactggaggggcagtgccacgatctcacctcactgcaac 29150
 ctctgcctcccggattgaagcgattctcctgcctcagcgtcctgagtagc 29200
 tgggactacaggagtatcacccgactgggctaatttttgtatttttagta 29250
 gagacgggggttttgacatgttgcccaggctggctgaaacccatgagctc 29300
 aagtgatctgcctacctcagccttccaaaatgctgggattacggacatga 29350
 gcaccacgcccggccacacccctgtctcttaaaaaaaaaaaaaaaatgcaag 29400
 ttagagcatattacagctttgtctctcaggaggatacttagtgtagtag 29450
 ctataattcatagattcccaagaagtttagagcctaagtagtagggtccc 29500
 accagaggggctatcattaaatttaaagatttgtaaatacatctcattgt 29550
 ccaacaccacaaacttgattgctttaaaataactgggttaggtacatttag 29600
 taactctatttagtgcttttaactctatactgctatatcctcacattgagat 29650
 tttttttctttctcttccatcttcattctttttctctcatcctcattc 29700
 ttataagcctagaatacatcacaaatcctttatgcccattggaagcaagag 29750
 gaaataagaatggagatgtttgttttgccattaaactaaagatctgggggtg 29800
 tcgggggagaagggggatagagaaggagaagtgggaagaggtgtccataat 29850
 agcttaggtgcaattctgcttattttacattttacccccgctgactgcca 29900
 ctttttcttcagccctcacacattgtttgtgcagggaacctcataggacca 29950
 ggaattgtctatagaggtgggaatttgtctcacccgtgaagggtacacctc 30000
 tagcatggttaatagctctcttaggatttgttatcatatggaagatgtaaa 30050
 gggaggagattctggttctgctgctgctgcatgcagtggcatttcat 3100
 ttaaatacttattttataattgatgacactttctggcttctgttaatt 30150
 cctccctcaaagatcaataaaaccagaaccaggcatggtggcatgcacttg 30200
 tggctcctgtaaccaccaacagggttcacctgctgctgcttagatagag 30250
 ccaattatcaagacccgggaattgcaaggagaaagagtaatttatgcag 30300
 agccagctgtgcaggagaccagagttttattattactcaaatcagctctcc 30350
 ccgaacattcgaggatcagagcttttaaggataatttggccggtaggggc 30400
 ttaggaagtggagagtgtggttggtcaggttgagatggaatcacaggg 30450
 agtggaagtgcaggtttctgtgctgctgctgctgctgagggatggcag 30500
 aactggttgggcccagattaccgggtctgggtggtctcaaatgatccacca 30550
 gttcagggtctgcaagatatctcaagcactgatcttaggtttacaacag 30600
 tgaatttatccccagggaacaatttggggagggttcagactcttgagccag 30650
 aggtgcattatccctaaaccgtaattcttaattgtttagtaatttgtt 30700
 agtctgcaaaaggtagacttgttccccaggcaagaagggggttttttcaga 30750
 aaagggtattatcatttttgtttcagagtcaaacatgaactgaatttc 30800
 ttcccaaagttagttcagcctacacccagggaatgaagaaggacagcttaa 30850
 aggttagaagcaagatggagtcagtgaaggtctgatctctttcactgtcat 30900
 aatttctcagttataatttttgcaaaaggcggtttcagctccagctactt 30950
 gggaggctgagacaggaggattaatggagcccaggagtttgaggttgacag 31000
 agagctatgatcacgccactgcactccagcctgggtgacagagtgagacc 31050
 ctgtctctaaataaaataaataaataaataaataaataaataaataaatac 31100
 aagatggtgtgcaattagaattgagcgattttgtttccaaacctcaagaa 31150
 agcttggtcttgcctgctgctccagGTGGCTGGATAAATTGGGCCTGTCAGC 31200
 W L D K L G L S A
 CCGAATGGGAATAGAAAGTGGTGATGAGGCAAGTATTCTTTGGAGCAGGAA 31250
 R M G I E V V M R Q V F F G A G
 ACTACCATTAGTGATGAAAACCTCGATCCTTACCTgtaagtgaacctat 31300
 N Y H L V D E N F D P L P
 tattttcctaattctagtggagtagattaaagtcaactcaggacctctgg 31350
 tgtaacctcctatgaacagtcagtcctctcagtaactagccaaatcatg 31400
 agatgatgaattagaaggagccttagatagcatccaatctaactttttt 31450
 tgtgtgtttgaagagaagaaatcaagagctaggaataactttttaagggt 31500
 aagccatttgcagtatagtgtggaattttgtttaaaaggggataaattgaa 31550
 attttatgactcattatacaagacaaaataagttggattttcaaatgttt 31600
 taaaaagtaaatcaaagttataattgcctacagtagcgaagccttcaaaa 31650
 cattttttatgttatgaaattgtaatttttaaccttaaaatgagccag 31700
 taccatgtgtttgtcttaaaaatctcatgctaagaatttactatgtgtta 31750
 ataactctcaagataatttatgaataaagtcctattttctaactcttctcc 31800

Fig. 16 (continued)

27/34

aactgtatctggtgctaaatcaggaaatgtttcttcccaaaaagcctcgt 31850
ggaagatctgtatgtctaaatatagtcagggaataacagatgtagccc 31900
tgcaagcatgaccttgatttttatagtctaaaatgtcatttgcagatat 31950
ctattttctaaagaataattcctaaaagaattatttgaatggtgtaggaaa 32000
gctaagaaattttgcaaagagcgtagctgaaaataaagctaggcttttg 32050
tggtttgtggatagacttcccaacaaaattgctttttatctatagtgatc 32100
caagcttgtggaacatatagtcattcttttttagaaaattcttagaaaa 32150
gtgatcttgcaaaaatggaatttatctttcccaagtataattctgtcatg 32200
tatagagttaaactaagcatagtaatttcaccagacaaacattccaaaatc 32250
tactcctgacctttttatctcatccaaattttccagggeccagacataa 32300
acctttgccttacgaactctttgtatatgcactaaaatagcttctccttc 32350
aaggttctcagtcagctagaaaaatgtgcaagagtaaatggtacccttct 32400
cactttagatccaagagaattagacttaaaactcactctacatgtctgtg 32450
actttattttatgtcatgacagtcctgtgaggtggcaaggcaggtatct 32500
tggatccatttttttagataaggaagttcaaattgagaagaggttgcata 32550
tttacaggaagccatactgtagtctctatgttactcttaaaaatcccatc 32600
aaatcctgcttctgaggcctgcatactttctaccctaccagtcattgacc 32650
catgcttatgtctcctttgaaaacattgattccactcttgtctccagtga 32700
aaaagtggaaattttaagcagagaaaacaaaagccatttgccttggtaagtct 32750
actttccctctactttcaagaaggaaagttggggatgtgttgatggtg 32800
attttattttattttatttttttaaaaattgatacaaggtcttactgta 32850
ttgtgcaggctggtctcaaaactcctgggctcaagtgatcatcccacctca 32900
gctcccagtggttgggattacaagcatgaaccattgtgcccaccaccgatc 32950
cgcagtttttaagaaaaacttttactatagaaaattttaatcatataca 33000
aaatacagaggaagatatgaacccactttaggagactagaataatgcca 33050
ccccaaaaatagccactttggcataaggattatttcgagctaaaggcaac 33100
tgggaagaaaacacataagaagaaaagttctctgtccttctccatttgcta 33150
aaagcaggacatgaatctttaaagtcctccctccttccctttctaccagga 33200
aaaacaagagtttaactcactgaagataaacttcagacccttatcagtgtaga 33250
gatggcactagaagaatctataattacatactcatttattttccttccac 33300
aacttgcaccccgagagataaaaatccttttcccttggtcagtctcttg 33350
tccaaaaatttgcctctataagctggagttctaaagccacctctttgagaat 33400
tacttgttccctggtattttctgttaacatacatgtatttaatacatgt 33450
taacaagcttctgtttgttttctcctgttttctgtcttgttacagaggt 33500
ccatcccaactaagaactaagagtaggaggaataataatttctcctctg 33550
catactttgatcttgttttaatccgtaacccttccacttttccacctcta 33600
cctatttagattactttgaagcaaatttcagatatattactttatctataa 33650
ataattcagtagtggtgtaggtgtggtggtcacacctgtaatcccaacac 33700
tttgggaagctgaggcaggagatcacttgagcccaggagttcaagacca 33750
gtacggcaacaaaaaatcaaaaacttatctgggcatggtggcacatgcc 33800
tgtgtcccagctacatgagaggctgaggcaggaggtcgcttttagccca 33850
ggaggttgaggctgcagtaagctgcattcacaccactgcactccagcctg 33900
ggtgacagagtaagaccatgtctcaaaaaatacatatttttagtatgtat 33950
cctttttgtaaaaacacaatacttttatcatactttaataataacaata 34000
attccttagtatcaccaaaatttttgcagtgctcacattttccttatt 34050
gtctaaaaatattgttgatagttattcaaatcagaatccaaacaaggtcca 34100
tatattacatttgggttgacaagtctcttaagtttgttcattcttaagttc 34150
ttcctccctctctttcatctcttgttaatttattaatgtgaaaaaacaggt 34200
aatttgttctatagtatttcctacattatagagtttgcacattttattcc 34250
ctatgatatacatttagcatgttccctctgtccctgtgtttcctgtaaaact 34300
ggtagttatacctagaaagcttgagtttattcaggttttttaattgtatttt 34350
ttttgcaagaattctttattatctgtctctggaagcacagaatgtctggt 34400
tgtgtctggttttgatcttgacagctactgatgaccattgcctaataccat 34450
tactttattgggtggggggaataaggttttaaaaataaatttttttaaa 34500
gatttttttaactgttatttttgagacagtgctctcatttctgtttccagge 34550
tggagtgagtgccacaatcacggctcactgcagccttgacctcctggga 34600
tcagggtgatcttctcacctcagcctcctgggtacctggaactacaggtgc 34650
acaccaccacacctggctaatttttgtattttgtgtacagaaggggttt 34700
catcatgtttccagactggctctgaaactcctgggttcaagtgtactacc 34750
cacttcagcttcccaaatcctgggattacactttggccaccgtgcctgg 34800
cctaaatgaaattatttgcctctaaacagacagaagttttactttaaaaa 34850

Fig. 16 (continued)

28/34

tttgtctttgtgtgtacatgtgtttgtgtatgtgtgtgtgtctaaaagtt 34900
 tggcttttgagctttgtctttgaattcttggatgaacaataaccaagaatac 34950
 ttaaactctgacattcttgcagatatccctacaggctatggcctttt 35000
 gaattgtgtcctccagtataaaaaagcagcaagcagatactgctctcag 35050
 attcatgggtgggcacatgtgaggtgaaaaaaaaaaaaaagatgaatccta 35100
 tttaaatgccccaggataaacagtgatactctttgtaggataactatttg 35150
 ctggccactggtttcatataaaggacataaagtaaatctatttttgt 35200
 ctcttctccccaaccaccacaactagGATTATTGGCTATCTCTTCTGTT 35250
 D Y W L S L L F
 CAAGAAATTGGTGGGCACCAAGGTGTTAATGGCAAGCGTGCAAGGTTCAA 35300
 K K L V G T K V L M A S V Q G S
 AGAGAAGGAAGCTTCGAGTATACCTTCATTGCACAAACACTGACAAgtaa 35350
 K. R. R. K. L. R V Y L H C T N...T. D N
 gtatgaaacacaccctttaccaatcatcaagtttttagtgggtaagcctgt 35400
 aactttactcaaacaccctgttgcatgtgtctatacattgcataagtata 35450
 ggcagttgcaatttttagtaaaagttttatacaacgattttattttattttat 35500
 ttttagaagaaaaatgctacttttgttgttgttgttttttgagacggggc 35550
 ctgcctcgctcaccaggtgagtgagtgagtgcaatctcagctcactgc 35600
 aacctccgcctcccggttcaagtgaattcttgaaggagagaacaataata 35650
 acaacaatatatttttcaaaagttgtgaccgcagtttctggagttgagaa 35700
 gacatcgagattttttagcctcatactcttgccttttaggtagcaaaaaat 35750
 gttcctaaatctcaggaatatctctagataggttcaatctatcattcc 35800
 tgataagatgatgtgaaataactaattctagccaaaaaagaccagctacc 35850
 atttccgattgttggggactgggaactctggatagtggagcccccagtag 35900
 gaagtagcgagggggaatggtttgaatggataaattcataaaaaatgtcag 35950
 tagatttaattttcttatacatttcagtcctttttataaggctaggaaaag 36000
 ccctgtttttatgggtttataatttgaattcacatgaaccacaaaaattt 36050
 gccttttaccttctatgtctgaaatggatagtctggctggcctcttaa 36100
 caaccagctggcagagctgtgaggatctcagtggtctctagcccagaca 36150
 ttggtagcatgaacggcaacatttttaattgtgttttcaaaataggagca 36200
 cactagcgggtctaaaacgatcataaaagaaggataactaagaggggccact 36250
 gtcatattggatcctaatacttaggatgcattatggattgtcattatgga 36300
 tactaatacttaggatcacatttgaattgagtttttaattgcttaaat 36350
 agatacatatttctattaagtttaacctctttgcttttagTCCAAGGTATA 36400
 P R Y
 AAGAAGGAGATTAACTCTGTATGCCATAAACCTCCATAATGTCACCAAG 36450
 K E. G. D. L. I. L Y A I N L...H N...V T K
 TACTTGCGGTACCTATCCTTTTTCTAACAAGCAAGTGGATAAATACCT 36500
 Y L R L P Y P F S N K Q V D K Y L
 TCTAAGACCTTTGGGACCTCATGGATTACTTTCCAAGtaagtaattttcc 36550
 L. R. E. L. G...P H G L L S K
 ttgttcattccaaactttcaataaatttattgggtgtttatcagaatagag 36600
 agtttggacagggagcaaaagacaaagtcaactatatcaagttctaataa 36650
 ttcttaatatccaggaaatttatgtatgaatacttactaatatgagtata 36700
 actcatcctaagagtctaaagcaaaaggatgtgaacacaaactagcagtt 36750
 atcttagagaataagtttgcattttcaaaataaacttgacatatcaagatcc 36800
 actcaacgcatttaattatttactctaaaaagacataattcttggtaac 36850
 acattcactaaagcaaaatatacctttatataattgctatcaaaggtag 36900
 tgggttgggtataaaaatatcataccatgtgagatcagtggtgattcctttac 36950
 agcattaatttttattgggttagagtaagaaaaagaatagctagagtatat 37000
 ttcttaagtagattctcatacactttgggtttcaaaaaccaattattgact 37050
 acatcttataaaagcctgtattcaatggagtgcacaaaaatgactatgag 37100
 tcttaaaagagtttaggcataaaatattttaagggtttctgttcaatgtatg 37150
 ttggaaggagttcctttctcatgactattctcatattggagcataaaaag 37200
 agtttacagggttggcgagtggtcatgcctgtaatcccaataactttgg 37250
 gaagctgaagcaggcagatcacttcagcccaggagtttgagaccagcctg 37300
 ggcaatatggcaaaactctctacaaaaatataccaaaaatagccaggcg 37350
 tgggtggtgcatgcctgtagtcccagctacttgggaagctgaggtgggagg 37400
 attgcttgagcccaggggggtcatggctgcagtgagctgtgatgggtgcct 37450
 ctgtcaccagcctgggtgacagagtgaagccctgtctcaaaaaataaaa 37500
 taaataaaaaatgaaggtttacaaaaattctcaccatctcctcccatcttt 37550

Fig. 16 (continued)

29/34

gcaaatgccacataagtgatgtgttccaggactattagcctcggaacctg 37600
 aggcagtagcagtaagcagcgtttctccaaagtcctgtcccccacagacaa 37650
 acattatttacactgggtactgctcttttattttttcccccctctatgcttt 37700
 attttactataactataatcatataacatgtaataggaanaaggcaggggt 37750
 cgggggagagatccagaagtcttcccaagagcctttcccaacatagcctct 37800
 gtagacattttttctttcttcttttttttttttttttttttttctgagaca 37850
 gagtctcactctgttgtccaggctagagtgagtgaggcgtgatctaggctc 37900
 actgcaacctccgctcctgggttcaagcaattctcccacctcagcctcc 37950
 ctagtgtctgggattagagcagcatcaccacgcctggctaatttttgt 38000
 attttttagtagagatgaggtttcaccatgtgggcccaggctggtcttgaa 38050
 tcctgacctcaagtgatccacctgccttagcctcccaagtgctaggatt 38100
 acacgagttagccaccgtgcccctgcccctattacattctgatcacacatt 38150
 tcatgtttttataattggaaactggtaaatatagacaatgttttgttc 38200
 ccctaaattctctttgatgagtataattacttacactcttctgtcttta 38250
 aaattttgcaaaatagtagatccctagataagtttatgagtgcacagtctgta 38300
 cgcttactcatattaatgacctcggagagttaaacaacagtcaccttta 38350
 aaattattactatcattatctttttgaggcggggtctcattctgt 38400
 ctcccaggctggagagtagtgggtgcgggtcacagctcactgcagccaccgc 38450
 tacctgggctcaagtgatccctcctcctcagccttctgagtagctgagac 38500
 cacaggcttatgctaccacacctggctaatttttttaactttttgtagaga 38550
 cgatgtctcattatgttggccaggctgggtcctcaaaactcctaagctcaagt 38600
 gatcttctcagcctcccaagtgctgggattacaggcatgaaaaactgc 38650
 acccagccctaaaaattattagggctcctgcatagtaagacttttaataaat 38700
 atttaaatgaacatctggttttttttaaaaaaaaaatagagacaaggtctc 38750
 actataattgcccgaagctggtctcgaaactcctggactcacgcaatcctgct 38800
 gccttagccgcccgaagtgctgggattacaggcatgaccacacctcatctg 38850
 ggctgagtgaacatatttttaacataaaggccgtattttatatttatctc 38900
 atacattttgcccagcatccccatttccgcccgaatctgttgcttgcta 38950
 tccttccagcttcatcttcatctgaaatttgacaaacatcttctatttctt 39000
 tgtcgtcatgttattgacttccagaatataaaataaaacactatacccaaa 39050
 ttaaaccaccacctcatgtgccagcctgatgtgaaaataatcagcataca 39100
 ttaagcttacccttgatataatgtgtagcatcttttagataaataatacagc 39150
 tgattaagcaatataagccttagtggtataatcttggccatgtacctcat 39200
 ctatctccagcaggattaattccacagtgatcagatttacctttaaaactt 39250
 tgtagcaaaatatacctctccaaaagcatatctaaaacttttgtgtgtact 39300
 cttgcaagttttcttaatttcatgcagaacaggctcttaccactgttagct 39350
 ggagatattttcaagacctatttttgggttcttggttcttgatgtggtca 39400
 tggcatttcccccttactccatctaaaaattgaggtgtacaggctttt 39450
 aaacaaaaccactcatatagactgagtacaactgcaatgcaggcatgct 39500
 aacctctgctacaatcatgggcgtgctattgatattgtcttaagtacaga 39550
 acacagggtgctgactcatttaggtcaaaatgtaaacagtttttctgc 39600
 tcactgagcttaagttaggcagggtgtgagagatttctttaaggaaaac 39650
 aaatatataaatgctacatggaaaaatatctaacattagagaattaag 39700
 taaataaaactaatatactcacaccatggaatcttgtgcagacattaaaat 39750
 tatgtagtggatggatgtttaatgggtgtgagaaaaagttaggatgtgctg 39800
 ggggtggggggaagaatcaagtttttaagaaaatacagtataccatactta 39850
 agtaaaaaaaaaaaaaaagggtatgtacagtcagtggttgccttaatgatgg 39900
 ggatacattccgagaaatgtgtcgatagggtgatcttcatccttgtgtgaac 39950
 atcatagagtgaacttacacaaacctagatggcttagcctactatgtatc 40000
 taggctatatgactagcctgttgcctcctaggctacaaacctgtaaagcat 40050
 gttactgtagcgaatatacaataacttaacacaaatggcaagctatcattg 40100
 tgttaagtagttgtgtatctaaacatatctaaaacatagaaaactaatgt 40150
 gttgtgctacaatgttacaatgactatgacattgctaggcaatagggaatt 40200
 ataattttatccttttatggaaaccacttatatatgcgggtccatgggtg 40250
 accaaaaacatccttatgtggcatatgactgtatacatgtacacaaaaaat 40300
 agatgaaagaatgaatatacatcaaaatatttaaaatggttataatgact 40350
 taggttactttttatcttagtaataataatgatgatagataatactt 40400
 ttatagtggttactatataaaagacactgttataagtggttctacatactt 40450
 tacatgtattacctaaatgatataaaataaactctgacagtaactaatct 40500
 tatacgttctcttttcttttttttttttttttttttttagacagaatctt 40550
 gctctaccaggctggagtgcagggtgcaatctcggctcactgcaacctcc 40600

Fig. 16 (continued)

30/34

gcctcccagggttcaaacgatttctcatgtctcagcctcctgagtagctggg 40650
actacaggcacacaccaccatgcccgctaatttttgtaatttttgggtag 40700
agatggagttttgccatgttggccaggctgatcttgaactcctggcctca 40750
agtgatctgcctgcctcagcctcccaaagtgcctgggattacagggtgtgaa 40800
ccactgtgctcggcctaattcttacaagttttcaatatttaaagagtgtca 40850
actttgttgacaatataaaacataatttgagaaaaagagatataagcatct 40900
tatttagaattatgaaaatatcaatagacctacagccgactaaagctttt 40950
cttcataagctcttgcctatatattgattcgctcctgtgaatatgcattaat 41000
ttgatttaataataagtagtattgataaagaaataacacttttccttaatttt 41050
taagaacgttoaacagtttttaatttgaattccaatagtgaatatcatag 41100
aaaatataaaattttctgtagtttagccaaattgtttttgtttcaccaca 41150
gcattctaccacaaatttcttaataacagtaagaaaatgaatgcatacctc 41200
ctgcaggagagggggagtttaggcagtttatgggcatagttacaagtgtgaga 41250
aatttctattggctaccattttagcgtaaattcataaaaaactgcattcaatt 41300
ctatatatctattttctttacataaaaaaggtttcaattattggccatta 41350
aataaaatagccaccattccagaagttgtgtcatgtttatcctttttata 41400
ccaccatcatatttgccctattatataagattgtgtgtgttccattttctgta 41450
atggggccagacagtaagtagtttctggctttggagtcctatgggtctctat 41500
cataactactcatctctgccattgtagcttaaagattatctagggtcaaat 41550
gcctaagtgtatagtggtgaataacaagttatataatataggtgccac 41600
aaaaaaaattttatttgggtctaaaaaagatttcatgacttttgtagcagc 41650
atgggtggggcatgcaccacttggttaactcggtgtatctttctcctttg 41700
cagATCTGTCCAACCTCAATGGTCTAACTCTAAAGATGGTGGATGATCAAA 41750
S V Q L N G L T L K M V D D Q
CCTTGCCACCTTTAATGGAAAAACCTCTCCGGCCAGGAAGTTCACTGGGC 41800
T L P P L M E K P L R P G S S L G
TTGCCAGCTTTTCTCATATAGTTTTTTTGTGATAAGAAATGCCAAAGTTGC 41850
L P A F S Y S F F V I R N A K V A
TGCTTGCATCTGAAAATAAAATATACTAGTCCTGACACTGaatttttcaa 41900
A C I *
gtatactaagagtaaaagcaactcaagttataggaaaggaagcagatacct 41950
tgcaaaagcaactagtgggtgcttgagagacactgggacactgtcagtgct 42000
agatttagcacagtaattttgatctcgctaggtagaacactgctaataata 42050
atagctaataataaccttgttccaaataactgcttagcattttgcatgtttt 42100
acttttatctaaagttttgttttgttttatttatttatttatttattt 42150
ttgagacagaatctctctctgtcaccaggctggagtgccatgggtgcgat 42200
cttggctcactgcaactttaagcaattctcctgcctcagcttccctgagta 42250
gctgggatttataggcggtgtgcccacacgcccagctactttctatatttt 42300
tgtagagatggagtttcgccatattggccaagctgggtctcgaactcctgt 42350
cctcgaactcctgtcctcaagtgtatccaccgcctcagcctctcaaagtg 42400
ctgggattacaggtgtgagccaccacacccagcagtggtttatttttgag 42450
acaggggtatcattctgttggccaggcttgagtgagtggtgcaatcatag 42500
atcactgcagccttttaactcctgggtcaagtcactcctcctgcttagcc 42550
tcccaagtagctaggaccacagacacatgccatcacacttggtattttt 42600
aaaaaattttttagagatgggtctcgctatgttacccaaactgggtcc 42650
tgaaactcctggactcaatttgatctcccaccttggccttccaggtgctgg 42700
gatttctttgggagtacagcatggtacagcaggagatcatttgatgttac 42750
ctctgtgcagtggttgctagtgcagaaagactataatacctgtggggaca 42800
gcgattagccaccacaaccagtctttatttaaagttattaaaaatggctg 42850
ggcgagtggtctcacacctgtaactcctagcactttgggagggccgaggcag 42900
atggatcacctgacgtgaggaatttgagaccagcctggccaacatgggtga 42950
aacccccatctctactaaaaaaatacaaaaaattagctgggtgtggtcctgta 43000
gtcccagctacttgggaggctggggcaggagaattacttgaaccaggag 43050
gcagaggttgcatgtgagccgagatttggtccactgcactccagcctgggtg 43100
acagagagagattccatctcaaaaaacaagttattaaaaatgtatatga 43150
atgctcctaataatggtcaggaagcaaggaagcgaaggatatattatgagt 43200
tttaagaaggtgcttagctgtatatatttctttcaaaatgtattagaaga 43250
ttttagaattctttcttctcatgtgccatctctacaggcaccatcagaaa 43300
aagcatactgcccgttacgtgaaactggttgtaaaagagaaactatctat 43350
ttgcaccttaaaagacagctagattttgctgattttcttcttctcggtttt 43400

Fig. 16 (continued)

31/34


ctttgtcagcaataatatgtgagaggacagattgttagatatgatagtat 43450
aaaaaatgggttaatgacaattcagaggcgaggagattctgtaaacttaaa 43500
attactataaaatgaaattgatttgtcaagaggataaaatttttagaaaaacac 43550
ccaataaccttataaactgtctgttaaatgcttgctttttctctacctttctt 43600
ccttgttttcagttgggaagcttttggctgcaagtaacagaaactcctaatt 43650
tcaaattggcttaagcaataaggaaatgtatattccacataactagacgt 43700
tcaaacaggccaggctccagcacttcagtacgtcaccagggatctgggtt 43750
cttcccagctctctgtctgtccatctttagcgctggttcattctcagac 43800
tctggttagcatgatggctgtagctgtttcatgggccccttcaaacctcat 43850
agcaaccagaggaagaaaatgagccattttttgagtctccttcatagact 43900
tgaataactctttttcagagcttctcagcaaaacctctcctcatgtctc 43950
ctcatgtcttattgttcagaaatgggtaattgtggccatttcaccagtcac 44000
tgccaacaacaacgaggttcctataattgtctctgagtaaccctttggaa 44050
tgagaggggtgttggctcagtcctacaaactgaacactgcagttctgcgctt 44100
tttaccagtgaaaaaatgtaattattttcccctcttaaggattaatattc 44150
ttcaaatgtatgcctgttatggatatagtatctttaaattttttatttt 44200
aatagcttttaggggtacacactttttgcttacaggggtgaattgtgtagt 44250
gggtgaagactcggcttttaattgtacttgtcacctgagtgtgtacattgt 44300
acccaataggtaatttttcatccattaccctccttccgcccctcttccctt 44350
ctgagtctccaacatcccttataccactgtgtatgttcttgtgtacctac 44400
agctaagcttccacttataagtgaagaacatgcagtatttggttttccatt 44450
cctgagttacttcccttaggataacagccccagttccgtccaagttgct 44500
gcaaaaatacattattcttctttatggctgagtaatagtccatggtacata 44550
tataccacatttttctttatccacttatcagttgatggacacttaggttaa 44600
ttccattcaatttcatccaatttaagtataatttgtaaggagctaaagctg 44650
aaaattaaatttttagatctttcaataactcttaaattttatatgtaagtgg 44700
tttttatattttcacatttgaaataaagtaatttttataaccttgatatt 44750
gtatgactattcttttagtaatgtaaagcctacagactcctacatttgga 44800
accactagtggtgtgtttcacccttggtatactatcaggatcctcga 44898

32/34

Figure 17

					50
human	MLLRSKPALP	PPLMLLLLG	LGPLSPGALP	RPAQAQDVVD	LDFFTQEP
mouse	~~~~~ML	RLLLLLWLG	LGALAQGAPA	GTAPTDDVVD	LEFYTKRPLR
rat	~~~~~	~LLLLLWLG	LRALTQGTPA	GTAPTKDVVD	LEFYTKRRLFQ
					100
human	LVSPSFLSVT	IDANLATDPR	FLILLGSPKL	RTLARGLSPA	YLRFGGKT
mouse	SVSPSFLSIT	IDASLATDPR	FLTFLGSPRL	RALARGLSPA	YLRFGGKT
rat	SVSPSFLSIT	IDASLATDPR	FLTFLSSPRL	RALSRGLSPA	YLRFGGKT
					150
human	FLIFDPKES	TSEERSYWQS	QVNQDICKYG	SIPPDVEEKL	RLEWPYQEQ
mouse	FLIFDPKES	TSEERSYWQS	QVNHDIRSE	PVSAAVLRKL	QVEWPFQELL
rat	FLIFDPNNEP	TSEERSYWQS	QDNNDICGSD	RVSADVL~~~	~~~~~
					200
human	LLREHYQKKF	KNSTYSRSSV	DVLYTFANCS	GLDLIFGLNA	LLRTADLQWN
mouse	LLREHYQKKF	KNSTYSRSSV	DVLYTFANCS	GLDLIFGLNA	LLRTADLQWN
rat	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
					250
human	SSNAQLLLDY	CSSKGYNISW	ELGNEPNSFL	KKADIFINGS	QLGEDYIQLH
mouse	SSNAQLLLDY	CSSKGYNISW	ELGNEPNSFW	KKAHILIDGL	QLGEDFVELH
rat	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
					300
human	KLLRKSTFKN	AKLYGPDVGQ	PRRKTAKMLK	SFLKAGGEVI	DSVTWHHYL
mouse	KLLQRSAPQN	AKLYGPDIGQ	PRGKTVKLLR	SFLKAGGEVI	DSLTVWHHYL
rat	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
					350
human	NGRTATREDF	LNPDVLDIFI	SSVQKVFQV	ESTRPGKKVW	LGETSSAYGG
mouse	NGRIATKEDF	LSSDALDTEI	LSVQKILKVT	KEITPGKKVW	LGETSSAYGG
rat	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
					400
human	GAPLLSDTFA	AGFMWLDKLG	LSARMGIEVV	MRQVFFGAGN	YHLVDENFDP
mouse	GAPLLSNTFA	AGFMWLDKLG	LSAQMIEVV	MRQVFFGAGN	YHLVDENFEP
rat	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
					450
human	LPDYWLSLLF	KKLVGTVKVM	ASVQGSKRRK	LRVYLHCTNT	DNPRYKEGDL
mouse	LPDYWLSLLF	KKLVGPRVLL	SRVKGPDERSK	LRVYLHCTNV	YHPRYQEGDL
rat	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
					500
human	TLYAINLHNV	TKYLRLPYPF	SNKQVDKYL	RPLGPHGLLS	KSVQINGLTL
mouse	TLYVNLHNV	TKHLKVPPPL	FRKPVDTYLL	KPSGPDGLLS	KSVQINGQIL
rat	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~L
					543
human	KMVDQTLPP	IMEKPLRPGS	SLGLPAFSYS	FFVIRNAKVA	ACI~
mouse	KMVEQTLPA	LTEKPLPAGS	ALSPLAFSYG	FFVIRNAKIA	ACI~
rat	KMVEQTXPA	LTEKPLPAGS	SLSVPAFSYG	FFVIRNAKIA	ACI~

Figure 18



34/34

Figure 19

```

      |MLLRSKPALPPPLMLLLGLGPLSPGALPRPAQAQDVVDLDFFTQEPHLVSPSFLSVT| 60
PHD |          EEEEE          HHH          EEEE    EEE|

      |IDANLATDPRFLILLGSPKLRTLARGLSPAYLRFGGTKTDFLIFDPKKESTFEERSYWS| 120
PHD |EEE          EEEEE    HHHHHH    HHHHE    EEEEE    HHHHHH|

      |QVNQDICKYGSIPPDVEEKLRLWPYQEQLLLREHYQKFKNSTYSRSSVDVLYTFANCS| 180
PHD |HHHHHHHH    HHHHHH    HHHHHHHHHHHHHHH    EEEEEEEEEEE|

      |GLDLIFGLNALRLTADLQWNSSNAQLLLDYCSSKGYNISWEIGNEPNSFLKKADIFINGS| 240
PHD | HHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHH    EEEEE    HHHHHH EEEE|

      |QLGEDYIQLHKLLRKSTFKNAKLYGPDVGQPRRTAKMLKSFLKAGGEVIDSVTWHHYL| 300
PHD | HHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHH    HHHHHHHHHHHH    EEEEEEEEEEE|

      |NGRTATREDFLNPDVLDIFISSVQKVQVVESTRPGKKVWLGETSSAYGGGAPLLSDTFA| 360
PHD |          HHHHHHHHHHEEEEEEE    EEEEE    HHHHHHH|

      |AGFMWLDKLGLSARMGIEVVMRQVFFGAGNYHLVDENFDPLPDYWLSLLFKKLVGTVKLM| 420
PHD |HHHHHHHHH    HHHH HHHHHHHHHHHH    EEEEE    HHHHHHHHHHHH    EEEEE|

      |ASVQGSKRRKLRVYLHCTNTDNPRYKEGDLTLYAINLHNVTKYLRPLYPFSNKQVDKYLL| 480
PHD |EEE    E EEEEEEE    EEEEE    EEEEE    HHHHHHHH|

      |RPLGPHGLLSKSVQLNGLTLKMVDDQTLPPLMKEPLRPGSSLGLPAFSYSFFVIRNAKVA| 540
PHD |HH    EEEEEEE    EEEEE    EEEEEEE EE|

      |ACI|
PHD | |

```

543

1

SEQUENCE LISTING

- (1) GENERAL INFORMATION:
- (i) APPLICANT: Iris Pecker, Israel Vlodavsky and Elena Feinstein
 - (ii) TITLE OF INVENTION: POLYNUCLEOTIDE ENCODING A POLYPEPTIDE HAVING HEPARANASE ACTIVITY AND EXPRESSION OF SAME IN GENETICALLY MODIFIED CELLS
 - (iii) NUMBER OF SEQUENCES: 47
 - (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: Mark M. Friedman c/o Anthony Castorina
 - (B) STREET: 2001 Jefferson Davis Highway, Suite 207
 - (C) CITY: Arlington
 - (D) STATE: Virginia
 - (E) COUNTRY: United States of America
 - (F) ZIP: 22202
 - (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: 1.44 megabyte, 3.5" microdisk
 - (B) COMPUTER: Twinhead* Slimnote-890TX
 - (C) OPERATING SYSTEM: MS DOS version 6.2, Windows version 3.11
 - (D) SOFTWARE: Word for Windows version 2.0 converted to an ASCII file
 - (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE:
 - (C) CLASSIFICATION:
 - (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: 08/922,170
 - (B) FILING DATE: 2 SEP 1997
 - (A) APPLICATION NUMBER: 09/109,386
 - (B) FILING DATE: 10 JUL 1998
 - (A) APPLICATION NUMBER: PCT/US98/17954
 - (B) FILING DATE: 31 AUG 1998
 - (A) APPLICATION NUMBER: 09/258,892
 - (B) FILING DATE: 1 MAR 1999
 - (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Friedmam, Mark M.
 - (B) REGISTRATION NUMBER: 33,883
 - (C) REFERENCE/DOCKET NUMBER: 910/62
 - (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: 972-3-5625553
 - (B) TELEFAX: 972-3-5625554
 - (C) TELEX:
- (2) INFORMATION FOR SEQ ID NO:1:
- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 27
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:
CCATCCTAAT ACGACTCACT ATAGGGC 27
- (2) INFORMATION FOR SEQ ID NO:2:
- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 24
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single

2

- (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:
GTAGTGATGC CATGTAAC TG AATC 24
- (2) INFORMATION FOR SEQ ID NO:3:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 23
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:
ACTCACTATA GGGCTCGAGC GGC 23
- (2) INFORMATION FOR SEQ ID NO:4:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 22
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:
GCATCTTAGC CGTCTTCTT CG 22
- (2) INFORMATION FOR SEQ ID NO:5:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 15
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:
TTTTTTTTTT TTTT 15
- (2) INFORMATION FOR SEQ ID NO:6:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 23
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:
TTCGATCCCA AGAAGGAATC AAC 23
- (2) INFORMATION FOR SEQ ID NO:7:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 24
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:
GTAGTGATGC CATGTAAC TG AATC 24
- (2) INFORMATION FOR SEQ ID NO:8:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 9
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:
Tyr Gly Pro Asp Val Gly Gln Pro Arg

3

5

9

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1721
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

CTAGAGCTTT CGACTCTCCG CTGCGCGGCA GCTGGCGGGG GGAGCAGCCA GGTGAGCCCA 60
 AGATGCTGCT GCGCTCGAAG CCTGCGCTGC CGCCGCCGCT GATGCTGCTG CTCCTGGGGC 120
 CGCTGGGTCC CCTCTCCCCT GCGGCCCTGC CCCGACCTGC GCAAGCACAG GACGTCGTGG 180
 ACCTGGACTT cTTCACCCAG GAGCGCGTGC ACCTGGTGAG CCCCTCGTTC CTGTCCGTCA 240
 CCATTGACGC CAACCTGGCC ACGGACCCGC GGTTCCTCAT CCTCTGGGT TCTCCAAAGC 300
 TTGCTACCTT GGCCAGAGGC TTGTCTCTCG CGTACCTGAG GTTTGGTGGC ACCAAGACAG 360
 ACTTCCTAAT TTTCGATCCC AAGAAGGAAT CAACCTTTGA AGAGAGAAGT TACTGGCAAT 420
 CTCAAGTCAA CCAGGATATT TGCAATATG GATCCATCCC TCCTGATGTG GAGGAGAAGT 480
 TACGGTTGGA ATGGCCCTAC CAGGAGCAAT TGCTACTCCG AGAACACTAC CAGAAAAAGT 540
 TCAAGAACAG CACCTACTCA AGAAGCTCTG TAGATGTGCT ATACACTTTT GCAAACCTGCT 600
 CAGGACTGGA CTTGATCTTT GGCCTAAATG CGTTATTAAG AACAGCAGAT TTGCAGTGGG 660
 ACAGTTCTAA TGCTCAGTTG CTCCTGGACT ACTGCTCTTC CAAGGGGTAT AACATTTCTT 720
 GGGAACTAGG CAATGAACCT AACAGTTTCC TTAAGAAGGC TGATATTTTC ATCAATGGGT 780
 CGCAGTTAGG AGAAGATTAT ATTCAATTGC ATAACTTCT AAGAAAGTCC ACCTTCAAAA 840
 ATGCAAAACT CTATGGTCCT GATGTTGGTC AGCCTCGAAG AAAGACGGCT AAGATGCTGA 900
 AGAGCTTCCT GAAGGCTGGT GGAGAAAGTGA TTGATTCACT TACATGGCAT CACTACTATT 960
 TGAATGGACG GACTGCTACC AGGGAAGATT TTCTAAACCC TGATGTATTG GACATTTTTA 1020
 TTTCATCTGT CCAAAAAGTT TTCCAGGTGG TTGAGAGCAC CAGGCCTGGC AAGAAGGTCT 1080
 GGTTAGGAGA AACAAAGCTT GCATATGGAG GCGGAGCGCC CTTGCTATCC GACACCTTTG 1140
 CAGCTGGCTT TATGTGGCTG GATAAATTGG GCCTGTCAGC CCGAATGGGA ATAGAAGTGG 1200
 TGATGAGGCA AGTATTCTTT GGAGCAGGAA ACTACCATT T AGTGGATGAA AACTTCGATC 1260
 CTTTACCTGA TTATTGGCTA TCTCTTCTGT TCAAGAAATT GGTGGGCACC AAGGTGTAA 1320
 TGGCAAGCGT GCAAGGTTCA AAGAGAAGGA AGCTTCGAGT ATACCTTCAT TGCACAAACA 1380
 CTGCAATCC AAGGTATAAA GAAGGAGATT TAACTCTGTA TGCCATAAAC CTCCATAACG 1440
 TCACCAAGTA CTTGCGGTTA CCCTATCCTT TTCTAACA GCAAGTGGAT AAATACCTTC 1500
 TAAGACCTTT GGGACCTCAT GGATTACTTT CCAAATCTGT CCAACTCAAT GGTCTAACTC 1560
 TAAAGATGGT GGATGATCAA ACCTTGCCAC CTTTAATGGA AAAACCTCTC CGGCCAGGAA 1620
 GTTCACTGGG CTGCCAGCT TTCTCATATA GTTTTTTGT GATAAGAAAT GCCAAAGTTG 1680
 CTGCTTGCAT CTGAAAAATA AATATACTAG TCCTGACACT G 1721

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 543
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Met Leu Leu Arg Ser Lys Pro Ala Leu Pro Pro Pro Leu Met Leu Leu
 5 10 15
 Leu Leu Gly Pro Leu Gly Pro Leu Ser Pro Gly Ala Leu Pro Arg Pro
 20 25 30
 Ala Gln Ala Gln Asp Val Val Asp Leu Asp Phe Phe Thr Gln Glu Pro
 35 40 45
 Leu His Leu Val Ser Pro Ser Phe Leu Ser Val Thr Ile Asp Ala Asn
 50 55 60

Leu Ala Thr Asp Pro Arg Phe Leu Ile Leu Leu Gly Ser Pro Lys Leu
 65 70 75 80

Arg Thr Leu Ala Arg Gly Leu Ser Pro Ala Tyr Leu Arg Phe Gly Gly
 85 90 95

Thr Lys Thr Asp Phe Leu Ile Phe Asp Pro Lys Lys Glu Ser Thr Phe
 100 105 110

Glu Glu Arg Ser Tyr Trp Gln Ser Gln Val Asn Gln Asp Ile Cys Lys
 115 120 125

Tyr Gly Ser Ile Pro Pro Asp Val Glu Glu Lys Leu Arg Leu Glu Trp
 130 135 140

Pro Tyr Gln Glu Gln Leu Leu Leu Arg Glu His Tyr Gln Lys Lys Phe
 145 150 155 160

Lys Asn Ser Thr Tyr Ser Arg Ser Ser Val Asp Val Leu Tyr Thr Phe
 165 170 175

Ala Asn Cys Ser Gly Leu Asp Leu Ile Phe Gly Leu Asn Ala Leu Leu
 180 185 190

Arg Thr Ala Asp Leu Gln Trp Asn Ser Ser Asn Ala Gln Leu Leu Leu
 195 200 205

Asp Tyr Cys Ser Ser Lys Gly Tyr Asn Ile Ser Trp Glu Leu Gly Asn
 210 215 220

Glu Pro Asn Ser Phe Leu Lys Lys Ala Asp Ile Phe Ile Asn Gly Ser
 225 230 235 240

Gln Leu Gly Glu Asp Tyr Ile Gln Leu His Lys Leu Leu Arg Lys Ser
 245 250 255

Thr Phe Lys Asn Ala Lys Leu Tyr Gly Pro Asp Val Gly Gln Pro Arg
 260 265 270

Arg Lys Thr Ala Lys Met Leu Lys Ser Phe Leu Lys Ala Gly Gly Glu
 275 280 285

Val Ile Asp Ser Val Thr Trp His His Tyr Tyr Leu Asn Gly Arg Thr
 290 295 300

Ala Thr Arg Glu Asp Phe Leu Asn Pro Asp Val Leu Asp Ile Phe Ile
 305 310 315 320

Ser Ser Val Gln Lys Val Phe Gln Val Val Glu Ser Thr Arg Pro Gly
 325 330 335

Lys Lys Val Trp Leu Gly Glu Thr Ser Ser Ala Tyr Gly Gly Gly Ala
 340 345 350

Pro Leu Leu Ser Asp Thr Phe Ala Ala Gly Phe Met Trp Leu Asp Lys
 355 360 365

Leu Gly Leu Ser Ala Arg Met Gly Ile Glu Val Val Met Arg Gln Val
370 375 380

Phe Phe Gly Ala Gly Asn Tyr His Leu Val Asp Glu Asn Phe Asp Pro
385 390 395 400

Leu Pro Asp Tyr Trp Leu Ser Leu Leu Phe Lys Lys Leu Val Gly Thr
405 410 415

Lys Val Leu Met Ala Ser Val Gln Gly Ser Lys Arg Arg Lys Leu Arg
420 425 430

Val Tyr Leu His Cys Thr Asn Thr Asp Asn Pro Arg Tyr Lys Glu Gly
435 440 445

Asp Leu Thr Leu Tyr Ala Ile Asn Leu His Asn Val Thr Lys Tyr Leu
450 455 460

Arg Leu Pro Tyr Pro Phe Ser Asn Lys Gln Val Asp Lys Tyr Leu Leu
465 470 475 480

Arg Pro Leu Gly Pro His Gly Leu Leu Ser Lys Ser Val Gln Leu Asn
485 490 495

Gly Leu Thr Leu Lys Met Val Asp Asp Gln Thr Leu Pro Pro Leu Met
500 505 510

Glu Lys Pro Leu Arg Pro Gly Ser Ser Leu Gly Leu Pro Ala Phe Ser
515 520 525

Tyr Ser Phe Phe Val Ile Arg Asn Ala Lys Val Ala Ala Cys Ile
530 535 540 543

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1721
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

CT AGA GCT TTC GAC 14

TCT CCG CTG CGC GGC AGC TGG CGG GGG GAG CAG CCA GGT GAG CCC AAG 62

ATG CTG CTG CGC TCG AAG CCT GCG CTG CCG CCG CCG CTG ATG CTG CTG 110
Met Leu Leu Arg Ser Lys Pro Ala Leu Pro Pro Pro Leu Met Leu Leu
5 10 15

CTC CTG GGG CCG CTG GGT CCC CTC TCC CCT GGC GCC CTG CCC CGA CCT 158
Leu Leu Gly Pro Leu Gly Pro Leu Ser Pro Gly Ala Leu Pro Arg Pro
20 25 30

GCG CAA GCA CAG GAC GTC GTG GAC CTG GAC TTC TTC ACC CAG GAG CCG 206
Ala Gln Ala Gln Asp Val Val Asp Leu Asp Phe Phe Thr Gln Glu Pro
35 40 45

CTG CAC CTG GTG AGC CCC TCG TTC CTG TCC GTC ACC ATT GAC GCC AAC 254
 Leu His Leu Val Ser Pro Ser Phe Leu Ser Val Thr Ile Asp Ala Asn
 50 55 60

CTG GCC ACG GAC CCG CGG TTC CTC ATC CTC CTG GGT TCT CCA AAG CIT 302
 Leu Ala Thr Asp Pro Arg Phe Leu Ile Leu Leu Gly Ser Pro Lys Leu
 65 70 75 80

CGT ACC TTG GCC AGA GGC TTG TCT CCT GCG TAC CTG AGG TTT GGT GGC 350
 Arg Thr Leu Ala Arg Gly Leu Ser Pro Ala Tyr Leu Arg Phe Gly Gly
 85 90 95

ACC AAG ACA GAC TTC CTA ATT TTC GAT CCC AAG AAG GAA TCA ACC TTT 398
 Thr Lys Thr Asp Phe Leu Ile Phe Asp Pro Lys Lys Glu Ser Thr Phe
 100 105 110

GAA GAG AGA AGT TAC TGG CAA TCT CAA GTC AAC CAG GAT ATT TGC AAA 446
 Glu Glu Arg Ser Tyr Trp Gln Ser Gln Val Asn Gln Asp Ile Cys Lys
 115 120 125

TAT GGA TCC ATC CCT CCT GAT GTG GAG GAG AAG TTA CGG TTG GAA TGG 494
 Tyr Gly Ser Ile Pro Pro Asp Val Glu Glu Lys Leu Arg Leu Glu Trp
 130 135 140

CCC TAC CAG GAG CAA TTG CTA CTC CGA GAA CAC TAC CAG AAA AAG TTC 542
 Pro Tyr Gln Glu Gln Leu Leu Arg Glu His Tyr Gln Lys Lys Phe
 145 150 155 160

AAG AAC AGC ACC TAC TCA AGA AGC TCT GTA GAT GTG CTA TAC ACT TTT 590
 Lys Asn Ser Thr Tyr Ser Arg Ser Ser Val Asp Val Leu Tyr Thr Phe
 165 170 175

GCA AAC TGC TCA GGA CTG GAC TTG ATC TTT GGC CTA AAT GCG TTA TTA 638
 Ala Asn Cys Ser Gly Leu Asp Leu Ile Phe Gly Leu Asn Ala Leu Leu
 180 185 190

AGA ACA GCA GAT TTG CAG TGG AAC AGT TCT AAT GCT CAG TTG CTC CTG 686
 Arg Thr Ala Asp Leu Gln Trp Asn Ser Ser Asn Ala Gln Leu Leu Leu
 195 200 205

GAC TAC TGC TCT TCC AAG GGG TAT AAC ATT TCT TGG GAA CTA GGC AAT 734
 Asp Tyr Cys Ser Ser Lys Gly Tyr Asn Ile Ser Trp Glu Leu Gly Asn
 210 215 220

GAA CCT AAC AGT TTC CTT AAG AAG GCT GAT ATT TTC ATC AAT GGG TCG 782
 Glu Pro Asn Ser Phe Leu Lys Lys Ala Asp Ile Phe Ile Asn Gly Ser
 225 230 235 240

CAG TTA GGA GAA GAT TAT ATT CAA TTG CAT AAA CTT CTA AGA AAG TCC 830
 Gln Leu Gly Glu Asp Tyr Ile Gln Leu His Lys Leu Leu Arg Lys Ser
 245 250 255

ACC TTC AAA AAT GCA AAA CTC TAT GGT CCT GAT GTT GGT CAG CCT CGA 878
 Thr Phe Lys Asn Ala Lys Leu Tyr Gly Pro Asp Val Gly Gln Pro Arg
 260 265 270

7

AGA AAG ACG GCT AAG ATG CTG AAG AGC TTC CTG AAG GCT GGT GGA GAA 926
 Arg Lys Thr Ala Lys Met Leu Lys Ser Phe Leu Lys Ala Gly Gly Glu
 275 280 285

GTG ATT GAT TCA GTT ACA TGG CAT CAC TAC TAT TTG AAT GGA CGG ACT 974
 Val Ile Asp Ser Val Thr Trp His His Tyr Tyr Leu Asn Gly Arg Thr
 290 295 300

GCT ACC AGG GAA GAT TTT CTA AAC CCT GAT GTA TTG GAC ATT TTT ATT 1022
 Ala Thr Arg Glu Asp Phe Leu Asn Pro Asp Val Leu Asp Ile Phe Ile
 305 310 315 320

TCA TCT GTG CAA AAA GTT TTC CAG GTG GTT GAG AGC ACC AGG CCT GGC 1070
 Ser Ser Val Gln Lys Val Phe Gln Val Val Glu Ser Thr Arg Pro Gly
 325 330 335

AAG AAG GTC TGG TTA GGA GAA ACA AGC TCT GCA TAT GGA GGC GGA GCG 1118
 Lys Lys Val Trp Leu Gly Glu Thr Ser Ser Ala Tyr Gly Gly Gly Ala
 340 345 350

CCC TTG CTA TCC GAC ACC TTT GCA GCT GGC TTT ATG TGG CTG GAT AAA 1166
 Pro Leu Leu Ser Asp Thr Phe Ala Ala Gly Phe Met Trp Leu Asp Lys
 355 360 365

TTG GGC CTG TCA GCC CGA ATG GGA ATA GAA GTG GTG ATG AGG CAA GTA 1214
 Leu Gly Leu Ser Ala Arg Met Gly Ile Glu Val Val Met Arg Gln Val
 370 375 380

TTC TTT GGA GCA GGA AAC TAC CAT TTA GTG GAT GAA AAC TTC GAT CCT 1262
 Phe Phe Gly Ala Gly Asn Tyr His Leu Val Asp Glu Asn Phe Asp Pro
 385 390 395 400

TTA CCT GAT TAT TGG CTA TCT CTT CTG TTC AAG AAA TTG GTG GGC ACC 1310
 Leu Pro Asp Tyr Trp Leu Ser Leu Leu Phe Lys Lys Leu Val Gly Thr
 405 410 415

AAG GTG TTA ATG GCA AGC GTG CAA GGT TCA AAG AGA AGG AAG CTT CGA 1358
 Lys Val Leu Met Ala Ser Val Gln Gly Ser Lys Arg Arg Lys Leu Arg
 420 425 430

GTA TAC CTT CAT TGC ACA AAC ACT GAC AAT CCA AGG TAT AAA GAA GGA 1406
 Val Tyr Leu His Cys Thr Asn Thr Asp Asn Pro Arg Tyr Lys Glu Gly
 435 440 445

GAT TTA ACT CTG TAT GCC ATA AAC CTC CAT AAC GTC ACC AAG TAC TTG 1454
 Asp Leu Thr Leu Tyr Ala Ile Asn Leu His Asn Val Thr Lys Tyr Leu
 450 455 460

CGG TTA CCC TAT CCT TTT TCT AAC AAG CAA GTG GAT AAA TAC CTT CTA 1502
 Arg Leu Pro Tyr Pro Phe Ser Asn Lys Gln Val Asp Lys Tyr Leu Leu
 465 470 475 480

AGA CCT TTG GGA CCT CAT GGA TTA CTT TCC AAA TCT GTC CAA CTC AAT 1550
 Arg Pro Leu Gly Pro His Gly Leu Leu Ser Lys Ser Val Gln Leu Asn
 485 490 495

GGT CTA ACT CTA AAG ATG GTG GAT GAT CAA ACC TTG CCA CCT TTA ATG 1598

8

Gly Leu Thr Leu Lys Met Val Asp Asp Gln Thr Leu Pro Pro Leu Met
 500 505 510

GAA AAA CCT CTC CGG CCA GGA AGT TCA CTG GGC TTG CCA GCT TTC TCA 1646
 Glu Lys Pro Leu Arg Pro Gly Ser Ser Leu Gly Leu Pro Ala Phe Ser
 515 520 525

TAT AGT TTT TTT GTG ATA AGA AAT GCC AAA GTT GCT GCT TGC ATC TGA 1694
 Tyr Ser Phe Phe Val Ile Arg Asn Ala Lys Val Ala Ala Cys Ile
 530 535 540 543

AAA TAA AAT ATA CTA GTC CTG ACA CTG 1721

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 824
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12

CTGGCAAGAA GGTCTGGTTG GGAGAGACGA GCTCAGCTTA CGGTGGCGGT GCACCCCTGTC 60
 TGTCACACAC CTTTGCAGCT GGCTTTATGT GGCTGGATAA ATTGGGCCTG TCAGCCCAGA 120
 TGGGCATAGA AGTCGTGATG AGGCAGGTGT TCTTCGGAGC AGGCAACTAC CACTTAGTGG 180
 ATGAAAACTT TGAGCCTTTA CCTGATTACT GGCTCTCTCT TCTGTCAAG AAACCTGGTAG 240
 GTCCACGGGT GTTACTGTCA AGAGTGAAAG GCCCAGACAG GAGCAAACCTC CGAGTGTATC 300
 TCCACTGCAC TAACGTCTAT CACCCACGAT ATCAGGAAGG AGATCTAACT CTGTATGTCC 360
 TGAACCTCCA TAATGTCACC AAGCACTTGA AGGTACCGCC TCCGTGTGTT AGGAAACCAG 420
 TGGATACGTA CCTTCTGAAG CCTTCGGGGC CGGATGGATT ACTTTCCAAA TCTGTCCAAC 480
 TGAACGGTCA AATTCTGAAG ATGGTGGATG AGCAGACCCT GCCAGCTTTG ACAGAAAAAC 540
 CTCTCCCCCG AGGAAGTGCA CTAAGCCTGC CTGCCTTTTC CTATGGTTTT TTTGTCTATA 600
 GAAATGCCAA AATCGTGCT TGTATATGAA AATAAAAGGC ATACGGTACC CTTGAGACAA 660
 AAGCCGAGGG GGGTGTTATT CATAAAACAA AACCCTAGTT TAGGAGGCCA CCTCCTTGCC 720
 GAGTTCCAGA GCTTCGGGAG GGTGGGGTAC ACTTCAGTAT TACATTCAGT GTGGTGTCT 780
 CTCTAAGAAG AATACTGCAG GTGGTGACAG TTAATAGCAC TGTG 824

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1899
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13

GGGAAAGCGA GCAAGGAAGT AGGAGAGAGC CGGGCAGGCG GGGCGGGGTT GGATTGGGAG 60
 CAGTGGGAGG GATGAGAAG AGGAGTGGGA GGGATGGAGG GCGCAGTGGG AGGGGTGAGG 120
 AGGCGTAACG GGGCGGAGGA AAGGAGAAAA GGGCGCTGGG GCTCGGCGGG AGGAAAGTGCT 180
 AGAGCTCTCG ACTCTCGCT GCGCGGCAGC TGGCGGGGGG AGCAGCCAGG TGAGCCCAAG 240
 ATGCTGCTGC GCTCGAAGCC TGGCTGCGG CCGCGCTGA TGCTGCTGCT CCTGGGGCCG 300
 CTGGTCCCC TCTCCCTGG CGCCCTGCCG CGACCTGCGC AAGCACAGGA CGTCGTGGAC 360
 CTGGACTTCT TCACCCAGGA GCCGCTGCAC CTGGTGAGCC CCTCGTTTCT GTCCGTCAAC 420
 ATTGACGCCA ACCTGGCCAC GGACCGCGG TTCCTCATCC TCCTGGGTTC TCCAAAGCTT 480
 CGTACCTTGG CCAGAGGCTT GTCTCCTGCG TACCTGAGGT TTGGTGGCAC CAAGACAGAC 540
 TTCCTAATTT TCGATCCCAA GAAGGAATCA ACCTTTGAAG AGAGAAAGTTA CTGGCAATCT 600
 CAAGTCAACC AGGATATTTG CAAATATGGA TCCATCCCTC CTGATGTGGA GGAGAAGTTA 660
 CGGTTGGAAT GGCCCTACCA GGAGCAATTG CTACTCCGAG AACACTACCA GAAAAAGTTC 720
 AAGAACAGCA CTAATCAAG AAGCTCTGTA GATGTGCTAT ACACCTTTTC AAACCTGCTCA 780

```

GGACTGGACT TGATCTTTGG CCTAAATGCG TTATTAAGAA CAGCAGATT GCAGTGGAAC 840
AGTTCTAATG CTCAGTTGCT CCTGGACTAC TGCTCTTCCA AGGGGTATAA CATTCTTGG 900
GAACTAGGCA ATGAACCTAA CAGTTTCCTT AAGAAGGCTG ATATTTTCAT CAATGGGTCG 960
CAGTTAGGAG AAGATTATAT TCAATTGCAT AACTTCTAA GAAAGTCCAC CTTCAAAAAT 1020
GCAAACTCT ATGGTCTCGA TGTTGGTCAG CCTCGAAGAA AGACGGCTAA GATGCTGAAG 1080
AGCTTCCTGA AGGCTGGTGG AGAAGTGATT GATTGAGTTA CATGGCATCA CTACTATTTG 1140
AATGGACCGA CTGCTACCAG GGAAGATTTT CTAAACCTG ATGTATTGGA CATTTTTATT 1200
TCATCTGTGC AAAAAAGTTT CCAGGTGGTT GAGAGCACCA GGCCTGGCAA GAAGGTCTGG 1260
TTAGAGAGAA CAAGCTCTGC ATATGGAGGC GGAGCGCCCT TGCTATCCGA CACCTTTGCA 1320
GCTGGCTTTA TGTGGCTGGA TAAATTGGGC CTGTCAGCCC GAATGGGAAT AGAAGTGGTG 1380
ATGAGGCAAG TATTCTTTGG AGCAGGAAAC TACCATTAG TGGATGAAA CTTGATCCT 1440
TTACCTGATT ATTGGCTATC TCTTCTGTT CAGAAATTGG TGGGCACCAA GGTGTTAATG 1500
GCAAGCGTGC AAGGTTCAA GAGAAGGAAG CTTGAGTAT ACCTTCATT CACAAACACT 1560
GACAATCCAA GGTATAAGA AGGAGATTTA ACTCTGTAT CCATAAACCT CCATAACGTC 1620
ACCAAGTACT TCGCGTTACC CTATCCTTTT TCTAACAAGC AAGTGGATAA ATACCTTCTA 1680
AGACCTTTGG GACCTCATGG ATTACTTTCC AAATCTGTCC AACTCAATGG TCTAACTCTA 1740
AAGATGGTGG ATGATCAAC CTGCCACCT TTAATGAAA AACCTCTCC GCCAGGAAGT 1800
TCACTGGGCT TGCCAGCTTT CTCATATAGT TTTTGTGA TAAGAAATGC CAAAGTTGCT 1860
GCTTGCATCT GAAAATAAAA TATACTAGTC CTGACACTG 1899

```

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 592
 (B) TYPE: amino acid
 (C) STRANDEDNESS: singl
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14

```

Met Glu Gly Ala Val Gly Gly Val Arg Arg Arg Asn Gly Ala Glu
      5              10              15
Glu Arg Arg Lys Gly Arg Trp Gly Ser Ala Gly Gly Ser Ala Arg
      20              25              30
Ala Leu Asp Ser Pro Leu Arg Gly Ser Trp Arg Gly Glu Gln Pro
      35              40              45
Gly Glu Pro Lys Met Leu Leu Arg Ser Lys Pro Ala Leu pro Pro
      50              55              60
Pro Leu Met Leu Leu Leu Gly Pro Leu Gly Pro Leu Ser Pro
      65              70              75
Gly Ala Leu Pro Arg Pro Ala Gln Ala Gln Asp Val Val Asp Leu
      80              85              90
Asp Phe Phe Thr Gln Glu Pro Leu His Leu Val Ser Pro Ser Phe
      95             100             105
Leu Ser Val Thr Ile Asp Ala Asn Leu Ala Thr Asp Pro Arg Phe
      110            115            120
Leu Ile Leu Leu Gly Ser Pro Lys Leu Arg Thr Leu Ala Arg Gly
      125            130            135
Leu Ser Pro Ala Tyr Leu Arg Phe Gly Gly Thr Lys Thr Asp Phe
      140            145            150
Leu Ile Phe Asp Pro Lys Lys Glu Ser Thr Phe Glu Glu Arg Ser
      155            160            165
Tyr Trp Gln Ser Gln Val Asn Gln Asp Ile Cys Lys Tyr Gly Ser
      170            175            180
Ile Pro Pro Asp Val Glu Glu Lys Leu Arg Leu Glu Trp Pro Tyr
      185            190            195
Gln Glu Gln Leu Leu Leu Arg Glu His Tyr Gln Lys Lys Phe Lys
      200            205            210
Asn Ser Thr Tyr Ser Arg Ser Ser Val Asp Val Leu Tyr Thr Phe

```

10

215	220	225
Ala Asn Cys Ser Gly Leu Asp Leu Ile	Phe Gly Leu Asn Ala Leu	
230	235	240
Leu Arg Thr Ala Asp Leu Gln Trp Asn Ser Ser Asn Ala Gln Leu		
245	250	255
Leu Leu Asp Tyr Cys Ser Ser Lys Gly Tyr Asn Ile Ser Trp Glu		
260	265	270
Leu Gly Asn Glu Pro Asn Ser Phe Leu Lys Lys Ala Asp Ile Phe		
275	280	285
Ile Asn Gly Ser Gln Leu Gly Glu Asp Tyr Ile Gln Leu His Lys		
290	295	300
Leu Leu Arg Lys Ser Thr Phe Lys Asn Ala Lys Leu Tyr Gly Pro		
305	310	315
Asp Val Gly Gln Pro Arg Arg Lys Thr Ala Lys Met Leu Lys Ser		
320	325	330
Phe Leu Lys Ala Gly Gly Glu Val Ile Asp Ser Val Thr Trp His		
335	340	345
His Tyr Tyr Leu Asn Gly Arg Thr Ala Thr Arg Glu Asp Phe Leu		
350	355	360
Asn Pro Asp Val Leu Asp Ile Phe Ile Ser Ser Val Gln Lys Val		
365	370	375
Phe Gln Val Val Glu Ser Thr Arg Pro Gly Lys Lys Val Trp Leu		
380	385	390
Gly Glu Thr Ser Ser Ala Tyr Gly Gly Gly Ala Pro Leu Leu Ser		
395	400	405
Asp Thr Phe Ala Ala Gly Phe Met Trp Leu Asp Lys Leu Gly Leu		
410	415	420
Ser Ala Arg Met Gly Ile Glu Val Val Met Arg Gln Val Phe Phe		
425	430	435
Gly Ala Gly Asn Tyr His Leu Val Asp Glu Asn Phe Asp Pro Leu		
440	445	450
Pro Asp Tyr Trp Leu Ser Leu Leu Phe Lys Lys Leu Val Gly Thr		
455	460	465
Lys Val Leu Met Ala Ser Val Gln Gly Ser Lys Arg Arg Lys Leu		
470	475	480
Arg Val Tyr Leu His Cys Thr Asn Thr Asp Asn Pro Arg Tyr Lys		
485	490	495
Glu Gly Asp Leu Thr Leu Tyr Ala Ile Asn Leu His Asn Val Thr		
500	505	510
Lys Tyr Leu Arg Leu Pro Tyr Pro Phe Ser Asn Lys Gln Val Asp		
515	520	525
Lys Tyr Leu Leu Arg Pro Leu Gly Pro His Gly Leu Leu Ser Lys		
530	535	540
Ser Val Gln Leu Asn Gly Leu Thr Leu Lys Met Val Asp Asp Gln		
545	550	555
Thr Leu Pro Pro Leu Met Glu Lys Pro Leu Arg Pro Gly Ser Ser		
560	565	570
Leu Gly Leu Pro Ala Phe Ser Tyr Ser Phe Phe Val Ile Arg Asn		
575	580	585
Ala Lys Val Ala Ala Cys Ile		
590	592	

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1899
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double

11

(D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:15

```

                                GGG      3
AAA GCG AGC AAG GAA GTA GGA GAG AGC CGG GCA GGC GGG GCG GGG  48
TTG GAT TGG GAG CAG TGG GAG GGA TGC AGA AGA GGA GTG GGA GGG  93
ATG GAG GGC GCA GTG GGA GGG GTG AGG AGG CGT AAC GGG GCG GAG  138
Met Glu Gly Ala Val Gly Gly Val Arg Arg Arg Asn Gly Ala Glu
      5              10              15

GAA AGG AGA AAA GGG CGC TGG GGC TCG GCG GGA GGA AGT GCT AGA  183
Glu Arg Arg Lys Gly Arg Trp Gly Ser Ala Gly Gly Ser Ala Arg
      20              25              30

GCT CTC GAC TCT CCG CTG CGC GGC AGC TGG CCG GGG GAG CAG CCA  228
Ala Leu Asp Ser Pro Leu Arg Gly Ser Trp Arg Gly Glu Gln Pro
      35              40              45

GGT GAG CCC AAG ATG CTG CTG CGC TCG AAG CCT GCG CTG CCG CCG  273
Gly Glu Pro Lys Met Leu Leu Arg Ser Lys Pro Ala Leu Pro Pro
      50              55              60

CCG CTG ATG CTG CTG CTC CTG GGG CCG CTG GGT CCC CTC TCC CCT  318
Pro Leu Met Leu Leu Leu Leu Gly Pro Leu Gly Pro Leu Ser Pro
      65              70              75

GGC GCC CTG CCC CGA CCT GCG CAA GCA CAG GAC GTC GTG GAC CTG  363
Gly Ala Leu Pro Arg Pro Ala Gln Ala Gln Asp Val Val Asp Leu
      80              85              90

GAC TTC TTC ACC CAG GAG CCG CTG CAC CTG GTG AGC CCC TCG TTC  408
Asp Phe Phe Thr Gln Glu Pro Leu His Leu Val Ser Pro Ser Phe
      95              100             105

CTG TCC GTC ACC ATT GAC GCC AAC CTG GCC ACG GAC CCG CGG TTC  453
Leu Ser Val Thr Ile Asp Ala Asn Leu Ala Thr Asp Pro Arg Phe
     110              115             120

CTC ATC CTC CTG GGT TCT CCA AAG CTT CGT ACC TTG GCC AGA GGC  498
Leu Ile Leu Leu Gly Ser Pro Lys Leu Arg Thr Leu Ala Arg Gly
     125              130             135

TTG TCT CCT GCG TAC CTG AGG TTT GGT GGC ACC AAG ACA GAC TTC  543
Leu Ser Pro Ala Tyr Leu Arg Phe Gly Gly Thr Lys Thr Asp Phe
     140              145             150

CTA ATT TTC GAT CCC AAG AAG GAA TCA ACC TTT GAA GAG AGA AGT  588
Leu Ile Phe Asp Pro Lys Lys Glu Ser Thr Phe Glu Glu Arg Ser
     155              160             165

TAC TGG CAA TCT CAA GTC AAC CAG GAT ATT TGC AAA TAT GGA TCC  633
Tyr Trp Gln Ser Gln Val Asn Gln Asp Ile Cys Lys Tyr Gly Ser
     170              175             180

ATC CCT CCT GAT GTG GAG GAG AAG TTA CGG TTG GAA TGG CCC TAC  678
Ile Pro Pro Asp Val Glu Glu Lys Leu Arg Leu Glu Trp Pro Tyr
     185              190             195

```

CAG GAG CAA TTG CTA CTC CGA GAA CAC TAC CAG AAA AAG TTC AAG	723
Gln Glu Gln Leu Leu Leu Arg Glu His Tyr Gln Lys Lys Phe Lys	
200 205 210	
AAC AGC ACC TAC TCA AGA AGC TCT GTA GAT GTG CTA TAC ACT TTT	768
Asn Ser Thr Tyr Ser Arg Ser Ser Val Asp Val Leu Tyr Thr Phe	
215 220 225	
GCA AAC TGC TCA GGA CTG GAC TTG ATC TTT GGC CTA AAT GCG TTA	813
Ala Asn Cys Ser Gly Leu Asp Leu Ile Phe Gly Leu Asn Ala Leu	
230 235 240	
TTA AGA ACA GCA GAT TTG CAG TGG AAC AGT TCT AAT GCT CAG TTG	858
Leu Arg Thr Ala Asp Leu Gln Trp Asn Ser Ser Asn Ala Gln Leu	
245 250 255	
CTC CTG GAC TAC TGC TCT TCC AAG GGG TAT AAC ATT TCT TGG GAA	903
Leu Leu Asp Tyr Cys Ser Ser Lys Gly Tyr Asn Ile Ser Trp Glu	
260 265 270	
CTA GGC AAT GAA CCT AAC AGT TTC CTT AAG AAG GCT GAT ATT TTC	948
Leu Gly Asn Glu Pro Asn Ser Phe Leu Lys Lys Ala Asp Ile Phe	
275 280 285	
ATC AAT GGG TCG CAG TTA GGA GAA GAT TAT ATT CAA TTG CAT AAA	993
Ile Asn Gly Ser Gln Leu Gly Glu Asp Tyr Ile Gln Leu His Lys	
290 295 300	
CTT CTA AGA AAG TCC ACC TTC AAA AAT GCA AAA CTC TAT GGT CCT	1038
Leu Leu Arg Lys Ser Thr Phe Lys Asn Ala Lys Leu Tyr Gly Pro	
305 310 315	
GAT GTT GGT CAG CCT CGA AGA AAG ACG GCT AAG ATG CTG AAG AGC	1083
Asp Val Gly Gln Pro Arg Arg Lys Thr Ala Lys Met Leu Lys Ser	
320 325 330	
TTC CTG AAG GCT GGT GGA GAA GTG ATT GAT TCA GTT ACA TGG CAT	1128
Phe Leu Lys Ala Gly Gly Glu Val Ile Asp Ser Val Thr Trp His	
335 340 345	
CAC TAC TAT TTG AAT GGA CGG ACT GCT ACC AGG GAA GAT TTT CTA	1173
His Tyr Tyr Leu Asn Gly Arg Thr Ala Thr Arg Glu Asp Phe Leu	
350 355 360	
AAC CCT GAT GTA TTG GAC ATT TTT ATT TCA TCT GTG CAA AAA GTT	1218
Asn Pro Asp Val Leu Asp Ile Phe Ile Ser Ser Val Gln Lys Val	
365 370 375	
TTC CAG GTG GTT GAG AGC ACC AGG CCT GGC AAG AAG GTC TGG TTA	1263
Phe Gln Val Val Glu Ser Thr Arg Pro Gly Lys Lys Val Trp Leu	
380 385 390	
GGA GAA ACA AGC TCT GCA TAT GGA GGC GGA GCG CCC TTG CTA TCC	1308
Gly Glu Thr Ser Ser Ala Tyr Gly Gly Gly Ala Pro Leu Leu Ser	
395 400 405	

13

GAC ACC TTT GCA GCT GGC TTT ATG TGG CTG GAT AAA TTG GGC CTG	1353
Asp Thr Phe Ala Ala Gly Phe Met Trp Leu Asp Lys Leu Gly Leu	
410 415 420	
TCA GCC CGA ATG GGA ATA gAA GTG GTG ATG AGG CAA GTA TTC TTT	1398
Ser Ala Arg Met Gly Ile Glu Val Val Met Arg Gln Val Phe Phe	
425 430 435	
GGA GCA GGA AAC TAC CAT TTA GTG GAT GAA AAC TTC GAT CCT TTA	1443
Gly Ala Gly Asn Tyr His Leu Val Asp Glu Asn Phe Asp Pro Leu	
440 445 450	
CCT GAT TAT TGG CTA TCT CTT CTG TTC AAG AAA TTG GTG GGC ACC	1488
Pro Asp Tyr Trp Leu Ser Leu Leu Phe Lys Lys Leu Val Gly Thr	
455 460 465	
AAG GTG TTA ATG GCA AGC GTG CAA GGT TCA AAG AGA AGG AAG CTT	1533
Lys Val Leu Met Ala Ser Val Gln Gly Ser Lys Arg Arg Lys Leu	
470 475 480	
CGA GTA TAC CTT CAT TGC ACA AAC ACT GAC AAT CCA AGG TAT AAA	1578
Arg Val Tyr Leu His Cys Thr Asn Thr Asp Asn Pro Arg Tyr Lys	
485 490 495	
GAA GGA GAT TTA ACT CTG TAT GCC ATA AAC CTC CAT AAC GTC ACC	1623
Glu Gly Asp Leu Thr Leu Tyr Ala Ile Asn Leu His Asn Val Thr	
500 505 510	
AAG TAC TTG CGG TTA CCC TAT CCT TTT TCT AAC AAG CAA GTG GAT	1668
Lys Tyr Leu Arg Leu Pro Tyr Pro Phe Ser Asn Lys Gln Val Asp	
515 520 525	
AAA TAC CTT CTA AGA CCT TTG GGA CCT CAT GGA TTA CTT TCC AAA	1713
Lys Tyr Leu Leu Arg Pro Leu Gly Pro His Gly Leu Leu Ser Lys	
530 535 540	
TCT GTC CAA CTC AAT GGT CTA ACT CTA AAG ATG GTG GAT GAT CAA	1758
Ser Val Gln Leu Asn Gly Leu Thr Leu Lys Met Val Asp Asp Gln	
545 550 555	
ACC TTG CCA CCT TTA ATG GAA AAA CCT CTC CGG CCA GGA AGT TCA	1803
Thr Leu Pro Pro Leu Met Glu Lys Pro Leu Arg Pro Gly Ser Ser	
560 565 570	
CTG GGC TTG CCA GCT TTC TCA TAT AGT TTT TTT GTG ATA AGA AAT	1848
Leu Gly Leu Pro Ala Phe Ser Tyr Ser Phe Phe Val Ile Arg Asn	
575 580 585	
GCC AAA GTT GCT GCT TGC ATC TGA AAA TAA AAT ATA CTA GTC CTG	1893
Ala Lys Val Ala Ala Cys Ile	
590 592	
ACA CTG	1899

(2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 594
- (B) TYPE: nucleic acid

14

- (C) STRANDEDNESS: double
(D) TOPOLOGY: linear
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16

ATTACTATAG GGCACGCGTG GTCGACGGCC CGGGCTGGTA TTGTCTTAAT GAGAAGTTGA 60
TAAAGAATTT TGGGTGGTTG ATCTCTTTCC AGCTGCAGTT TAGCGTATGC TGAGGCCAGA 120
TTTTTTCAGG CAAAAGTAAA ATACCTGAGA AACTGCCTGG CCAGAGGACA ATCAGATTTT 180
GGCTGGCTCA AGTGACAAGC AAGTGTATAT AAGCTAGATG GGAGAGGAAG GGATGAATAC 240
TCCATGGAG GCTTTACTCG AGGGTCAGAG GGATACCCGG CGCCATCAGA ATGGGATCTG 300
GGAGTCGGAA ACGCTGGGTT CCCACGAGAG CGCGCAGAAC ACGTGCCTCA GGAAGCCTGG 360
TCCGGGATGC CCAGCGCTGC TCCCCGGGCG CTCCTCCCCG GCGCTCCTC CCCAGGCCTC 420
CCGGGGCTT GGATCCCGGC CATCTCCGCA CCCTTCAAGT GGGTGTGGGT GATTTCGTAA 480
GTGAACGTGA CCGCCACCGG GGGGAAAGCG AGCAAGGAAG TAGGAGAGAG CCGGGCAGGC 540
GGGGCGGGT TGGATTGGGA GCAGTGGGAG GGATGCAGAA GAGGAGTGGG AGGG 594

(2) INFORMATION FOR SEQ ID NO:17:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17
CCCCAGGAGC AGCAGCATCA G 21

(2) INFORMATION FOR SEQ ID NO:18:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18
AGGCTTCGAG CGCAGCAGCA T 21

(2) INFORMATION FOR SEQ ID NO:19:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 22
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19
GTAATACGAC TCACTATAGG GC 22

(2) INFORMATION FOR SEQ ID NO:20:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 19
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20
ACTATAGGGC ACGCGTGGT 19

(2) INFORMATION FOR SEQ ID NO:21:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21
CTTGGGCTCA CCTGGCTGCT C 21

(2) INFORMATION FOR SEQ ID NO:22:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 23
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22
AGCTCTGTAG ATGTGCTATA CAC 23

(2) INFORMATION FOR SEQ ID NO:23:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 22
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23
GCATCTTAGC CGTCTTTCTT CG 22

(2) INFORMATION FOR SEQ ID NO:24:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 23
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24
GAGCAGCCAG GTGAGCCAA GAT 23

(2) INFORMATION FOR SEQ ID NO:25:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 23
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25
TTCGATCCA AGAAGGAATC AAC 23

(2) INFORMATION FOR SEQ ID NO:26:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 23
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26
AGCTCTGTAG ATGTGCTATA CAC 23

(2) INFORMATION FOR SEQ ID NO:27:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27
TCAGATGCAA GCAGCAACTT TGGC 24

16

(2) INFORMATION FOR SEQ ID NO:28:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 22
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28

GCATCTTAGC CGTCTTCTT CG 22

(2) INFORMATION FOR SEQ ID NO:29:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29

GTAGTGATGC CATGTAAC TG AATC 24

(2) INFORMATION FOR SEQ ID NO:30:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 22
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30

AGGCACCCTA GAGATGTTCC AG 22

(2) INFORMATION FOR SEQ ID NO:31:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31

GAAGATTCT GTTCCATGA CGTG 24

(2) INFORMATION FOR SEQ ID NO:32:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 25
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32

CCACACTGAA TGTAATACTG AAGTG 25

(2) INFORMATION FOR SEQ ID NO:33:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 22
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33

CGAAGCTCTG GAACTCGGCA AG 22

(2) INFORMATION FOR SEQ ID NO:34:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 22

17

- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34
GCCAGCTGCA AAGGTGTTGG AC 22

(2) INFORMATION FOR SEQ ID NO:35:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 23
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35
AACACCTGCC TCATCAGGAC TTC 23

(2) INFORMATION FOR SEQ ID NO:36:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 22
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36
GCCAGGCTGG CGTCGATGGT GA 22

(2) INFORMATION FOR SEQ ID NO:37:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 22
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37
GTCGATGGTG ATGGACAGGA AC 22

(2) INFORMATION FOR SEQ ID NO:38:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 22
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38
GTAATACGAC TCACTATAGG GC 22

(2) INFORMATION FOR SEQ ID NO:39:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:39
ACTATAGGGC ACGCGTGGT 19

(2) INFORMATION FOR SEQ ID NO:40:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 27
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

18

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:40
CCATCCTAAT ACGACTCACT ATAGGGC 27

(2) INFORMATION FOR SEQ ID NO:41:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 23
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:41
ACTCACTATA GGGCTCGAGC GGC 23

(2) INFORMATION FOR SEQ ID NO:42:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 44848
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:42

GGATCTTGGC	TCACTGCAAT	CTCTGCCTCC	CATGCAATTC	TTATGCATCA	50
GCCTCCTGAG	TAGCTTGGAT	TATAGGCTCG	CGCCACCACT	CCTGGCTACA	100
CCATGTTGCC	CAGGCTGGTC	TTGAACTCTT	GGGCTCTAGT	GATCCACCCG	150
CCTTGGCCTC	CCAAAGTGCT	GGGATTACAG	GTGTGAGCCA	TCACACCCGG	200
CCCCCGCTTT	CCATATTAGT	AACACACATG	TAGACCACAA	GGATGCACTA	250
TTTAGAAAAC	TTGCAATGGT	CCACTTTTCA	AATCACCCAA	ACATGTTAAA	300
GAAATTGGTA	TGACTGGGCA	TGGCACAGTG	GCTCATGCCT	GCAATCCTAG	350
CATTTTGTGA	GGCTGAGACG	GGCAGATCAC	GAGGTCAGGA	GATTGAGACC	400
ATCCTGACAG	ACATGGTGAA	ATCCCATCTC	TACTAAAAAT	ACAAAACRAAT	450
TAGCCCGGGG	TGATGGCAGG	CCCCTGTAGT	CCCAGCTACT	CGGGAGGCTG	500
AGGCAGGAGA	ATGGCGTGAA	TCCAGGAGGC	AGAGCTTGCA	GTGAGCCGAG	550
ATGGTGCCAC	TGCACTCCAG	CCTGGGCGAC	AGAGCGAGAC	TCCGTCTCAA	600
AAAAAATAAA	AAAGAAAGAA	ATTGGTATGA	CTGTTGACTC	ACAACAGGAG	650
TCAGGGGCAT	GGGGTGGGGT	GTAAGATTAA	TGTCATGACA	AATGTGGAAA	700
AGAAACTTCT	GTTTTTCCAA	CTCCACGTCT	GCTACCATAT	TATTACACTC	750
TTCTGTAGT	GTGGTGTATA	TGTTGTAATT	TTTTTTCATA	TGTATACAGT	800
AATTGTAGGA	TATGAACCTG	ATTCTAGTTG	CAAACTCAC	TATGAGCTTA	850
GCTTTTAAGT	TGCTTAAGAA	TAGGTAGATC	TATGCAATA	ATGATAATTA	900
TTATTATTAT	TTTAAGAGAG	GGTCTCACTT	TGTCACCCAG	GCTGGAGTGC	950
AGTGGTGTGA	TTAAGGGTCA	CTGCAACCTC	CACCTCCAG	GCTCAAAATA	1000
ACCTCCACAC	TCAGCCTCCC	CAGTAGCTGG	AACCACAGGC	ACGGGCCACC	1050
ACGCCCTGGT	AATTTTTTGT	ATTTTTTTGTA	GAGATGGGGT	TTTCATCATGT	1100
TGCCCAGGCT	GTCTTTGAAT	TCCTCGGCTC	AAGCAATCCT	CCCACCTTGG	1150
CCTCCCAAAA	TGCTGGCATC	ACAGGCATGA	TGGCATCACT	GGCATCACAT	1200
ACCATGCCTG	GCCTGATTTA	TGCAAAATTAG	ATATGCATTT	CAAAATAATC	1250
TATTTTATAT	TGTTGCCTTA	TTGGTGGTAC	AATCTCAAGT	GGAAAAATCT	1300
AAGGGTTTTG	GTGTTATTTG	CTTACTCAAC	CAATATTTAT	TAGACTCTTA	1350
CTAAGCACCA	ACATGATCAC	ATGCCTGAGC	TATGGCTAGC	ATAGCGTGTG	1400
AGACAAACTT	AATCTCTGTT	TTGGTGGAGC	ATATAATCTA	GTAGATGAAG	1450
CCAATGTTGA	GCAACATCAC	AATACTAACA	AATTGAGGAT	GCTACGAGAG	1500
TGTCTAACAA	ATTGAGGATG	CTACGAGAGT	GTCTAACAAA	TTGAGGATGC	1550
TATGAGAGTG	TGTCATGGAG	AGCTGCCTGG	AGATTGAGAG	AAAGCTTCCT	1600
TGAGGGAAGT	TACATTTTCA	CTGAAACACA	CTGCCATCTG	CTCGAGGTTT	1650
TGTAACCTGCA	TTACATATCC	GATTCTGACA	CTTCACATCC	CGATTCTGAC	1700
ACTTCACCCA	GTTACTGTCT	CAGAGCTTGG	GTCCGCATGT	GTAAAAAAG	1750
GACAGTATGC	ACTTGGCAGG	GTGTGAGAA	GGGAAGAGAA	CACAAGTAAA	1800
GCACCTGTAT	CAGGCATACA	GTAGGCATA	AGCGTGCGAT	GCTTGCTATG	1850
ATTATACATC	AGTGTAAGCA	TCAAGGAAAA	GCTGAAGAAA	AGTCTGACCA	1900
ACAGCGAAAG	ATAAATGCGC	AGAGGAGAAA	TTTGGCAAA	GCTCCTCAAT	1950
CAGGGGCAGT	CCGTACTCTA	CACCTTTGTAT	GGGGGCTTCA	GGTCTGAGT	2000
TCCAGACATT	GGAGCAACTA	ACCCTTTAAG	ATTGCTAAAT	ATTGCTCTAA	2050
TGAGAAGTTG	ATAAAGAATT	TTGGGTGGTT	GATCTCTTTC	CAGCTGCAGT	2100
TTAGCGTATG	CTGAGGCCAG	ATTTTTTCAA	GCAAAAGTAA	AATACCTGAG	2150
AAACTGCCTG	GCCAGAGGAC	AATCAGATTT	TGGCTGGCTC	AAGTGACAAG	2200
CAAGTGTTTA	TAAGCTAGAT	GGGAGAGGAA	GGGATGAATA	CTCCATTGGA	2250
GGTTTACTC	GAGGGTCAGA	GGGATACCCG	GCGCCATCAG	AATGGGATCT	2300
GGGAGTCGGA	AACGCTGGGT	TCCCACGAGA	GCGCGCAGAA	CACGTGCGTC	2350
AGGAAGCCTG	GTCCGGGATG	CCCAGCGCTG	CTCCCGGGC	GCTCCTCCCC	2400

GGGCGCTCCT	CCCCAGGCCT	CCCGGGCGCT	TGGATCCCGG	CCATCTCCGC	2450
ACCCCTCAAG	TGGGTGTGGG	TGATTTCGTA	AGTGAACGTG	ACCGCCACCG	2500
AGGGGAAAGC	GAGCAAGGAA	GATAGGAGAG	GCCGGGCGAG	CGGGGCGGGG	2550
TTGGATTGGG	AGCAGTGGGA	GGGATGCAGA	AGAGGAGTGG	GAGGGATGGA	2600
GGGCGCAGTG	GGAGGGGTGA	GGAGGCGTAA	CGGGGCGGAG	GAAAGGAGAA	2650
AAGGGCGCTG	GGGCTCGGCG	GGAGGAAGTG	CTAGAGCTCT	CGACTCTCCG	2700
CTGCGCGGCA	GCTGGCGGGG	GGAGCAGCCA	GGTGAGCCCA	AGATGCTGCT	2750
GCGCTCGAAG	CCTGCGCTGC	CGCGCGCGCT	GATGCTGCTG	CTCCTGGGGC	2800
CGCTGGGTCC	CCTCTCCCCT	GGCGCCCTGC	CCCGACCTGC	GCAAGCACAG	2850
GACGTCGTGG	ACCTGGACTT	CTTCACCCAG	GAGCCGCTGC	ACCTGGTGAG	2900
CCCCTCGTTC	CTGTCCGTCA	CCATTGACGC	CAACCTGGCC	ACGGACCCGC	2950
GGTTCTCAT	CCTCTGGGG	TAAGCGCCAG	CCTCTGGTC	CTGTCCCCTT	3000
TCCTGTCTTC	CTGACACCTA	TGCTGCCCC	GCCAGCGGCT	CTCCTTCTTT	3050
TGCGCGGAAA	CAACTTCACA	CCGGAACTTC	CCCGCTGTCT	TCTCCCCACC	3100
CCACTTCCCG	CCTCTCATT	TCCCTCTCCC	TCCCTTACTC	TCAGACCCCA	3150
AACCGCTTTT	TGGGGGGTAT	CATTAAAAA	ATAGATTTAG	GGGTTACAAG	3200
TGCAGTTCCT	TTCATGGGT	ATATTGCATT	GTGGTGGCAT	CTGGGCTCTT	3250
AGTGTAACTG	TACCCGGAAT	GTTGTACATT	GTATCTAATA	GGTAATTCTT	3300
CATCCTCAT	CCCTCTCCCA	CCCTCCACC	TTTTGGAGTC	TCCAGTGTCT	3350
ACTATTCCAC	TAAGTCCATG	TGTACACATT	GTTTAGCGCC	CACTCTAAAT	3400
GAGCCTTTTT	GTTTCATTCA	TTCTGTAAGT	GTTGAATAGG	CACCACTTAA	3450
GGTCAGGTAT	AAGTGGAAAT	TTGAAAAAGA	AAGTGGCCAC	TTGCCCCAGT	3500
ACTTCCCTAG	CCAAGAGGAG	GGAAACCAGG	CAGGTGCACC	TGAAGGCCTG	3550
TGAGTGCTTG	ATTTGCTGTG	CAGTGTAGGA	CAAGTAAGAT	TGTGCATAGC	3600
CTTCGTGATT	TAAGACTGTG	TTAGGAAGAT	TTCTCTTCT	TTTCTTTTCT	3650
TTTTCTTTTT	TCCTTTCTTT	TTTTTTTTTA	GGCAGATGAA	AAGGGCGTCA	3700
CAGAACAGGA	ATAAAATCT	AAATATTCAA	TAAATGAGAC	CTAGGAGACT	3750
ACTGCAGTGA	CTTACAAAGT	CCTAATAAAA	AGATGTCTCT	CCAAAATGGG	3800
GTCGCAAAAT	GTGGTGCTGC	CTTATCAGCT	CTAAGTTTTT	TCCTTACCTG	3850
AGAAAGAAAG	AACCTGATGC	AGGTTGAGGG	CTCCTGCCCC	ATGAATGCAG	3900
GCTGACTCCA	AGATGGGGAG	CTACAGGGAC	AATCCCAGGT	CTTCTAGGCC	3950
TCTTATTTAG	GCCTTGGGAG	CCTCCAGAGA	TGGCCACATC	TGACCAGGCC	4000
CAGATAGAGG	GAAAGATCAC	CATTATCTCA	CCTCTGTGTC	AAATACCTAG	4050
ATGCTGTCTC	CCCTGAGCCC	ACACTATAGT	TGCCAGCGCT	AATTTAATGG	4100
GTAGTGTACT	GGTTAAGAGA	TGGACAGACC	ATCCTGGCTT	GACTCTCAGC	4150
TCTGGCAAAAG	ATGAGTGACT	TGGTTTTTCC	ATATCTCTTG	GCCACACCAA	4200
CCTTGATTTC	TTCCAGCTGA	GAATGGAAAT	TCTCAAGCTT	GCCTCAAGGA	4250
TTATTGCCCG	AGGATTTGAT	GATATGGTAA	GAGCTTCTCA	GTGTTTGACC	4300
CATAGTAAGT	GTTTGACGTT	TCAAACGAAT	TGTTTCTTTC	TAGGACATGG	4350
TGAGCATTTG	GTAGCCATTTC	ACCGGTTTTTC	TGTTTCTTTG	GATCATAGTT	4400
AACCTCTCCT	TTTCTTCTG	GCACCTACAAT	TTTCTGTGTG	GGAAGAATCC	4450
TTACTTCTG	CCCTTCCCCT	TAAGGATAGG	AAGCTGATAC	TAGGCAGCAA	4500
CTAGTTGGGG	GATAGGAAGA	TTGTTCCAGA	GAAATGCTGA	ACCATAGGGC	4550
TCCAGATCAC	AGGACCCGAG	TCTTAGCTTG	CTGGGGGTGT	GGGTGGGGGG	4600
GGCGGGTTAC	TGAACATGGG	TATGAAGTAG	ATGTCCATTT	ACTGAAATGT	4650
GAGGACCTGA	GGCCTCTTCT	ATTGCTGTAG	CCAGCATATT	CCCCAACCTC	4700
TCCCCAAGAA	AGGACAGATG	GGGGTTCCCC	CCTGAGTAA	CAGGTCCAAA	4750
AGAAAAAACA	TACAGTGGGA	CTTCCAGGAT	CTGGGCCCTGA	TCACCCAGCA	4800
GTCAAGCTCC	CCGCAATTGA	CTAACACCCC	CCTAACACGT	AGAAATTCCA	4850
ATCTGCAATT	TAGTGAGGAT	GATACCTTTA	TTCTTCTTAA	ATACATCTCT	4900
TCATTTCCCA	GAGCACCCCT	TTTTCCCCTC	CTCTGCACCT	TTTTGTAAAA	4950
GACTGGAGTA	TAATGAAATA	CCAAGAGAGC	ATAACATGTG	ATACATAAAA	5000
CTTTTCTTCT	GGTTTACAAA	ACAGTTCATT	CTGTGCCATA	CGTGCTTCTC	5050
TCCAAGGCTG	GCTGCTGTCT	GTTCCAGCCC	GCTTCGCTTG	GAGAGGCCAT	5100
CTGCCATACC	TGCTCCCCAG	ACGCATCGAC	AAGCACACCC	AGAGTGTAT	5150
CTGCTAAGAC	CTAAAAGAGG	GAGGAACCCC	CTCTCCTCAT	CTAAGACCTA	5200
GCTTCTAAAT	TAGAGTGTGA	GGGTCCATCT	CCCCAGGAGG	GGCACAGGGC	5250
CCAAACAGCC	CAGCCATCTC	AGAAGACAAC	ACTAAGCTTT	GTAGGGGTCC	5300
ACAGTAGAGG	AGAGTAAGAC	GCCTGTTGTT	TAAATTATTA	CAGTTCTCTA	5350
AAAGTGAAGA	TGTGTGGGCG	GGATGGCAAG	AGCTGAGCAG	ACGAAAGCTG	5400
AAGGAATAAG	GAAAGAGAGG	AGGACACAAA	CAGCTGACAC	TTCTCTAGTT	5450
CTTGTCATTT	GCCTGGCCCT	GTTCTAAGCA	CCTTCTAGGT	ATTAATCCAT	5500
TTAGTCTTGG	CTACAACACT	GTGAGTAACT	AGTTTGTCTA	CCCCCATTTT	5550
AAAAATGAAG	AAAGTGAGGC	TCAGGGAGGT	TAAGTAACTT	GGCCACAGTT	5600
TGAAACTAGA	CTTGATCAC	ATGAGATAAT	AGTGCCCAATA	AAAAGGGAAA	5650
GCAGATTATA	TTTTTTAAAG	GAAAGAGAGT	AGGATATGGT	AGAAAAAGAT	5700
TGTTTGGAAA	GGAATTGAGA	GATTGATATA	ATGAAAAGAA	GCATTACAT	5750
GAGAGTAACA	GTATCAGGGC	CCAAACCTTC	ATCTAAGGTA	CTTCAAGAG	5800
GCCTAAGCAA	ACTTAGTCA	TGGCGTGGTT	CTAGTCTCCA	TGATGGCAAA	5850
TACATTGTGT	ACAGCCCAAC	TCCACACAAA	ACTTAAATAC	CAATGATAGA	5900
GCAATCTAAA	ATTTGAAAGA	AAAAATCTTT	CAATTTGTCT	TCTTCCCAGA	5950
GGGACTTAAT	CAAGAAACCA	ATCAAAATAC	TTCTTAAGCC	TAAGTGTGTG	6000
CAGAACTCCA	AAGAGAGCCC	AGCCCTAAAT	CAACACTGTC	CAATGGAAAT	6050
ATAATATAAT	GTGGCCCTCA	TATGCAAGGT	CATATGTAAT	TTTAAATTTT	6100
CTAGTAGCCA	TATTAAAAAG	GTA AAAAGAA	ACAAGTGAAA	TAAATTTTAA	6150

TAATTTTATT	TAGTTCAATA	GATCCAAAAT	GTTTCTCAG	CATGTAATCA	6200
ATATAAAAAT	ATTAATGAGG	TATTTATTAT	TCCTTTTCTC	AAACCAAGTC	6250
TATTTCTATAA	TCTGGCGTGT	ATTATTTACA	GCACTTCTCA	GACTATATTT	6300
CTTTCTTTCT	TTTTTTTTTC	CGAGACAAAT	TTGCTCTTGT	CACCCAAGCT	6350
AGAGTACAAT	GGCGTTACCT	CGGCTCACTG	CAACCTCCGC	CTCCCGGGTT	6400
CAAGTTATTCT	TCCTGSCCTCA	GTCTCCCAAG	TAGCTGGGAC	TAGAGGCATG	6450
CACCACCACG	CCTGGCTAAT	TGTGTATTTT	TAGTAGAGAC	AGGGTTTCAC	6500
CATGTTGGCC	AGGCTAACTC	CAAACTCCTG	AGCTCAGGTG	ATATGCCAC	6550
CTCGGCTCC	CAAAGTGTG	GGATTACAGG	CGTGAGCCAC	TGCACCCGGC	6600
CTCAGATTAA	CTATATTTCA	AGCGTTCAGT	AGCCACATGT	AGCTAGTGCT	6650
ATGGTAGTGG	ACAGTACAGA	TCTGCATTTC	AATTAAGACA	CGTATACAAG	6700
CATAGTTTAC	TAATGCACGG	TAAAAAAAAG	TATAGTGTG	AGTCGGTGGT	6750
AGAAATCCTA	AATACTGCAG	AGCAAAAGTG	GTACGAACAG	CAATCTCAGT	6800
GATAATGCAA	CCATGCTTGC	TTTTTCATTG	AATTTGCTTA	TTTTCTCTCA	6850
GCAAAAGTTCA	TCCATTTTTG	CCAATTCAAT	AAATATTTAC	TGATAAAAAAC	6900
TTTCAATATT	AGATTCTTGC	ATCTTCATAG	ACAGAGTTGC	TTTTACATT	6950
TAGAAAAATTA	CTTATCAATG	TTAAACACAC	GTTTTGATAA	CCAGTGTGGG	7000
AAAGAGGTGC	AGACTCCCCA	TGTGCTTATT	GATGGCAGAA	ATATTCACAG	7050
CCAAAGGGAA	ACAAAGGGCT	GGGGACAATC	ACACACCTCA	TGTCTCTTAA	7100
CTCCTGGGAA	GTGCTGTCCC	TCTGATTGAG	CTCTTATTAT	TGCCTTCCCC	7150
ACTAACCTTG	TCCACTGTGC	CCTGGAGCCC	TTTGCAGGGT	TACCTGCTCT	7200
GCTCCTCTCA	CAGAATATCT	CCTCTACCTC	CTTGTCCAAG	CTACAACCTG	7250
GCTATTCTCT	GATGACACTG	TCTTCCCTGT	AGCCCTTTTG	AGTAATGGGT	7300
GCATATTCTC	CCATAGTCCA	GTTCTTTTCC	TGTTCTCCAG	TCTGGCTTCT	7350
GGATGACAGC	CCACTAGTTT	GAACCTCCATA	CTGCTATAGT	TCAAGTCCCT	7400
TTTGACTTGT	TACCTTGGGC	AAATTACCTC	CTTTTGTTCA	GGTTCCTTGT	7450
TTGTAAATG	ACGATAATAA	TGCCATTTCG	TTCAGTGGGT	TATTTTGAAA	7500
TTGAGTGAAA	GAAGGCGGGT	AGCTTCCCTA	CACGCTCAGT	GTAGACTAGC	7550
CTGATGTGCA	TTACGGGTGA	TGCCATGACT	CAGTGTGTTT	TCCTCATCTC	7600
CACATCTGGC	TCTCATCCAG	TGCTCCTGCT	TACGGCACTC	TGTCCCCCTC	7650
TTACTTACTC	CCCCTTATTA	ACTGAAGACT	GGCACTGATC	TCACAGTTTC	7700
CTCTCCACTT	CTAGTCTCA	CCATCATCCT	AGATGACTTC	AAGTCACCTA	7750
GATAAACTGT	CTCAGTTTCT	TCACTCAGAT	TTTTTTATAA	CAGATAATGT	7800
TACACTCAAG	TTGTAACAGA	ACCAGCTTAT	CCAGCTCATG	AAATGTATGC	7850
ATTTTCATCTC	AACTCTGTAT	TCAGTGACAT	CCTGTGGGTA	TCTGGAAATC	7900
AGCCATGSGT	AGAATATTTA	CCATGGAAAT	TGGCAAAATAC	TAAAAAGCAG	7950
AGCACCTTTT	TTTCTGAGAG	CCAGACCATA	GCTCTTCTAC	TCCATAGCAC	8000
CCATCATTAAC	AATTTTAAAT	TACCTCCACT	GAACAGCTTC	TTCTCTCTCT	8050
TACTTCTTCC	ATATCTGATT	TGAGCTTCTT	AATTTATCAT	GTGAACCACT	8100
CTTGTAATAA	TAAACCCAAA	TCCCTGTTC	ATTGTTCTTC	CTGCTAAAAAT	8150
ACTAAACCTG	GTTTAGTCCA	ACCATATTTT	CTCTCTTTGG	AATCTACAGG	8200
GTGGCCCAAA	AACCTGGAAA	TGGAAAAATA	TTACTTATTA	ATTTTAAATGT	8250
ATATTAATAA	GCCATTTTAA	TGCTTCATTT	CCAGTCTCAG	TGGCCACCCT	8300
GTATAGCTGG	GCTATTGAGC	TCTTGGGGGA	GGAGGGAGTG	GACAGTCTCC	8350
CAGCCACACA	GACTGATGTT	GCACCAAAACA	TTTTTTAGCT	TCCAGACTTC	8400
CCTGGCCCTT	AGTGTATCCC	TTAACTCTCC	ATTTCTCTGC	CTTTCACATT	8450
CTCTACTTTT	TAAAAATCTC	TGACTCCACC	TTACCTTAT	CATTCTTAGC	8500
ACATGACCAT	ACTTCTGCTT	CCCAAGAAA	ATGAGCAATT	ACTTCTTTT	8550
CCTTTTCTCT	CTGTCATCAA	ATCTGCAGAC	ATGTCATGCC	TAAGTCCAGC	8600
TTTCTCTCTT	TCTCTGATCT	CAGTCTGCTT	CTTCCATTTC	TGCCCTGAAT	8650
CCCGTCCCTT	CCCCAACCCC	CAAGGACTTC	GCTCTATCAG	TCACCTCTTC	8700
CCTCTCTCTG	ATCTTCAACT	CCTCCCATTT	TACTGGCTTC	TTCTCTCAAGC	8750
CTTTCCCCAA	GCCTTTCCCA	TCTCAATTAC	CTCCTCGCAC	ATGCCTCTGC	8800
AGAAACCACC	CCGTTTCTTC	CCTCCCTCG	GCAGCCTGTT	CTTCTGTGTT	8850
TGCCCTCATG	ATGGCACCAT	CATTGTGTCA	CTAAAAATCAA	TCTCTCCGAC	8900
ATCATCAATG	GCCTTCTTTT	GTTGGGAAAC	CTAATAAACA	CTTTATCTTA	8950
TTTGTCTTTT	GTTATGGGTT	GAATGAGGTT	ACCCCGAAAT	CCATATTAGA	9000
AGTCCTAAC	CCCAGTACCT	CAGAAATGTA	CTTTATTTGG	GAATAGGGTC	9050
ATTGCAGAGC	TTATTAGTTA	GGATGAGGTC	ATACTGGAAT	GTGATGGGCT	9100
GCTTATCTAA	TATGACTGAT	GTCCTTATAA	CAAGGAGAAA	TTTGGAGACA	9150
GACACGCACA	TAGGGAGAAT	ACCATGTGAT	GACAGGAGTT	ATGGAGTTGG	9200
AGTCAAAAAG	CTATGGGAAC	TTAGGAGAAA	GACCTGGAAC	AAATCCTTTC	9250
CTGCGCTTAG	AGAGGGAGTA	TGGCCCTGCC	ACTACCTTGA	ATTCAACGTT	9300
TCGGCTTTTC	AAAACGTGTA	GACAATACAT	TTCTGTTGTT	CAAACCAATT	9350
AGTTTGCAGT	ACTCTGCGAC	TGCAGCCCTA	ACAAACTAAT	ACAGTCTCTT	9400
GGAGGCAATT	GGCAAGGTTG	ACAAATGGAAG	CACCTTCTTA	CCCTTTTAGG	9450
TCTGTGCGCT	TTCTTGTGG	GGGGTGTGTT	CTAACAATTC	CTCTCCATCT	9500
CTCTCTCTCT	AGTTTGTCTT	AAACATTGGT	GTTCTTCAGA	CTTCTGACCT	9550
AGGCCTTCTT	TTCACTTCAC	ATATTCCCTT	GGGTGGTCTC	ACCCACTTCC	9600
AGAAATTAAT	TAAATTAATG	CTCATGCACT	ACTGTGCTGG	AAACTGTTTA	9650
ACAACCTGGCT	CTCTGGGAAG	AGGGGAGACT	GTTTGTATGGT	TTTGTCTGAT	9700
TTCTGTGGTG	TAAATACTCC	CTCCATGGCC	AATTCCAAAC	TGCCAACAGT	9750
TTAACCAACTG	GCTCAAAAT	TTTCTCCAAA	TTTAACTATT	GGCTTTTACA	9800
GGCCAAACAAC	GTGGTACAGC	CAACTCCAGC	ACACCTCTGC	TTTTGTGTCA	9850
GAGAGAAGTA	ACTTATTTTT	GTACAAAAGG	TAAAAATAAA	ACACCTGCAG	9900

GCCCCCTTTT	TTTCCCTAAC	AAACTGCTCT	AGAAATAGAA	TAGCTGAAGC	9950
TTCTTTTATG	CATTCATCTG	TTAITTCCAT	GTCACGTGGG	TGGTGGGATT	10000
ATTTTTCCCT	TATTTTTCTT	GTATATGGTT	GAAATACTGT	ACCTTTGATC	10050
AGTTTTAGTT	TTAIGGCATG	TTTTGCACCC	ATATTAAATC	TAGTTTTTGT	10100
CAGAGGGCGT	CAATATTATT	TTCTCAAAAC	AAGAAAATAT	TTCAATTGCAA	10150
AGGAGACAAA	CAAAAAGGTC	CTTAATACCA	AAACTTTGAA	ATGTGATTTC	10200
TTGTACTTGG	CAGTGTCCAA	GTGGTAAACC	CAAACAGTAT	TGGGTTTTCA	10250
TTTTGTTCAG	GAAAGTCTTT	GTCTGGCAGC	GACTTACCCT	TACATCAGGC	10300
GGGCCTTGCT	CATTCAATCA	CTTAAGTATT	TATTAAACAC	CAGCGGTGTG	10350
CCAAGTACTT	ATCTAGGTAT	CGGTAGATT	CTGATAAGTC	AGTCAGGTCC	10400
CTGCTCTCAG	GGAGCTTGCA	GCAGAGATGG	GGGCTGCAAT	AGAGAGTAAG	10450
CCAAGGAAAT	GAAAAAGGAA	GTTGATTTC	GAGAGTGATG	AATGCTATGA	10500
AGAAAATGAA	GGCAGCGCAG	TGTGATGGAG	AGTGACCCAA	GGTGGTACAG	10550
TTTGTACCTC	TAAGGACCAG	ACTGTGACCC	AGGTCACTCA	CAGATGCCCG	10600
TCATGTGATG	CACACGCAAC	TTTTCCAGGT	GCTCGTTTCC	TCCCCTTCC	10650
CAGTCTCTTG	CCCAGCGCGG	ACTGCTTACA	AATACAGCTA	GAGGAATCTA	10700
AATGAGGTTT	CTCTATCATC	AAACCCAAATC	AAAATGCCAA	GGAACAGAAT	10750
CAGTGCCTGG	CTGAAGGCAG	TGGAACAGGG	CCAGCCTGGA	GTGGTTCTCT	10800
CTGAGGAAGT	TCCTCATCTT	GGTTTTAGGG	CCATACCTTG	TGACCTGTGA	10850
GCTAGGGGTT	GCCAGTCCCT	GACATTCTTA	CTGAGGACTC	GCCTGTCTAT	10900
ATTCCCCGCC	TGTATGTGTC	TCCTGAGTTC	CAGACACACA	GGGCGAAGCG	10950
CCTGATGGAT	GGAAAGTATG	TTTTTGGTGT	TCCATTGGTA	TCTCAAATTC	11000
TACAAAACCT	AGTGCCCTTT	CTCCTCCCTG	TTCCCTCCCA	TCTTCAGTCT	11050
ATCACCTGTT	CCTCATCCAG	CAATGATAT	TACCATCTTC	CAAGGAGCTT	11100
CCCAGGAGTA	ATCCTTGACT	CCTCCTCAAC	ATCCAAATTA	TAATCAAATC	11150
TAGGCCAGGT	ACAATGAGTC	ACGCCATATA	TCCAGCACT	TGGGAGGCT	11200
GAGGCAGGTG	GATCATTTGA	GGCCAGGAGT	TCAAGACCAG	CCTGGCCAAC	11250
AAGGTGAAC	CTGTCTCATT	TAAAAAAGT	TATTTTAAAA	ACTCAAATCT	11300
ATTATTTCTA	CCTCTAAGTG	TGCTTGAAAT	TTATCCATCT	CTCTCCATCT	11350
CTGAGCTGTT	ACCTTACCTC	AGTCCATCAC	GTTTTGTCTA	CGTTAACATG	11400
ACCAGAGTCT	TGTTCTTAGT	CTGGTGAGGT	CACCTCAGCT	GCTTCAGATC	11450
CTTCCATGGC	TCACCGTTGC	CCTCATATAA	AGTTGGCACT	CCTGGACATG	11500
TGGCTTACGG	GGCCCTCCGT	GATGTGGCCC	TATTTGCTTC	TCCATTCTGT	11550
TCTCTCCAG	CCTCTCTGCC	CCCATCTCTA	GGCACCAACC	ACACCTTCT	11600
GCTCGTCAAT	GGTGCCAGCT	TCTCTTCTAT	CTCTGGTCTT	TGGACAGACT	11650
TTTCCCTTCA	CCTGGAATGC	TTTCTTCAAT	CCTACCCAC	TCTCTTAAAT	11700
CTAGATAAGG	TTTTATTCTT	TGAAATGTCT	AGCAGTGAAA	CCATTTCCTC	11750
TGAAAAACCT	TCTCTAACCA	ACCCCTTACC	CTCAGCCCAA	GGTCTAGATT	11800
AGGAGTCCCT	CTGAATGTTT	CCATAGCAAT	TTTAAAGAAT	TGCCTATTTA	11850
CTTGTTTCGT	TCTATCACTA	AACTACAAAT	TGTATGAGAA	CAGCCACTAT	11900
CTCTGCCTGG	TTCAACCATC	ATCTCCAGCA	ACTAGCATAA	TGCCTGGCAG	11950
AGTCAGCCTG	CAACAAATAT	TTGTTGAATA	AATTAACAGA	TGGCTTTATC	12000
TCCTTAAAGTA	AATCTTGCTT	TTTTCACCTA	TTAAACAGAA	CGCACAGGCC	12050
AGGTGTGGTG	GGCCATGCCCT	GTAATCCCAG	CACCTTGGCA	GGCTGAGGTG	12100
GGCGGATCAC	CTGAGGTCAG	GAGTTCAAGA	CCAGCCTGGC	CAACATGGTG	12150
AAACCCCATC	TCTAATAAAA	ATACAAAAAT	TAGCTGGGCA	TGGTGGTGGG	12200
TGCGTATAGT	CCCAGCTACT	AGGGAGGCTG	AGGCAAGAGA	ATCGCTTGAA	12250
CCCAGGAGGC	AGAGGTGGCA	GTGAGCCGAG	ATCATGCCAC	TGTACTCCAG	12300
CCTGGATGAC	AGAGACCCTG	TCTCAAAACA	CACACACACA	CACACACACA	12350
CACACACACA	CACACACACA	CACACACACC	AAGTTGTATA	ATTTAAAAAT	12400
TAACGTGCTT	GTTATGGAAC	ACTTGTAATA	TACAGGAAAG	TAATGAAAAA	12450
GTCTACCATC	TAGCTCACCA	CATAATGACC	ATTGCTATCA	TCCTGGCATA	12500
ATTCTCTCCT	GTATATAAAT	ATATATCTTT	TTATTGTTAA	AATTACACTA	12550
TGAGTACTAT	TTATTTATTT	TACTGTGGCA	AAATGCGCAA	AACATAAAAT	12600
CTTGCCATTT	TAAGGTATGC	AGTTTGGTGC	ATTCACCACA	CTCACATTGT	12650
TGTGCAAAAT	TCACCACTAT	CTATCTCAGA	ACTTCTTCGT	CTTCCCAAAC	12700
TGAAACTCTG	TACCCATTAA	ACAATAGTGC	ATCTCTGTTT	TTCCCTCTCC	12750
TACAATTTAT	TTTTATTITG	GTTTGTACCA	AACTGAAAAAT	AGCTGCTTCT	12800
TCCTTACTTA	GTTTCAGATTA	GCATTTCCAT	TTATTTAGCC	GTGGTTTITGA	12850
GGATGCCATG	ACAGATGCCA	TCCTTCCTAG	AGCTCTTTGG	GGCTGTCAGG	12900
TATTTTCAGT	AGGGTGAATT	CGGGTTGATA	ACATTTTAAA	ATCTCACTTT	12950
ATTCTGAGGT	TCCTAGTGTG	AGAGCCCAAC	GTATTTTITAG	GGACTCCCAA	13000
GTTACAAACA	AAAATATGGT	GAGGAGGAAT	CACCTGAAGTT	TTAACACAAG	13050
AGACTTACAT	TTTGTTCAT	TTCTATCTTT	TAGTTTATTT	CCTAAGCATA	13100
AAGAAATACT	TTGAAAATTT	TACATAGCAT	TATACATATT	TAATTAAGCA	13150
TGAGCACATC	TTAAAACCTT	AAATTTTAGA	TCAGATCTTT	AATTCCTAGG	13200
ATATTAAGAG	GTAATGGCAA	TTTGGCCAGG	TGTGGTGGTT	CACGCCATTA	13250
ATCCCAACAC	TTTGGGAGGG	TGAAGTGGGC	GAATTGCTAG	AGCCGAGGAG	13300
GTGGAGGCTG	CAATGGCCTG	AGATCACGCC	ATCGTACTCC	AGCCTGGATG	13350
ATGAGAAATGA	AATCTGTCTT	CAAAAAAATA	AAAAAATAAA	AAAAAGAGAA	13400
GAAGAAGTAT	TGGCAATCAG	TGCTCCAGGA	ATAATTTCTT	GACTTGAAAT	13450
AAACCTACAT	GTAGACAAAC	TAATTAGGCC	ATTCCAAGAG	TTGCTAGCAT	13500
TGGTTTAAATA	TGTTTTCAGA	GCATTCAGG	AAGCAGTGTG	GCCAGCATTG	13550
CATGTTTGAT	ACTTCAGAAA	TGTATGACAG	GTGTTTCTCT	TACCCAGGTC	13600
TTCTGTTTTT	TTAGTTTTGC	TCATGTAAAT	ATTTATGAAC	ATCCTCATCT	13650

TTTTGAGGGA	AGGGATTATA	GATCATTCTA	ATTCCATTTT	CTAGCATTTG	13700
GTACCATTCT	AAGCACATGA	TAGGCACCCA	TTTGGAGCAT	TTTGGGCTTG	13750
ACAGAATATG	CATTTAGAAT	TGTTCAAATT	AGAGGTGTCA	GTGATGGGAA	13800
TTAGAATACT	ATATAATTCT	AAGTCATTTG	ACTTAAATAC	AAAAGAATGA	13850
TTTTCTTGG	TGGGGAATGG	TGAAGGGAGG	CAGGAGTTAA	GAAGAGGAGA	13900
AGAGATCCTA	AGTCATTAT	AACTTCTCT	GGAAAGACAG	GTGTGTGAAG	13950
ACTTTTTAAA	AAGTCATTCA	CCAAATTGTG	TGTGTGTGTG	TGTGTGTGTT	14000
TTAAATAGAC	TTTATTTTTT	AGAGCAGTTT	TAGGTTTCACA	GCAAAATTGA	14050
ATGCAAGGAC	AGAGATTTC	CATAAACCCC	CTGCCACAC	ACATGCATAG	14100
CCTCCCTCAT	TATCAACATC	CCCACCAGAG	AGGTGTTTGT	TCTAGTTGAT	14150
GAACCTACAC	TGACACATCA	TTATCACCCA	AAGTCCATAG	TTACCGGCAG	14200
GGTTCACGTG	CGGTGTACAT	TCTATGGGTT	TGAGCAAATG	TATAATGACA	14250
TGTATCCACC	ATTATAGTAA	CATACAGAGT	ATTTTCAGTG	CCCTGCAAAT	14300
CCCTGTGTCT	CCACCTATTTC	ATCCCTCCCT	CTCTGCATTT	CCACCCCTAG	14350
CCCTGGGTAA	CCGCTGATCT	TTTTACTGTC	CCATAGTTTC	GGACGATCTA	14400
TTTTTCAGAC	AGACACAGAG	CTGTCTTTCC	CTTAGTTTCT	ATTCTATCAT	14450
TTCTTTCTCC	CCATCCATCA	TAAAAGGCTA	TGAGTTTTTT	TTAAGTGTG	14500
AACACCATCC	TACTTGTCAA	GTTAAACAT	AAGCTCCTGG	CTGGGTACAG	14550
TGGCTCATGC	CTGTAATCTC	AGCATTTTGG	GAGGCTGTGG	CAGAAGCATC	14600
ACTTGAAGCC	AGAAGTTTGA	GACCAGCCTG	GGCAACATAG	CAAGACCCCA	14650
TCCCTCCACA	CACAAACACA	CACACACACA	CACACACACA	CACACACACA	14700
CACACACACA	CACAAAAACA	AGCTCTTGCC	AGAATTAGAG	CTACAAATTG	14750
CCCTCAGGTT	CCTAGAAGAT	CAGTCCTTCA	ATTAGATTCA	GATTGAGATG	14800
CTTCTCTTT	TAAACAATGA	TTCCCTTTCT	ATCATGCCCA	ATAAGAAAAC	14850
AAATAAAAA	TAAACAATAC	TGCCCTGTAAT	CTCAGCTACC	CAGGAGGCAG	14900
AAGCAGAAT	GCTTCAACCC	GGCAAGCAGA	AGTTGCAGTG	AAGTGAGATC	14950
GCGCCACTGC	ACTCCAGCCT	GGGAAACAGA	GCAAGATTCT	GTCTCAAAAA	15000
CAAAACAATG	TGATTTCTC	CTCTAAGTCC	TGCACAGGGA	AATGTTAAGA	15050
AATAGTCCCA	CCAGGAAAGA	AGGAAGTAAG	AATGTTTGG	TAGATTGTCT	15100
TGGAAAAAAT	AGTTATACTT	TCTTGCTTGT	CTTCTAACA	GTTCTCCAAA	15150
GCTTGGTACC	TTGGCCAGAG	GCTTGTCTCC	TGCGTACCTG	AGGTTTGGTG	15200
GCACCAAGAC	AGACTTCTTA	ATTTTCGATC	CCAAGAAAGGA	ATCAACCTTT	15250
GAAGAGAGAA	GTTACTGGCA	ATCTCAAGTC	AACCAGGGTG	AAATTTTTTA	15300
AAGATTCACT	CTATATTTTA	ATTAACGTCA	GTCCGTCATG	AGAATGCTTT	15350
GAGAAAACTG	TTATTTCTCA	CACCTAACAA	TTAATGAGAT	TAAGTTCTCT	15400
TCCCCTCATC	TGACCTGTGG	AGGAATCTGA	ACAAGAGGAG	GAGGCAGTGG	15450
GCAGGTTTCC	TTATCTGAT	GTTTGTCTATG	TTCAGTGTGA	GGCCTCACAA	15500
AAAAAATAAA	AAAAAATAAA	GGCGTCTCTG	ATATAACTGA	GAGCTCATTG	15550
TACAGTAAAT	ATTAATAAAA	CAGTGATTGT	AGCTGAAGGA	TAGAAGTCT	15600
TGGAGGGAGC	AAGTGGGTAG	AATCGCGTCA	AACTAAAGAG	CATTTCTAGC	15650
CAAGAGACACA	ATGATAGATT	GAAGGATATT	TATTCTAAAT	ATAGAATATG	15700
GGTGAACGAG	ATCTGTGGAC	TTCTGGGCTC	CAACGTTAGA	TTCTGATTTT	15750
AGCAAGCTTG	TCAGGGGATT	CTGATATTGA	AAGGCTGTGG	CCTTCACCTG	15800
AGAAACCTGC	CCTAGGGGGC	CATGAAAATT	TGTCCTGTCT	TTCAGAAGTG	15850
CTATCAGACA	TCAAAATGGA	GTTAAATCGT	ATCTTAACAA	TTACTAGGAT	15900
GGGCGCAGTG	ACTCACACCT	GTAATCCCAA	CACTTTGGGA	GGCTGAGGCA	15950
GGAGGATCAC	TTGAGCCGAG	GAGTTCGGGA	CCAGCCTGGG	CAACATAGAG	16000
AGACGTTGTC	TCTATTTTTT	AATAATTATA	AGAGAAAAAA	ATACTGAAAA	16050
TATTGTATAC	ACCACCTGAAT	TATAATAATG	TGTATATAAT	GTATATATTC	16100
ATTATGAGGA	ATATTGATT	ATTTTATATA	TTATATCTTT	TCCTTCTGTT	16150
TATTTTATCC	AGTTATGAG	TATTTAGAAC	AATTCATCAG	TAATTGGGGC	16200
TAAATTGACA	GAATAGTAA	CAGAGAAAAT	AGAAAAAGAC	AGATGGGTTA	16250
TCTTTGAATA	CCAGGTTGGA	GTTGTTTATG	GGTTTGTTTT	TTGTTTGGG	16300
GGCGTTTTTT	TAGACAGAGT	CCCACCTGT	TGCCCAGGCT	GGAGTGCAGT	16350
GGCACAAGCA	TGGCCCACTG	CATCCTTGAC	CTCTTGGGCT	CAAGCAATCT	16400
TCCCACCTTA	GCCTCCTGAG	TAGCTGGGAC	CACAGGTGCA	TGTCACCACA	16450
CCCAGCTAAT	TTTTTTATTT	TTGTAGAGA	CAGTCTTTCT	ATGTTATCCA	16500
GGCTGATCTC	AAACTCCTGC	ACTCAAGTGA	TCCCCCTGCC	TTGGCGTCCC	16550
AAAGTATTGG	GATTATAGGC	ATAGCCACCA	CACCCAACTT	AGTTTCTATT	16600
TAGACTTGGC	CCTTCCACAC	CAGTCATTG	TGTCCAAAAG	ATCTCATAAA	16650
TGTAGACAGG	AACTGTCTCT	TTGCTCATCA	GTTTTCTTCA	TCCTGTGTCT	16700
AGGGGGATGG	TCGGTGGGGG	AAACTGGGGT	TATGCAAGTT	CCTCTGAAAC	16750
ATCCTCTGTG	AGCCAGGGGA	TGGATGAGGC	ACCAGCCGCC	AGCGAGTCAG	16800
TGTGCAGCTT	TCCAGAAAGG	AAGTCATCAG	CCAGTCAGCC	GGCCCTGGCA	16850
GCCAGCACCC	GGCAACCCTG	CTGTCTTGTG	ATAAAGAAAT	GGCTGCCTG	16900
ACAGGATGGT	GTGGATTTTT	CTTTTTCTTT	TTTTTTTTTT	TTGAGACAGG	16950
GTCTGGCTCT	GTGCCCCAGG	CTGGAGTGCA	ATGGCGGGAT	CTTGGCTCAC	17000
TGCAGCTCT	GCCTCCACAG	CTCAAGGCAT	CCTCCCACTT	CGGTCTCCCG	17050
AGTAGCTGGG	ACCACAGGCA	CACACCACCA	CGCCCACTA	AGTTTTCGTA	17100
TTTTTAGTAG	AGGCAGGGTT	TTACTATGTT	GTCCAGGCTA	GTCTCAAACT	17150
CCTGAGCTCA	AGCTATCCAT	CTGCCTTGGC	CTCCCAAAGA	GCTGGAATTA	17200
CAAGCGTGAG	CCACTGTGCC	TGACCAGGGT	GGATTTTTTC	AAGTGCACAT	17250
GTGTGGTCC	CAGAAGCTCT	GATGGTACCA	AATTCCAAGC	GAAAAAAAGT	17300
CAATGGTCC	CACCCATCCT	ACCTCCCATG	ATGGCAAGAG	GAAATCACCA	17350
CAGTCAGAT	ACAGTCCATG	TAAACAAAAT	TGCTATGGAT	TTTGAAAGTG	17400

AACCTTAAGA	GAAGTGCAC	ATGTTTTCTT	CATTAGAGTT	CTCTGGTAAT	17450
TTCCAGCTTT	TTTTTTTTTT	TTTTTTAGAC	AGTGTCTCGC	TTTGTCCGCC	17500
AGTGTACCCC	AGGCTGGAGT	GCAGTGACGT	GATCTCGGCT	CAGTGCAACC	17550
TCCGCCTCGT	GGGTTGAAGT	GATTCTCCTG	CCTCAGCCTC	CTGAGTAGCT	17600
GTATTTTAGT	AGAGACGAGG	TTTCACCATT	TGGCCAGGCT	GGTCTCGAAC	17650
TCCTGACCTC	AAGTGATTGG	CCCATCTCAG	CCTCCCAAAG	TGCTGGGATT	17700
ACAGGTGTGA	GCCACTGCAC	CCGGCCAGTA	ATTTCAAGCT	TCTGAGGAGC	17750
CCTTTGAATT	GTTAAATAAC	TTGTAGCTAT	GTCCAACATA	TCCATGTTCA	17800
GTGTATGTTT	GATATTTCTT	AGGAAACCTG	CCCTTGGTTG	TTTTCTTTGT	17850
GGTAATTTCAT	GAGCCGGCAA	ATTTGACATG	TGTTACAGAA	TATACCTTTT	17900
CTCTGCTCTC	CTACCTCATA	ACCAGAACTT	AATTATCCTG	CTTTAGTCAC	17950
ATAAATAGCT	AACATAAATA	ATATATGAGA	TTTCAGTCTG	CTCAGTGTGA	18000
AAATAGACCT	TCTAAATGAT	CTCTTCCACT	TGCAGATATT	TGCAAAATATG	18050
GATCCATCCC	TCCTGATGTG	GAGGAGAAGT	TACGGTTGGA	ATGGCCCTAC	18100
CAGGAGCAAT	TGCTACTCCG	AGAACACTAC	CAGAAAAAGT	TCAAGAACAG	18150
CACCTACTCA	AGTAAGAAAT	GAAAGGCACC	CTAGAGATGT	TCCAGCCCCA	18200
AAGATATTTG	AATAGGTTGG	ACTCGGGCAC	CAATCTAGCA	AGTCTACCGG	18250
AAGTTGTATA	AAGCTGAAAA	TACTGAAGCA	TTTCCCAAAAT	GGGAAATCCT	18300
AAACTCAAAA	CTTGCTTTTT	GGTTTTTTTG	TTTGTTTGTT	TTTTCTTCAT	18350
CTGACATTGC	TTAGTAGTCA	CAGAAATGAA	GATAAATCAA	TCATTTCATGA	18400
TCTAACCAATG	ACCTTCAGTG	CTCTAAAAAA	CTACGGAGTC	AAGGAAACA	18450
TGAATATATT	CCTCATGTAA	AATTAAAAATA	CAGACATATA	AAGGGCAAAA	18500
CATGAACATC	ATTCAATCCT	TGAGGTCCGT	CCCCCTCCCA	GAAATAACCC	18550
CCAGTATGCC	TTGGTTTAGA	GCATTAAAGCA	GGAGGGCCCT	GAGTCACTCC	18600
AGACAGTCTT	GACCAACCAAG	CAGCATTCTC	TTTTTGTTC	CTCTGTGGCT	18650
TTTGCAACA	CAGGGCTAGC	TCAGCTACCC	ATTAGTATGT	TTTCAGTCAC	18700
TAAACAGTC	TTCCAGTCTT	CAAAATAGGA	TGACATTGTC	ACATGGGGCT	18750
TAAAGCAAG	TGAAACAAGG	AACCCCTTT	TTTTTTTTTT	TTGAGATGGA	18800
ATCTCACTCT	TGTCGCCAG	CCTGGAGTGC	AATGGCGCAA	TCTTGGCTCA	18850
CTGCAACCTC	CACCTCCAG	GTTCAGAGA	TTCTCCTGCC	TTAGCCTCCT	18900
ATTCATTATG	AGGAATATT	GATTATTGAG	TTCTGTAGG	GTAAAGATAT	18950
TACCCCGAT	CATATTATTG	ATTATTGAGT	AGCTGAGATT	ACAGGTGCCT	19000
GCCACCAGA	CCGGCTAATT	TTTTGTATTT	TTTAGTAGAG	ACAGGGTTTC	19050
ACCATGTTGG	CCAGGCTCCA	GGCTCGTCTC	GAACCTCTGA	CCTCAGGTGA	19100
TCCACCCACC	TCAGCCTCCC	AAAGTTCTGG	GATTACAGGC	GTGAGCCACC	19150
ACTCCTGGCC	ACAATCTTTT	TTTAACTATG	AAATATATTT	TTATCTGAAG	19200
TTTGATGTTT	ATACCCAAT	GAGGGATGAT	GTTCCTATAT	CTCAGTTAAA	19250
GAAATAACCT	GCTCAGATAC	TTCAAGCTCT	TCTTTTGACT	TTTGAAATA	19300
AATGATCTTG	AAGTTACTAT	ACITTTGTTG	GGTTAGTTAA	CATTATTATA	19350
AGTATATTAT	TTTAATTAAT	TATCTTTGTA	AGATTTTACT	GTATACTACC	19400
TGGAGTTCAA	TGATCAGAT	GGATTTCAA	TTTATGTACA	TTTTTTATGT	19450
ATATGGTACA	GAAAAAATG	TGATCCATAA	GAAATCAGAA	AATAGCGCAT	19500
ATGCTAATAG	CTAATGTTGT	CCTCTAAAAA	ACTTATTTTT	GCATTTTTAA	19550
GAGGGGGATA	TACTCTGACA	CTTTAATAAG	TGTAATTAAT	TATTGACTGG	19600
AATTGGCAT	GAGGCAGGCG	CATTTCAGAT	CCCATTAAG	GAATGACACA	19650
TACCAGAGAA	CCACAGAAGT	AAGGCCACAT	TTGTAATAAA	TCATTATAGC	19700
TCTGCTAGGA	GAGACCCAG	TGTATTAGG	TAATTAATGG	ATTTGCTCTT	19750
AAAACACATG	TCCCGGAAGA	TATAGGTGAG	TCTTGGGGGG	CCGCATTAAA	19800
CATTATACCA	ATGTATCTTA	CATTCTAAG	AAAGTTTTAC	TACTTTACAG	19850
GATCTTTCTG	TTACCAAAAT	GGAAGGTTTC	CAACTCCAGG	ACTTGGCTTT	19900
CATAGTTCTT	ACACCAGGGG	AAATGCCTTC	CTTTGCTAAC	TATGCAACCA	19950
GGTTAGTTAG	TGTAAGTCCA	GCCACCCTGT	TGGCAATGCT	AAAAGGTACA	20000
ACAAACACAG	AATTTTATTT	GCATTGTAA	ACATTTGATT	TCTGGCTCGA	20050
AATTTTCAGT	TTTCATGGGC	ACGTATGGA	AACAGAAATC	TTCTGTGTTT	20100
AGTTTGGCCA	CCTACTCATT	GTAGTGACAA	ATATTTTCAGA	AGCCAATAGG	20150
GGATTCCACA	AATTGTTCTG	AACCTGTGGC	TGAGACTGGT	AATGGCTGAG	20200
TGACATGGGG	ACATACCACA	AAAGAAGAGG	TAGCAAAAGG	CTGCTGAGAT	20250
AAGGACATGT	TCATTGCTTA	GCTAGTGGCC	TGCACCCTTA	AAACACATGT	20300
CCCAGGCTGG	GTGCTGTGGC	TCACGCCTGT	AATCCCAGCA	CTTTGGGAGG	20350
CTGAGGCGGG	TGATTACCTT	GAGGTACAGG	GTTCGAGACC	AACCTGGCCA	20400
ACATAGTGAA	ACCTCATTTT	TACTAAAAAT	ACAAAAATTA	GCCAGGCATG	20450
GTGGCGGGCG	CCTGTAGTCC	CAGCTACTCA	GGAGGCAGGC	AGGAGAATTA	20500
CTTGAATCTG	GGAGGCAGAG	GTGTGTGTA	GCCGAGATTG	CGCCACCGCA	20550
CGTAGCCTTG	GGCGACAAAG	TGAGACTCTG	TCTCAAAAAA	ACAAAAACAA	20600
AAAAACAAACA	AACAAAAAAC	AACAACAACA	AAAAAACGGG	TATCCCAGAA	20650
GATACAGGTA	AGTTTCTTAA	CACAGGTCCT	CTGTATGGT	GCGTTCCACT	20700
TAAGTAGAAG	ATGACAAAAA	CATTGTCTAT	GAGAATATAG	ACTCACATTT	20750
TAAACCTGTT	TGAGCAGGAA	AAGGAAGCAA	TGTTACAGAT	GTAATCTCTG	20800
GTGTGACTGC	AGAAAGGATG	ACTCCCTTAT	TAAAGTAGTC	ATCCTGAGTG	20850
AGCTAACTCT	TTGTACTTCC	TCTTCTCTC	CTGTCCCTCT	CATCACCCCA	20900
TTCTTCCGTT	GCCTACACCC	AGGCCACAT	TGGATGCTGA	CATAGACTTA	20950
CATGGTACAG	TCCAAGGGAA	AGATCTGCCA	TTTTTTTCAA	TGTGTCTCTT	21000
TGGTTATCTT	CATTCCAAGG	ATCTCTCCAC	TCTTTATACA	GTAAGAGATG	21050
AGAGTCTGGA	AAGGATTGGG	AATAAGATAA	TGAATTGTAA	GTTTTAAATT	21100
GTTCTTCGTA	TTTTGGGGAA	GGAGTAGGCT	AGGTGGTCCT	TCTGTTTTTT	21150

TTTTGTTTTT	TTTTTTAAAG	TAGATGTGGC	CAGACGTGGT	GGCTCACGCC	21200
TGTAATCCCA	GCACTTTGAG	AGGCTGAGGC	AGGTGGATCA	CTTGATGTCA	21250
GGAGTTCAAG	ACCAGCCTGG	CCACACAGT	GAAACCCCGT	CTTTACTAAA	21300
AATACAAAAA	CTAGCCGGGC	TTGGTGGCGT	CCACCTGTAG	TCCCAGCTAC	21350
TGCAGAGGTG	GAGGCAGGAG	AATCACTTGA	ACCCGGGAGG	TGGAGGTTCG	21400
AGTGAGCCAA	GATCATGCCA	TTGTACTCCA	GCCTGGGCGA	CAGAACAATA	21450
CTCTGTCTCA	AAAAAAAAGA	GAAAAAGAAA	GAAAAAAGA	ATGGATTGTA	21500
ACTCAGTCGT	CAATAGCCTC	TATTCAGGA	GATGTTACAG	TTGATTATGT	21550
TATAGGGGGT	GTATAATAGA	ATTTGAGCT	ATGTAAATTC	CAAGTGCAAT	21600
TGGAAGAAATG	AAGAAATGGA	GGAAGGGTAA	AGTATGAGTG	CAAGCAATCC	21650
AGGTTTTTTG	AAAATGCTAT	AATCTTTGTT	CAGGGCTAGT	ACAAAGTGCT	21700
ATTTAGCTGT	AAGGGTTTTT	TGTGATTAC	AGACAGTTTT	CACATGTGTC	21750
ATTTCAACCT	TGGTTTTATG	GCGAAGGCAT	GTGATGGTGC	TTGTCCCAGG	21800
ACTTTAGATC	CATATCTGAG	GTTCCTGTCG	GGCAAAGATA	TTACCCCTGA	21850
TCATATTATA	GTCTATAAGT	GGGAGAGTTG	TGCCTGGAGC	TCAAGTCTTA	21900
TGATTTCTGA	TCCAGGGCAC	TTCTTACAAC	ATGATTTTGC	AATATAAAAAG	21950
CCCTATAATGT	GTGACTAAAG	CAGGTCACTC	ACCCCTTGTA	ACAGACTCTA	22000
GTAAATGTAC	TGCCACCAAA	CGGCTGCGTG	ATATTGGGCA	AAGACTTACC	22050
TTATTTGAAT	CTCAGTTTCC	TCCTAGAAAA	ATGAGGGTGG	AGGTAAAGCA	22100
TAGGCTGATG	ATCCTAAAGC	CTCCATACTG	CCCTAAACTG	TGGCTCTAAG	22150
ATCCAGTAGA	ATGCTGGGTC	ACAGGACTCT	AGGGAGCTTT	TCAAACCCAA	22200
ATGTCTGTCA	TTCTTGTATG	GTAGGCAGCA	GTTTATGGAA	GTGGGCGACA	22250
CAGCAAAATAT	CAAAATACCT	AAAGCAGCTT	GCAAGAGTTG	TTTCTGCCTA	22300
GTGGTCTTTA	TAGTTAATAT	TAAATAGTTA	ATTTTTTTTT	TTTTTGAGAC	22350
AGAGTCTTGC	TCGTTAACCC	AGGCTGCAGT	GCASTGGCAC	AATCTCGGCT	22400
CACTGCAACC	TCCACCTCCC	GGGTTTGAGC	AATTCTGTCT	CAGCCTCCCA	22450
AGTAGCTGGG	ACTACAGGTG	CATGCCACTG	CACCCAGCTA	ATTTTGTGAT	22500
TTTTAGTAGA	GACGGGGTTT	CACCATATG	GGCAGGCTGG	TCTCGAACTC	22550
TGACCTCAG	GTGATCCACC	TGCCTCAGCC	TCCCAAAGTG	CTGGGATTAC	22600
AGGCATGAGC	CACCTGACCC	AGCTTAAATA	GCTAATATTT	AATATTATTC	22650
TATAGTTAAT	CAAGTAAATC	AGGCCAAAGA	CTTAGAAACA	AAACAAAAG	22700
CCACTTTTAA	GGAGAAAGGG	TGTAAGTTTG	CCAGATAGAT	AGAGATCTTT	22750
CTTTTTTAAC	TACAAGAGTT	CAGGAATGAA	TTACTCTTTA	ACAAACGACT	22800
ATAGATATAC	ATGAAATTTG	GAAGGACTTA	TTATGCATAT	GATAATCAAT	22850
TAAAGACAA	CACCTTAAAT	TATATTGTTG	CACTCTCAA	AAAGTGGTAA	22900
TAGAACAGCT	AATGGTTTAA	AAAGCAGAGT	ACAGAAGTTC	CCAACTTAT	22950
GGCACCTTAA	TATCGCAGAA	AACTTTTTAA	AGCATGCCTA	GGCCACAAAA	23000
AATACCTGTA	TTTTGATTAT	TAAATTGTAA	GGTCTACACA	ACCTAATAGT	23050
AATAGGTCCA	ATAGTATGTC	TGTCCAATAG	ATGTTGATGT	TTTTTTCCTT	23100
GCAAACTTAA	AAGATCTTAC	AGTGCCTCTG	TAAATAGCAC	TGCCTGGTTA	23150
GAGTTGAATT	TCAGATAAAT	AATTTTTTTC	ATGTTAATTA	TTTTTCTTTT	23200
CTTTACTTTT	TTTTTGTGTT	TTTTGTTTTT	TGTTTTTTTT	TTTTGAGACA	23250
GGGTCTCAAT	CTGTGCCCCA	GGCTGCTGTG	CAATGGCATG	ATCATGGCTC	23300
ACTGCAGCCT	TGACCTCCCT	GGGCTCAGGT	GATCCTCCCA	CCTCAGCCTC	23350
CCAAGTAGCT	AGCTGGGACT	ACAGGTGCTT	ACCATCATGC	CCGGCTAATT	23400
TTTGTGTTTT	TTGTAGAGAT	GTGGTTTTGC	CATGTTGCC	AGGCTGGTCT	23450
TGAACCTCTG	GGCTCAAGTG	ATCCGCCCGC	CTCGGCCTCC	CAAAGTGCTA	23500
GGATGACAGG	CATGAGCCAC	TGCACCTGGC	CCCTGGGCGA	AGTATTTCCT	23550
AATGGTTACA	TAGGACATAC	ACTAAACATT	ATTTATTGTC	TATATGAAGT	23600
TCAAGTTTAA	CTAGGTGCC	TGCACTTTTA	GTTGCTAAAT	CCTGTAGCTG	23650
TACCCATGCA	TTCCTGGTG	CTCCCCAGCT	TGCCTTGAC	AGAGTTTGGA	23700
AACCATAGTC	CTATACTCT	AGGCCAATTT	TTTAATGTAA	AATTTGATTC	23750
ATTTTAAATT	AATAAATAAT	AACAGGAATT	TTTTTAAAAA	TGTTTTTAAA	23800
TATAATTAAA	ATTATCAAAA	TATTTTTTAA	CTGAACITGT	GACTAGAGAT	23850
ATTTAGATT	TGAAGAGTGG	GGTTTTATGCT	AACTAATGAC	AGTCTGGCTA	23900
TGCATGTGGA	GCACTGAGCT	ATAAATTGTG	GCTTCCCAA	TTCTCCTGAT	23950
GTCACTTGAA	CAAAACCTAA	GTGTGAGACC	AGAGCTTCTG	GTATCTTCCA	24000
TGGGATTICA	TTCAACAGCT	GGAGCAAATG	AAGTCAGATT	GATTTTTTTT	24050
AAATTGTCCA	ATTTTGTGTT	CTCAAAAAACA	TAATTATAAT	CATTTATTAG	24100
AACTAGAATT	TCTTCAGTTT	AACAACAGAA	ATAGTTATTC	ATTATGAAAA	24150
GCGAATCTGG	AGGCCTTCAT	TGTGGTGCCA	ATCTAACCAT	TAAATTGTGA	24200
CGTTTTTCTT	TTAGGAAGCT	CTGTAGATGT	GCTATACACT	TTTGCAAACT	24250
GCTCAGGACT	GGACTTGATC	TTTGGCCTAA	ATGCGTTATT	AAGAACAGCA	24300
GATTTGAGT	GGAACAGTTC	TAATGCTCAG	TTGCTCCTGG	ACTACTGCTC	24350
TTCCAAGGGG	TATAACATTT	CTTGGGAAC	AGGCAATGGT	GAGTACCCCA	24400
GGGAACAATT	CATTAATAAG	GAGATTCCTC	ACTAGCATT	TTTCTTTTCT	24450
TTTCTTTTCT	TTTTCTTTT	TTTTTTTTTT	GAGACAGAGT	CTCGCACTGC	24500
TGCCCCAGGCT	GGAGTGAGT	GGGCCACCT	CGGCTCACTT	GAAGCTCTGC	24550
CTCCCCAAAC	GCCATTCTCC	TGCCCTCAGC	TCCCGAGTAG	CTGGGACTAC	24600
AGGCACCCGC	CACCGCGCCC	GGCTAATTTT	TTTTTTTTTT	TTTTTTTTTT	24650
TTTTTTTGCA	TTTTTAGTAG	AGACGGGGTT	TCACCGTGT	AGCCAGGATG	24700
GTCTTGATCT	CCTGACCTCG	TGATCTGCC	TCCTCGGCT	CCCAAAGTGC	24750
TGGGATTACA	GGCGTGAGCC	ACCAGGCCCG	GCTAGCATT	TTTCTTATGA	24800
CACTTTTTTT	TTTTTTTGA	GACGGAGTCT	CGCTCTGTG	CCCAGGCTGG	24850
AGTGCACTGG	GCCTATCTCG	GCTCACTGCA	AGCTCCACCT	CCCAGGTTCA	24900

CGCCATTCTC	CTGCCTCAGC	CTCCCGAGTA	GCTGGGACTA	CACGCACCCG	24950
CCACCAACGCC	CGGCTAATTT	TTTTGTATTT	TTAGTAGAGA	CGGGGTTTCA	25000
CCGTGTTAGC	CAGGATGGTC	TCTATATCCT	GACCCCATGA	TCTGCCCGCC	25050
TCGGCCTCCC	AAAGTGGTGG	GATTACAGGC	GTGAGCCACT	GCGCCCGGCC	25100
AACACTCTTT	TTATTATTAG	CAAATATACT	TCTGCCTGGG	CACATTCTTG	25150
CAAGTGCTCA	ACAATGCAAC	TTTTGGAAGT	GCATGTGGCA	GAAACTCCTG	25200
CTGTATTTAT	TCCAGAACCT	ATTATTGCTA	ATCCCACTTT	ATGTTACATT	25250
TGAAGTGAGA	ACCAGTTGGA	GCCAGCAACG	TTCCCACTC	CAAAGTTCCC	25300
TTGAGATTTT	CAGAACTACT	TAACCCCTATT	ATGCTTGGCA	ACCTGGACTC	25350
AGCAAACTG	GGAAGTCAGC	AGTTTGTTTT	ATTTCATCCCT	TCCTTTCTCA	25400
GTTTCTCAAA	TGTGTCACTT	AATCTCAGTA	ACCCCATTCG	AACCTTCATT	25450
ACCTGCCCAA	GCGGCTAGA	ACTTGCCAGT	ATAGAATCCT	ACGTGGGTCA	25500
AGCTCCTGAC	TGTCTCCTTC	TTCACCTCTT	TTTTGCAAA	AACCTTGAAA	25550
TTTTAACTAT	AAGTATTCAT	GATTCGCCAC	ATTTATTCAA	AACATAGAGT	25600
GCTTTTTCCA	CATATCAGCC	AATGGAAATA	AGGATTAAAT	GGGAAATGAA	25650
ATGTAGTAAT	AGGATAAGCA	CAAGTCTTCT	TCCTGCTCAA	ACTTTTTTTT	25700
TTTTTTTTTT	CAGACAAGAT	CTTGCTCTGT	TACCCAGGCT	GGAGTGCAGT	25750
GGCGTGTTC	TAGCTCAATG	TAACTCCCAA	CTCCTGGGCT	CATGCAATCT	25800
CTCACACCTC	AGCCCCCTGA	TAGCTAGGA	CTACACTATG	CCTAGCCAAT	25850
TTTTTTTCTT	TTGTCTGGTT	GTGTGCCCCA	GGCTGTCTCG	ATCTCCTGGC	25900
CTCAAGTAAT	CCTCCTGCCT	CGGCCCTCTA	AAGTGTCTGG	ATTATAGGCA	25950
TGAGCCACTG	TGCCCGGTCT	CAAACTTTTT	TTTCCAAAGT	AAATGAAAGT	26000
ATTAGATATG	GAATATAGTC	TAGTTCCAG	ATATCCATAT	CCATTGGTTT	26050
ATTACCTCA	TTATTAACCT	CAAATTGTTT	AATAGACCCT	CATATCTCAG	26100
TTATACAGTT	AAAAATTTTG	TTTTGTTTTT	CTGGAGTATC	TTATTATATA	26150
CTATGAGTTT	TACTTTACTT	ATTTATTTTA	TTTTTTGAGA	CAGACGCTTG	26200
CTCTGTCACT	CAGGCTGGAG	TGCGTTTGG	TGATCATGGC	TCACTATGGC	26250
CTCGACCTTC	TGGGCTCAAG	TGATCCTCTC	CCTCAGCCTC	CCAAGCTGAG	26300
ACTACAGGCA	TGCACCACCA	CATCTAGCTA	ATTTTTTTTT	TTCCCCATGG	26350
AACAAGGCTT	TACTATGTTA	CCCAGAGTGG	TCTCAAACTC	CTGGCCTCAG	26400
GGGATCTCC	TGTCTCAGCC	TACCAAAATG	CTGGGATTAC	AGGCATGAGC	26450
CATAGCGCCA	GACCTGGTTT	TACTTTTTCT	GACTTTGAAT	TACAAGTTTT	26500
TGTAATTTGG	AAAATGTTTT	GTTGCTTTTA	AATACTGCTG	TATGTTTGCT	26550
TTTAAATACA	ACATTTCTCG	ATATATATTT	TGAGAATTGC	TGCTTTTCAG	26600
AACCTAACAG	TTTCCTTAAG	AAGGCTGATA	TTTTCATCAA	TGGGTCCGAG	26650
TTAGGAGAAG	ATTTTATTCA	ATTGCATAAA	CTTCTAAGAA	AGTCCACCTT	26700
CAAAAATGCA	AAACTCATGT	GTCTGTATGT	TGGTCAGCCT	CGAAGAAAGA	26750
CGGCTAAGAT	GCTGAAGAGG	TAGGAACCTAG	AGGATGCAGA	ATCACTTTAC	26800
TTTTCTTCTT	TTTCCTTTTG	AGACAGAGTC	TCACTCTGTC	AGCCAGACTG	26850
GAGTGCAGTG	GTACAACTAT	GGCTCACTGC	AACTTCGACC	TCCCAGGCTC	26900
AAGCAATCCT	CCCATCTCAG	TCCCACAAAT	AGCTGGGACT	ACAGGTGCAC	26950
ATCACACAC	CTGGCTACTT	TAAAAAAATT	TTTTGTAGA	GATGGGGTCT	27000
CCCTGTGTTG	CCCAGGCTGG	TCTCTTGAAT	TCCTGTGCTC	AAGCCATCCT	27050
TCCACCTCAG	CCTCCAGGAT	TGCCAGGATT	ACAGGCATGA	GCCACCACAC	27100
CCAGCCACCA	CTTTTCTTAA	AAAAAAATAA	AGATTCTCTC	TGGTAGACAA	27150
TCCTCAATAG	TCCACATGTT	ATTAAACAAT	CTGCTGCCTG	AATACATGAT	27200
TTACCAAAAA	AAGGAAATTT	TGACGGGTTT	AGAAATATCA	GGGATCTGAG	27250
GCAAAATGTA	CCTATGTATA	AATTTGCTAT	CAAAATTAGG	AAGTTTGTGT	27300
TTACCTGATC	CTAAAGCAGT	AACCAGCCCA	TTTCTAGGGA	ATAAACTCT	27350
CATGCGTATA	TTGTGCATAT	ATATGTATTA	TATGACTGAG	TGATAATAAA	27400
ATTTTTTTTC	TAGCTTCTTG	AAGGCTGGTG	GAGAAGTGAT	TGATTCACTG	27450
ACATGGCATC	AGTAAGTATG	TCTCCTATTC	TTAATACTAG	GAAAGTAAGG	27500
CTAGCTTTAT	TTATTACCTA	GTATTCAAAA	AGTTAGTTCA	TTAACTGCC	27550
AATTGACTGC	AGTTCAAATA	AGAAACAAAT	AGTGTCTCAA	GTAGCACTGT	27600
ACTCCAATTT	TAATATTAAAT	AAAAAAATTT	TTAAGTTATT	TTAAATAATG	27650
TAGTGGTTTC	TATAAAGATC	ACTTTATACA	GAAGAACAGT	GCCAAATTAAC	27700
CCATGGGAACA	TATAAGTAGC	TAAAACCAAT	TGCTTGCCAA	AGAACCAGTA	27750
ACCCAGGAGT	ACATGTCTTT	GCCACTGTGT	TTTTTCAAGA	CAGAGTAAC	27800
GATTTCTAGT	TACTTGCTATA	GAATGGACTC	CTCCTCATAA	CTCCCTTCCA	27850
TCTTGGTCTT	TCCCTAGTAG	AACCTCTACC	TTTTTTTAGT	AACAGGTGAG	27900
TGGGAGAGGT	AAGAAGGAGA	ATAAGGTCAG	CAATTAACTT	AAAAGCAGAA	27950
AGTAAATTTT	GTTATTTTTT	TTCTGAATAT	TTCTGTGTA	ATTTAGCTAC	28000
TATTGGAATG	GACGGACTGC	TACCAGGGAA	GATTTTCTAA	ACCCTGATGT	28050
ATTGGACATT	TTTATTTTCT	CTGTGCAAAA	AGTTTTCCAG	GTAATAGTCT	28100
TTTTAACTTT	TTTAAATGTA	AACCAGAAATC	CTTATTTTAT	AGTCTAGCTA	28150
GTTCTAAAT	CTATAGGTAT	GTATATTTAC	ATGTTTTTCT	AATTTTAGAG	28200
AACAAGCACT	ATGACTTATC	CACGTGTAAT	TTCCCTTCTA	GCATTGGGTC	28250
TTACCCCATG	TACGTGATTA	GAAATTTGAA	ATATTTCCAA	TAGCCTTTAG	28300
TAGAATTAAC	TCACATAGAT	GATAAGAAATG	GGTTGGTTCA	CTTCATGTTT	28350
CTTCCACAGC	CTACTATTTT	AATAAAAGAA	AGTTTCCCAA	GACCTAAATG	28400
ACTATGAACA	TATTTTATAA	CTATATAGGA	GGGGTGGGTC	TAGGAATACA	28450
AAGTTTGA	TGCTGTTAAT	CTTCAACACC	ACAGTTGAAA	CCACAGGTCA	28500
GCTTTTTTGC	AATTACCATG	GATACCTTTT	TGTTCTATAG	GTGGTTGAGA	28550
GCACCAGGCC	TGGCAAGAAG	GTCTGGTTAG	GAGAAACAAG	CTCTGCATAT	28600
GGAGCGGGAG	CGCCCTTGCT	ATCCGACACC	TTTGCACTG	GCTTTATGTG	28650

AGTGAAGCAG	CGCTGGCCTT	AGGGGTGAGA	GTGCAGCTCT	TCTCCATCCT	28700
TCTATTCTGC	TGAATAGCT	CCCCAGCCAA	AAAGCAGATC	AAAGACCGTT	28750
TCAGTGGCTG	AGCCCCAAA	TTATGCCAG	ATTTTGCAAG	AAAATGATT	28800
ACTAAGCTT	GAGGGACATC	TTTAAACAAT	GTTCCAAAT	AATCACTATA	28850
AGGATGAAT	GTTCAGAAA	TTTTGGCCTT	TAATTATGGC	CCATAAATAT	28900
GTCAAGTAGT	CCTTACTCTA	AAGAAGTACA	CTGTAAAAGA	ATGCATATAG	28950
CCGGATATGG	TAGTTCCTTG	TAATCCCAAT	ACTTTGGGAG	GCCAAGGTGG	29000
GAGGATTGCT	TGAGCCGAGG	AGTTTGAGGC	TGCAGTGAGT	TATGATGGTG	29050
CCACTGCACT	CTAGACTGGG	CAACAGAGTG	AGACTGTCTT	TTTTTTTCCC	29100
CTCTGTCAAC	CAGACTGGAG	GGCAGTGGCA	CGATCTCACC	TCACTGCAAC	29150
CTCTGCCTCC	CGGATTGAAG	CGATTCTCCT	GCCTCAGCGT	CCTGAGTAGC	29200
TGGGACTACA	GGAGTATCAC	CGCACTGGGC	TAATTTTTGT	ATTTTTAGTA	29250
GAGACGGGGT	TTTGACATGT	TGCCCAGGCT	GGTCTGAAAC	CCATGAGCTC	29300
AAGTGATCTG	CCTACCTCAG	CCTTCCAAA	TGCTGGGATT	ACGGACATGA	29350
GCTACCACGC	CCGGCCACAC	CCTGTCTCTT	AAAAAAAAAA	AAAATGCAAG	29400
TTAGAGCAT	TTACAGCTTT	GTCTCTCAGG	AGGATACCTA	GTGTATGTAG	29450
CTATAATTCA	TAGATTCCCA	AGAAGTTTAG	AGCCTAAAGT	ATGAGGTCCC	29500
ACCAGAGGGG	CTATCATTAA	ATTTAAAGAT	TTGTAAATC	ATCTCATTTG	29550
CCAACACCAC	AAACTTGATT	GCTTTAAAT	ACTGGTTTAG	TTACATTTAG	29600
TAACCTATT	AGTGTCTTTA	ATCTATACTG	CTATATCCTC	ACATTGAGAT	29650
TTTTTTTCTT	TTCTCTTCCA	TCTTCATTCT	TTTTTCTCTC	ATCCTCATT	29700
TTATAAGCCT	AGAATACATC	ACAAATCCTT	TATGCCCATG	GAAGCAAGAG	29750
GAATAAGAA	TGGAGATGTT	TGTTTTGCCA	TTAACTAAAG	ATCTGGGGTG	29800
TCGGGGAGAA	GGGGGATAGA	GAAGGAGAAG	TGGGAAGAGG	TGTCCATAAT	29850
AGCTTAGGTG	CAATTCTGCT	TATTTTACAT	TTTACCCCG	CTGACTGCCA	29900
CTTTTCTTCT	AGCCCTCACA	CATTGTTTGT	GCAGGGACCT	CATAGGACCA	29950
GGAATTGTCT	ATAGAGGTGG	GAATTGTCT	CACCCGTAAA	GGGATACCTC	30000
TAGCATGGTA	ATAGTCTTCT	AGGATTGTTT	ATCATATGGA	AAGATGTAAA	30050
GGGAGGGGAT	CTGCTGTCTG	TGCTGTCTG	GCATGCAGTT	GCCATTTTCT	30100
TTAAATGACT	TATTTATAAT	TGATGACACT	TTTCTGGCTT	CCTGTTAATT	30150
CCTCCCTCAA	AGATCAATAA	ACCAGAACCA	GGCATGGTGG	CATGCACTTG	30200
TGGTCTGTGA	ACCACCCAAC	AGGTTCACTT	TGCCTGCTGT	CTAGATAGAG	30250
CCAATTATCA	AGACAGGGGA	ATTGCAAGG	AGAAAGAGTA	ATTTATGCAG	30300
AGCCAGCTGT	GCAGGAGACC	AGAGTTTTAT	TATTACTCAA	ATCAGTCTCC	30350
CCGAACATTC	GAGGATCAGA	GCTTTTAAAG	ATAATTGGC	CGGTAGGGGC	30400
TTAGGAAGTG	GAGAGTGTCT	GTTGGTCAAG	TGGAGATGG	AATCACAGGG	30450
AGTGGAAAGT	AGGTTTTCTT	GCTGTCTTCT	GTTCTCTGGT	GGGATGGCAG	30500
AACTGGTTGG	GCCAGATTAC	CGGTCTGGGT	GGTCTCAAAT	GATCCACCCA	30550
GTTCAGGGTC	TGCAAGATAT	CTCAAGCACT	GATCTTAGGT	TTTACAACAG	30600
TGATGTATATC	CCCAGGAACA	ATTGGGGAG	GTTCAGACTC	TTGGAGCCAG	30650
AGGCTGCATT	ATCCCTAAAC	CGTAATCTCT	AATGTTGTAG	CTAATTGTGT	30700
AGTCTCTCAA	AGGTAGACTT	GTCCCCAGGC	AAGAAGGGGG	TCTTTTCAGA	30750
AAAGGGCTAT	TATCATTTTT	GTTCAGAGT	CAAAACATGA	ACTGAATTTT	30800
TTCCCAAAGT	TAGTTCAGCC	TACACCCAGG	AATGAAGAAG	GACAGCTTAA	30850
AGGTTAGAAG	CAAGATGGAG	TCAATGAGGT	CTGATCTCTT	TCACTGTCT	30900
AAITTCCTCA	GTTATAATTT	TTGCAAAAGC	GGTTTCAGTC	CCAGCTACTT	30950
GGGAGGCTGA	GACAGGAGGA	TTAATGGAGC	CCAGGAGTTT	GAGGTTGCAG	31000
AGAGCTATGA	TCACGCCACT	GCACTCCAGC	CTGGGTGACA	GAGTGAGACC	31050
CTGTCTCTAA	ATAAATAAAT	AAGTAAATAA	ATAAATACAT	AAATAAATC	31100
AAGATGGTGT	GCAATTAGAA	TTGAGCGATT	TTGTTTCCAA	ACCTCAAGAA	31150
AGCTTGGTCT	TGCTCTGTCC	CAGGTGGCTG	GATAAATGG	GCCTGTGAGC	31200
CCGAATGGGA	ATAGAAGTGG	TGATGAGGCA	AGTATTCTTT	GGAGCAGGAA	31250
ACTACCATT	AGTGGATGAA	AACTTCGATC	CTTACCTGT	AAGTGACCAT	31300
TATTTTCTTA	ATTCTAGTGG	AGTAGATTAA	AGTCAACTCA	GGACCTCTGG	31350
TGTTAACCTC	CTATGAACAG	TCAGTCCTCT	CAGTAACCTAG	CCAAATCATG	31400
AGATGATGAA	TTAGAAGGAG	CCTTAGATAG	CATCCAATCT	AACATTTTTT	31450
TGTGTGTTTG	AAGAGAAGAA	ATCAAGAGCT	AGGAATAACT	TTTTAAAGGT	31500
AAGCCATTG	CAGTATAGTG	TGGATTTTGT	TTAAAGGGG	ATAATTTGAA	31550
ATTTTATGAC	TCATTATACA	AGACAAAATA	AGTTGGATTT	TCAAATGTTT	31600
TACAAAGTAA	ATCAAAGTTA	TAATTGCCTA	CAGTACGCAA	AGCTTCAAAA	31650
CATTTTTTAT	GTTATGAAAT	TGTAATTTAT	TTAACCTTAA	AATGAGCCAG	31700
TACCATGTGT	TTGCTTAAAA	ATCTCATGCT	AAGAATTTAC	TATGTTGTTA	31750
ATAATCTTCA	AGATATTTAT	GAATAAAGTC	TTATTTCTAA	TCCTTCTCTC	31800
AACTGTATCT	GGTGCTAAAT	CAGGAAATGT	TTCTTCCCAA	AAAGCCTCGT	31850
GGAAGATCTG	TATGTCTAAA	TATATGTCAG	GGATAATACA	GATGTAGCCC	31900
TGCGAAGCAT	GACCTTGATT	TTTATAGTCT	AAAATGTCT	TTGCAGATAT	31950
CTATTTTCTA	AGAATAATTC	CTAAAAGAAT	TATTTGAATG	TTGTAGGAAA	32000
GCTAAGAAAT	TTTGCAAGA	GCGTACGTGA	AAATATAAGC	TAGGCTTTTG	32050
TGGTTTGTGG	ATAGACTTCC	CAACAAAAT	GCTTTTATC	TATAGTGATC	32100
CAAGCTTGTG	GAACATATTA	GTCATCTTTT	TTTAGAAAAT	TCTTAGAAAA	32150
GTGATCTTGC	AAAAATGGAA	TTTATCTTTC	CCCAAGTATA	TTCTGTCTAT	32200
TATAGAGTTA	AACTAAGCAT	AGTAATTTCA	CCAGACAAAC	ATTCAAAATC	32250
TACTCTGTAC	CTTTTATCT	CATCCAAAT	TTCCAGGGGC	CCAGACATAA	32300
ACCTTTGCCT	TACGAATCT	TTGTATATGC	ACTAAATATG	CTTCTCTTCT	32350
AAGGTTCTCA	GTGAGCTAGA	AAAATGTGCA	AGAGTAAATG	GTACCCCTCT	32400

CACTTGTAGA	TCCAAGAGAA	TTAGACTTAA	ACTCACTCTA	CATGTCTGTG	32450
ACTTTATTTT	ATTTGCATGA	CAGTCCTGTG	AGGTGGCAAG	GCAGGTATCT	32500
TGGATCCATT	TTTTAGATAA	GGAASTTCAA	ATTGAGAAGA	GGTTGCATGA	32550
TTTACAGGAA	GCCATACTGT	AGTCCTATGT	TACTCTTAAA	AAATCCCATTC	32600
AAATCCTGCT	TCTGAGGCCT	GCATACCTTC	TACCTTACCA	GTCAATTGACC	32650
CATGCTTATG	TCTCCTTTGA	AAACATTGAT	TCCACTCTTG	TCTCCAGTGA	32700
AAAAGTGGAA	TTTAAGCAGA	GAACAAAAG	CCATTGTCT	TGTTAAGTCT	32750
ACTTTCCTC	TACTTTCAAG	AAGGAAAGTT	GGGGTATGTG	TTGAATGGTG	32800
ATTTATTTAT	TTTATTATTA	TTTTAAAAAT	TGATACAAGG	TCTTACTGTA	32850
TTGTGCAGGC	TGCTCTCAA	CTCCTGGGCT	CAAGTGATCA	TCCCACCTCA	32900
GCCTCCCACT	GTTGGGATTA	CAGCATGAAC	CATTGTGCCC	ACCACCGATC	32950
CGCAGTTTTT	TAAGAAAAAC	TTTTACTATA	GAAAATTTTA	ATCATATACA	33000
AAATACAGAG	GAAAGTATAT	GAACCCACTT	TAGGAGACTA	GAATATGCCA	33050
CCCCAAAATA	TGCCACTTTG	GCATAAGGAT	TATTTGAGC	TAAAGGCAAC	33100
TGGGAAGAAA	CACATAGAAG	AAAAGTTCTC	TGTCCTTCTC	CATTTGCCTA	33150
AAAGCAGGAC	ATGAATCTTA	AAAGTCCCCC	TCCTTCCCTT	TCTACCAGGA	33200
AAAACAAGAG	TTAATCACTG	AAGATAACTT	CAGACCCCTA	TCAGTGTAGA	33250
GATGGCACTA	GAAGAATCTA	TATTACATAC	TCATTTATTT	TCCTTCCCAC	33300
AACTTGCCAC	CCAGAGACT	AAAAATCCTT	TTCTTTTGTG	ATGTCCTCTG	33350
TCCAAAAATT	TGCTCTATAA	GCTGGAGTTC	TAAGCCACCT	CTTTGAGAAT	33400
TACTTGTTC	CTGGTATTTT	CTGTTAACAT	ACATGTATTA	ATATACATGT	33450
TAACAAGCTT	CTGTTTGTGT	TTCTCTGTG	TTCTGTCTTG	TTACAGAGGT	33500
CCATCCCAAC	TAAGAACTAA	AGAGTAGGAG	GAAAATATAA	TTTCTCTCTG	33550
CATACCTTGA	TCTTGTTTAA	TCCGTAACCC	TTCCCACTTT	TCACCTCCTA	33600
CCTATTAGAT	TACTTTGAAG	CAAAATTCAG	ATATATTACT	TTATCTATAA	33650
ATATTTCAGT	ATGTGCTAGG	TGTGGTGGCT	CACACCTGTA	ATCCCAACAC	33700
TTTGGGAAGC	TGAGGCAGGA	GGATCACTTG	AGCCCAGGAG	TTCAAGACCA	33750
GCTACGGCAA	CAAAAAATCA	AAAACCTTATC	TGGGCATGGT	GGCACATGCC	33800
TGTGCTCCCA	GCTACATGAG	AGGCTGAGGC	AGGAGGATCG	CTTTAGCCCA	33850
GGAGGTTGAG	GCTGCAGTAA	GCTGCATTCA	CACCACTGCA	CTCCAGCCTG	33900
GGTGACAGAG	TAAGACCATG	TCTCAAAAAA	ATACATATTT	TAGTATGTAT	33950
CCTTTTTGTA	AAAACACAAT	ACTTTTATCA	TACTTTAAAT	AATAACAATA	34000
ATTCCTTAGT	ATCACCAAAT	ATTTTGTGAG	TGTCTCACAT	TTTCTCTATT	34050
GTCTAAAAATA	TGTGTGATAG	TTATTCRAAT	CAGAATCCAA	ACAAGGTCCA	34100
TATATTACAT	TGGTTGACA	AGTCTCTTAA	GTTTGTTCAT	CTTTAAGTTC	34150
TTCTTCCCTC	TCTTTTCATCT	CTTGTAATTT	ATTAATGTGA	AAAAACAGGT	34200
AAATTGTCT	ATAGTATTTT	CTACATTATA	GAGTTTGCTA	CATTTATTC	34250
CTATGATATC	ATTTAGCATG	TTCTCTGTG	CCCTGTGTTT	CCTGTAAACT	34300
GGTAGTTATA	CCTAGAAGCT	TGAGTTTATT	CAGGTTTTTA	ATTGTATTTT	34350
TTTTGCAAGA	ACTCTTTATT	ATCTGCTTCT	GGAAGCACAG	AATGTCTGGT	34400
TGTGTCTGGT	TTTGATCTTG	ACAGCTACTG	ATGACCATTG	CCTAATCCAT	34450
TACTTTATTG	GGGTGGGGGG	AATAAGGTTT	TAAAAATAAT	TTTTTTTAAA	34500
GATTTTTTTA	ACTGTTATTT	TGAGACAGTG	TCTCATTTTG	TTTCCCAGGC	34550
TGGAGTGAG	TGGCAACATC	ACGGCTCACT	GCAGCCTTGA	CCTCCTGGGA	34600
TCAGGTGATC	TTCTCACCTC	AGCCTCCTGG	GTACCTGGAA	CTACAGGTGC	34650
ACACCAACCAC	ACCTGGCTAA	TTTTTTGTAT	TTTGTGTACA	GAAGGGGTTT	34700
CATCATGTTT	CCCAGACTGG	TCTTGAACCTC	CTGGGTTCAA	GTGATCTACC	34750
CACCTCAGCT	TCCCAGAAATC	CTGGGATTAC	ACTTTGGCCA	CCGTGCTGGG	34800
CCTAAATGAA	ATTATTTGTC	TCTAAACAGA	CAGAAGTTTT	ACTTTAAAAA	34850
TTTGTCTTTG	TGTGTACATG	TGTTTGTGTA	TGTGTGTGTG	TCTAAAAGTT	34900
TGGCTTTGAG	CTTTGCTTTG	AATTCTTGGA	TGAACAATAA	CCAAGAATAC	34950
TTAAACTCTG	ATCATTCTTG	ACAGATATCC	CCTACAGGCT	ATGGCCTTTT	35000
GAATTGTGTC	CTCCAGTGAT	AAAAAGCAGC	AAGCACGATA	CTGCTCTCAG	35050
ATTCATGGTG	GTACATGTG	AGGTGAAAAA	AAAAAAAAG	ATGAATCCTA	35100
TTTAAATGCC	CCCAGGATAA	CAGTGATACT	CTTTGTAGGA	TAACATTTTG	35150
CTTGCCACTG	GTTTCATTAA	ATAAGGACAT	AAGTAAAGAT	CTATTTTGTG	35200
CTCTTCTCTC	CCAACCACCA	CAACTAGGAT	TATTTGGCTAT	CTCTTCTGTT	35250
CAAGAAATTG	GTGGGCACCA	AGGTGTTAAT	GGCAAGCGTG	CAAGGTTCAA	35300
AGAGAAAGGAA	GCTTCGAGTA	TACCTTCATT	GCACAAACAC	TGACAAGTAA	35350
GTATGAACA	CACCTTTTAC	CAATCATCAA	GTTTTAGTGG	GTAAGCCTGT	35400
AACTTTACTC	AAACACCTG	TTGCATGTG	CTATACATTG	CATAAGTATA	35450
GGCAGTTGCA	ATTTAGTAAA	GTTTATATCA	ACGATTTTAT	TTTATTTTAT	35500
TTTTAGAAGA	AAAATGCTAC	TTTTGTGTTT	GTGTTTTTTT	GAGACGGGGC	35550
CTCGCTCGTC	ACCCAGGCTG	GAGTGCACTG	GTGCAATCTC	AGCTCACTGC	35600
AACCTCCGCC	TCCCGGGTTC	AAGTGATTCT	TGAAGAGGAG	AACAATAATA	35650
ACAACAATAT	TATTTTCAAA	AGTTGTGACC	GCAGTTTCTG	GAGTTGAGAA	35700
GACATCGAGA	TTTTTGTAGC	CTCATACTCT	TGCTTTAGGT	AGCAAAAAAT	35750
GTTCTCTAAAT	CTCAGGAATA	TTCTCTAGAT	AGGTTTCAAT	CTATCATTCC	35800
TGATAAGATG	ATGCTGAAAT	ACTAATCTTA	GCCAAAAAAG	ACCAGCTACC	35850
ATTTCCGATT	GTTGGGACT	GGGAACCTG	GATAGTGAGG	ACCCAGTAG	35900
GAAGTAGCGA	GGGGAATGGT	TTGAATGGAT	AAATTCATAA	AAAATGTCAG	35950
TAGATTTAAT	TTTCTTATAC	ATTTCACTCT	TTTTATAAGG	CTAGGAAAAAG	36000
CCCTGTGTTT	TATGGTTTAT	AATTTGAATT	CACATGAACC	CACAAAAATT	36050
GCCTTTTACC	TTCTTATGTC	TGAAAAATGA	TAGTCTGGCT	GGCTCTTAA	36100
CAACCCAGCT	GGCAGAGCTG	TGAGGATCTC	AGTGTGCTCT	AGCCCAGACA	36150

TTGGTAGCAT	GAACGGCAAC	ATTTTAAATT	GTGTTTTCAA	AATAGGAGCA	36200
CACTAGCGGT	CTAAAAAGAT	CATAAAAGAA	GGATACTAAG	AGGGCCCACT	36250
GTCAATTATGG	ATCCTAATAC	TTAGGATGCA	TTATGGATTG	TCATTATGGA	36300
TACTAATACT	TAGGATCACA	TTTGTAATTG	AGTTTTTAAT	TGCTTAAATT	36350
AGATACATAT	TTCTATTAAAG	TTAACCTCTT	TGCTTTTAGT	CCAAGGTATA	36400
AAGAAGGAGA	TTTAACTCTG	TATGCCATAA	ACCTCCATAA	TGTCACCAAG	36450
TACTTGCGGT	TACCTATACC	TTTTTCTAAC	AAGCAAGTGG	ATAAATACCT	36500
TCTAAGACCT	TTGGGACCTC	ATGGATTACT	TTCCAAGTAA	GTAATTTTCC	36550
TTGTTTCATT	CAAACTTTCA	ATAAATTTAT	TGGTGTTTAT	CAGAATAGAG	36600
AGTTTGGACA	GGGAGCAAAA	GACAAAGTCA	ACTATATCAA	GTTCTAATAA	36650
TTCTTAATAT	TCAGGAAATT	TATGTATGAA	TACTTACTAA	TATGAGTATA	36700
ACTCATCCTA	AGAGTCTAAA	GCAAAAGGAT	GTGAACACAA	ACTAGCAGTT	36750
ATCTTAGAGA	ATAAGTTTGC	ATTTCAAAT	AACCTGACAT	ATCAAGATCC	36800
ACTCAACGCA	TTTAAATTAT	TTACTCTAAA	AAGACATAAT	TCTTGGTAAAC	36850
ACATTCACTA	AAGCAAAATA	TACCTTTATA	TAATTGCTAT	CAAAGGTATG	36900
TGGGTTGGTA	TAAATATATCA	TACCATGTGA	GATCAGTGTG	ATTCTTTTAC	36950
AGCATTAAAT	TTTATTGGTT	AGAGTAAGAA	AAAGAATAGC	TAGAGTATAT	37000
TTCTTAAGTA	GATTCTCATA	CACTTTGGTT	TCAAAAACCA	ATTATTGACT	37050
ACATCTTATA	AAAGCCTGTA	TTCAATGGAG	TGCCAAAAAA	TGACTATGAG	37100
TCTTAAAGAG	TTAGGCATAT	AAATATTTTA	AGGTTTCTGT	TCAATGTATG	37150
TTGGAAGGAG	TTCTTTCTC	ATGACTATTG	TCATATTGGA	GCATAAAAAG	37200
AGTTTACAGG	CTTGGCGCAG	TGGCTCATGC	CTGTAATCCC	AATACTTTGG	37250
GAAGCTGAAG	CAGGCAGATC	ACTTCAGCCC	AGGAGTTTGA	GACCAGCCTG	37300
GGCAATATGG	CAAACTCTC	TCTACAAAAT	ATACCAAAAT	TAGCCAGGCG	37350
TGGTGGTGCA	TGCGTGTAGT	CCCAGCTACT	TGGGAAGCTG	AGGTGGGAGG	37400
ATTGCTTGAG	CCCAGGGGGG	TCATGGCTGC	AGTGAGCTGT	GATGGTGCCT	37450
CTGTCACCCC	GCCTGGGTGA	CAGAGTGAGA	CCCTGTCTCA	AAAAATATAA	37500
TAAATAAAAA	TTAAGAGTTT	ACAAAATTCT	CACCATCTCC	TCCCATCTTT	37550
GCAAATGCCA	CATAAGTGAT	GTGTTCCAGG	ACTATTAGCC	TGGGAACCTG	37600
AGGCAGTACA	GTAAGCAGCG	TTTCTCCAAA	GTCTGTCTCC	CCACAGACAA	37650
ACATTATTTA	CACCTGGGTAC	TGCTCTTTTA	TTTTTTCCCC	TCTATGCTTT	37700
ATTTTACTAT	AACATAATC	ATATAACATG	TAATAGGAAA	AAGGCAGGGT	37750
CGGGGGAGAG	ATCCAGAAAT	CTTCCCAAGA	GCCTTTCCAA	CATAGCCTCT	37800
GTAGACATTT	TTTCTTTCTT	CTTTTCTTTT	TTTTTTTTTT	TTCTGAGACA	37850
GAGTCTCACT	CTGTTGTCCA	GGCTAGAGTG	CAGTGGGGTG	ATCTAGGCTC	37900
ACTGCAACCT	CCGCTCCTG	GGTTCAAGCA	ATTCTCCCA	CTCAGCCTCC	37950
CTAGTAGCTG	GGATTAGAGG	CATGCATCAC	CACGCTGCG	TAATTTTTGT	38000
ATTTTTAGTA	GAGATGAGGT	TTCAACCATG	GGGCCAGGCT	GGTCTTGAAC	38050
TCCTGACCTC	AAGTGATCCA	CCTGCCTTAG	CCTCCCAAAG	TGCTAGGATT	38100
ACACGAGTGA	GCCACCGTGC	CCTGCCCTTA	TTACATTCTG	ATCACACATT	38150
TCATGTTTTA	TAATTGGAAA	ACTGGTGAAA	TTATAGACAA	TGTTTTGTTC	38200
CCCTAAATTC	TCCTTGATGA	GTATATATTA	CTTACACTCT	TCTGTCTTTA	38250
AAATTTTGCA	AAATAGTATC	CTAGATAAGT	TTATGAGTGC	ACAGTCTGTA	38300
CGCTTACTCA	TATTAATGAC	CTCGGAGAGT	TAAACAACAG	TCACCTTTAA	38350
AAATATTATC	TATCATGATC	ATTATTTTTG	AGGCGGGGGT	CTCATTCTGT	38400
CTCCAGGCT	GGAGAGTAGT	GGTGCGGTCA	CAGCTCACTG	CAGCCACCGC	38450
TACCTGGGCT	CAAGTGATCC	TTCTCTCTCA	GCCTTCTGAG	TAGCTGAGAC	38500
CACAGGCTTA	TGCTACCACA	CCTGGCTAAT	TTTTTAACTT	TTTGTAGAGA	38550
CGATGTCTCA	TTATGTTGCC	CAGGCTGGTC	TCAAACCTCT	AAGCTCAAGT	38600
GATCTTCTCT	AGCTCCCAA	AGTGCTGGGA	TTACAGGCAT	GAAAACTGTC	38650
ACCCAGCCCT	AAAAATTATT	AGGGTCTGTC	ATAGTAAGAC	TTTAATAAAT	38700
ATTTAAATGA	ACATCTGGTT	TTTTTAAAAA	AAAAATAGAG	ACAAGGTCTC	38750
ACTATATTGC	CCAAGCTGGT	CTCGAACTCC	TGGACTCAGC	CAATCTGTCT	38800
GCCTTAGCCG	CCCAAAGTGC	TGGGATTACA	GGCATGACCC	ACCTCATCTG	38850
GGCTGAGTGA	ACATATTTTT	AACATAAAGG	CCGTATTTTA	TATTTATCTC	38900
ATACATTTTG	CCAGCATACC	CCATTTCCGC	CGAATCTGTT	GCTTGCTAAT	38950
TCCTTCCAGC	TTCAATTCAT	CTGAAATTTG	ACAAACATCT	TCTATTTCTT	39000
TGTCGTCAAT	TTATTGACTT	CAGAATATAA	AATAAAACAC	TATACCCAAA	39050
TTAAACCCCA	CCCTCATATG	CCAGCTGAT	GTGAAAATAA	TCAGCATACA	39100
TTAAGCTTAC	CCTTGATATA	TGTGTAGCAT	CTTTTAGATA	AATATACAGC	39150
TGATTAAGCA	ATATAGCCTG	ATGGTATAAT	ATCTTGCCCA	TGTACCTCAT	39200
CTTATCTCCA	GCAGGATTAA	TTACAGTGA	TCAGATTTAC	CTTTAACTT	39250
TGTAGCAAAA	TATCTCTCC	AAAAGCATAT	CTAAACCTTT	TGTGTGTAAT	39300
CTTGCAAGTT	TCTTAATTTT	ATGCAGAACA	GGCTCTTACC	ACTGTTAGCT	39350
GGAGATATTT	TCAAGACCTA	TTTTTGTGTT	TGGTTTCTG	ATGATGGTCA	39400
TGGCATTTCC	CCCTTCACTC	CATCTAAAAA	TTGAGGTGAT	ACAGGCTTTT	39450
AAACAAACCC	AACCTATATA	GACTGAGTAC	AACTGCAATG	CAGGCATGCT	39500
AACTCTGCT	ACAATCATGG	GCGTGCTATT	GATATGTCTT	AAGTTACAGA	39550
ACACAGGCT	GAGCGTCTCA	TTAGGTCAAA	ATGTAACCA	GTTTTTCTGC	39600
TCAGTGATGC	TTAATGAGGA	CAGGGTGTGA	GAGATTTCTT	TAAGGAAAAC	39650
AAATATATAA	TAATGCTACA	TGGAAAAATA	TCTAACATTA	GAGAATTAAG	39700
TAAATAAACT	AATATATCA	CACCATGGAA	TCTTGTGCAG	ACATTAATAA	39750
TATGTAGTGG	ATGGATGTTT	AATGGTGTGA	GAAAAAGTTA	GGATGTGCTG	39800
GGGTGGGGGG	AAGAAATCAAG	TTTTAAGAAA	ATACAGTATA	CCCATACTTA	39850
AGTAAAAAAA	AAAAAAAAGG	TATGTACAGT	CATGTGTTGC	TTAATGATGG	39900

29

GGATACATTC CGAGAAATGT GTCGATAGGT GATTTTCATCC TTGTGTGAAC 39950
 ATCATAGAGT GAACCTTACAC AAACCTAGAT GGTCTAGCCT ACTATGTATC 40000
 TAGGCTATAT GACTAGCCTG TTGCTCCTAG GCTACAAACC TGTAAGCAT 40050
 GTTACTGTAG CGAATATACA AATACTTAAC ACAATGGCAA GCTATCATTG 40100
 TGTTAAGTAG TTGTGTATCT AAACATATCT AAAACATAGA AAACCTAATGT 40150
 GTTGTGCTAC AATGTTACAA TGACTATGAC ATTGCTAGGC AATAGGAATT 40200
 ATAATTTTAT CCTTTTATGG AACCACACTT ATATATGCGG TCCATGGTGG 40250
 ACCAAAACAT CCTTATGTGG CATATGACTG TATACATGTA CACAAAAAAT 40300
 AGATGAAAGA ATGAATATAC ATCAAAATAT TTAAAAATGGT TATAATGACT 40350
 TAGGTTACTT TTATTTATCT TAGTAATAAT AATGATGATA GATAATACTT 40400
 TTATAGTGTT TACTATATAA AAGACACTGT TATAAGTGTT CTACATACTT 40450
 TACATGTATT ACCTAAATGA TATAAATATA ACTCTGACAG TAACTAATCT 40500
 TATACGTTCT CTTTTCTTTT TTTTTTTTTT CTTTTTTTAG ACAGAATCTT 40550
 GCTCTACCAG GCTGGAGTGC AGGGTGCAAT CTCGGCTCAC TGCAACCTCC 40600
 GCCTCCCAGG TTCAAACGAT TCTCATGTCT CAGCCTCCTG AGTAGCTGGG 40650
 ACTACAGGCA CACACCACCA TGCCCGGCTA ATTTTGTAT TTTTGGGTAG 40700
 AGATGGAGTT TTGCCATGTT GGCCAGGCTG ATCTTGAAT CCTGGCCTCA 40750
 AGTGATCTGC CTGCCTCAGC CTCCCAAAGT GCTGGGATTA CAGGTGTGAA 40800
 CCACGTGTCT CGGCCTAATC TTCAAAGTTT TCAATATTTA AAGAGTGTCTA 40850
 ACTTGTGTTGA CAATATAAAA CATATTTGAG AAAAAGAGAT ATAAGCATCT 40900
 TATTTAGAAT TATGAAAATA TCAATAGACC TACAGCCGAC TAAAGCTTTT 40950
 CTTTATAAGC TCTTGCTTAT ATTGATTGCG TCCTGTGAAT ATGCATTAAT 41000
 TTGATTTAAA TAATAAGTAT GTATAAGAAA TAACACTTTT CCTTAATTTT 41050
 TAAGAAGCTT CAACAGTTTT TAATTTGAAT TCCAATAGTG AAATACATAG 41100
 AAAATATAAA ATTTTCTGTA GTTTAGCCAA ATTGTTTTTG TTTCCACCAC 41150
 GCATTCTACC AAAATTTCTT AATAACAGTA AGAAAATGAA TGCACTACCTC 41200
 CTGCAGGGAG AGGGGAGTTA GGCAGTTTAT GGGCATAGTT ACAAGTGAGA 41250
 AATTTCAATTG GCTACCATTT ACGCTAATTT CATAAAAACT GCATTCAATT 41300
 CTATATATCT ATTTTCTTTA CATAAAAAAG GTTTCAATTA TTGGOCATTA 41350
 AATAAAATAG CCACCATTC AGAAGTTGTG TCATGTTTTT CCTTTTATA 41400
 CCACCATCAT ATTGCTTATT ATATAGATTG TGTGTGTTCC ATTTTCTGTA 41450
 ATGGGCCAGA CAGTAAGTAT TTCTGGCTTT GGAGTCCATA TGGTCTCTAT 41500
 CATAACTACT CATCTCTGCC ATGTAGCTT AAAGATTATC TAGGTCAAAT 41550
 GCCTAAGTGA TATAGTGTG AAATACAAGT TATATAATAT AGGCTGCCAC 41600
 AAAAAAAAT TTATTTGGTC TAAAAAAGAT TTCTAGACTT TTGTAGCAGC 41650
 ATGGGTGGGG CATGCACAC TTGGTTAACT CGGTGTATCT TTCTCCTTTG 41700
 CAGATCTGTC CAACTCAATG GTCTAACTCT AAAGATGGTG GATGATCAAA 41750
 CCTTGCCACC TTTAATGGAA AAACCTCTCC GGCCAGGAAG TTCCTGGGC 41800
 TTGCCAGCTT TCTCATATAG TTTTTTGTG ATAAGAAATG CCAAAGTTGC 41850
 TGCTTGATC TGAAATAAAA ATATACTAGT CCTGACACTG AATTTTTCAA 41900
 GTATACTAAG AGTAAAGCAA CTCAAGTTAT AGGAAAGGAA GCAGATACCT 41950
 TGCAAGGCAA CTAGTGGGTG CTTGAGAGAC ACTGGGACAC TGTCAGTGCT 42000
 AGATTTAGCA CAGTATTTTG ATCTCGCTAG GTAGAACACT GCTAATAATA 42050
 ATAGCTAATA ATACCTTGTT CCAAATACTG CTTAGCATTT TGCAATGTTT 42100
 ACTTTTATCT AAAGTTTTGT TTGTTTTAT TATTTATTTA TTTATTTATT 42150
 TTGAGACAGA ATCTCTCTCT GTCAOCCAGG CTGGAGTGCC ATGGTGCGAT 42200
 CTTGGCTCAC TGCAACTTTA AGCAATCTCT CTGCCTCAGC TTCCTGAGTA 42250
 GCTGGGATTA TAGGCGTGTG CCACCACGCC CAGCTACTTT CTATATTTTT 42300
 TGTAGAGATG GAGTTTCCGC ATATTGGCCA AGCTGGTCTC GAACCTCTGT 42350
 CCTCGAACTC CTGTCTCAAA GTGATCCACC CGCCTCAGCC TCTCAAAGTG 42400
 CTGGGATTAC AGGTGTGAGC CACCACACCC AGCAGTGTTT TATTTTTGAG 42450
 ACAGGGTATC ATTCTGTTGC CCAGGCTTGA GTGCAGTGGT GCAATCATAG 42500
 ATCACTGCAG CCTTTTAACT CCTGGGCTCA AGTCATCCTC CTGCTTAGCC 42550
 TCCCAAGTAG CTAGGACCAC AGACACATGC CATCACACTT GGCTATTTTT 42600
 AAAAAATTTT TTGTAGAGAT GGGGTCTCGC TATGTTACCC AAACCTGGTCC 42650
 TGAACCTCTG GACTCAATTG ATCCTCCAC CTGGCCTTC CAGGTGCTGG 42700
 GATTTCTTTG GGAGTACAGC ATGGTACAGC AGGAGATCAT TTGATGTTAC 42750
 CTCTGTGCAG TGTGCTAGT CAGCGAAAGA CTATAATACC TGTGGGGACA 42800
 GCGATTAGCC ACCACAACCA GTCTTTATTT AAAGTTATTA AAAATGGCTG 42850
 GGCGCAGTGG CTCACACCTG TAATCCTAGC ACTTTGGGAG GCCGAGGCAG 42900
 ATGGATCACG TGACGTGAGG AATTGAGAC CAGCCTGGCC AACATGGTGA 42950
 AACCCCATCT CTAATAAAAA ATACAAAAAT TAGCTGGGTG TGGTCTGTGA 43000
 GTCCAGCTA CTTGGGAGGC TGGGGCAGGA GAATTACTTG AACCCAGGAG 43050
 GCAGAGGTTG CAGTGAGCCG AGATTGTGCC ACTGCACTCC AGCCTGGGTG 43100
 ACAGAGAGAG ATTCCATCTC AAAAAAACAA GTTATTAATA ATGTATATGA 43150
 ATGCTCCTAA TATGGTCAGG AAGCAAGGAA GCGAAGGATA TATTATGAGT 43200
 TTTAAGAAAGG TGCTTAGCTG TATATTTATC TTTCAAAATG TATTAGAAGA 43250
 TTTTAGAATT CTTTCTCTCA TGTCCATCT CTACAGGCAC CCATCAGAAA 43300
 AAGCATACTG CCGTTACCGT GAAACTGGTT GTAAAAGAGA AACTATCTAT 43350
 TTGCACCTTA AAAGACAGCT AGATTTTGCT GATTTCTTC TTTGGGTTTT 43400
 CTTTGTGAGC AATAATATGT GAGAGGACAG ATTGTTAGAT ATGATAGTAT 43450
 AAAAAATGGT TAATGACAA TCAGAGCGGA GGAGATTCTG TAACTTAAA 43500
 ATTACTATAA ATGAAATTGA TTTGTCAAGA GGATAAATTT TAGAAAACAC 43550
 CCAATACCTT ATAACCTGCT GTTAATGCTT GCITTTTCTC TACCTTTCTT 43600
 CCTTGTTC A GTTGGGAAGC TTTTGGCTGC AAGTAACAGA AACTCCTAAT 43650

30

TCAAATGGCT	TAAGCAATAA	GGAAATGTAT	ATTCCACAT	AACTAGACGT	43700
TCAAACAGGC	CAGGCTCCAG	CACITCAGTA	CGTCACCAGG	GATCTGGGTT	43750
CTTCCAGCT	CTCTGCTCTG	CCATCTTTAG	CGCTGGCTTC	ATTCTCAGAC	43800
TCTGGTAGCA	TGATGGCTGT	AGCTGTTTCA	TGGGCCCTT	CAAACTCAT	43850
AGCAACCAGA	GGAGAAAAT	GAGCCATTTT	TTGAGTCTCC	TTACATAGACT	43900
TGAATAACTC	TTTTTCAGAG	CTTCTCACAG	CAAACCTCTC	CTCATGTCTC	43950
CTCATGTCTT	ATTGTTTCAGA	AATGGGTAAT	GTGGCCATTT	CACCAGTCAC	44000
TGCCAACAAAC	AACGAGGTTT	CTATAATTGT	CTCTGAGTAA	CCCTTTGGAA	44050
TGGAGAGGGT	GTGGTTCAGT	CTACAACTG	AACACTGCAG	TTCTGCGCTT	44100
TTTACCAGTG	AAAAAATGTA	ATTATTTTCC	CCTCTTAAGG	ATTAATATTTC	44150
TTCAAATGTA	TGCCTGTTAT	GGATATAGTA	TCTTTAAAT	TTTTTATTTC	44200
AATAGCTTTA	GGGTACACA	CTTTTGTCTT	ACAGGGTGA	ATTGTGTAGT	44250
GGTGAAGACT	CGGCTTTTAA	TGTACTTGTC	ACCTGAGTGA	TGTACATTGT	44300
ACCCAATAGG	TAATTTTTC	TCCATTACCC	TCCTTCCGCC	CTCTTCCCTT	44350
CTGAGTCTCC	AACATCCCTT	ATACCACTGT	GTATGTTCTT	GTGTACCTAC	44400
AGCTAAGCTT	CCACTTATAA	GTGAGAACAT	GCAGTATTTG	GTTTTCATT	44450
CCTGAGTTAC	TTCCCTTAGG	ATAACAGCCC	CCAGTTCCGT	CCAAGTTGCT	44500
GCAAAATACA	TTATTTCTT	TTATGGCTGA	GTAATAGTCC	ATGGTACATA	44550
TATACCACAT	TTCTTTATC	CACITATCAG	TTGATGGACA	CTTAGGTTAA	44600
TTCCATTCAA	TTTCATTCAA	TTTAAGTATA	TTTGTAAGGA	GCTAAAGCTG	44650
AAAATTAAAT	TTTAGATCTT	TCAATACTCT	TAAATTTTAT	ATGTAAGTGG	44700
TTTTTATATT	TTACATTG	AAATAAGTA	ATTTTATAA	CCTTGATATT	44750
GTATGACTAT	TCTTTTAGTA	ATGTAAGCC	TACAGACTCC	TACATTGGA	44800
ACCACTAGTG	TGTTGTTTCA	CCCCTTGTTA	TACTATCAGG	ATCCTCGA	44848

(2) INFORMATION FOR SEQ ID NO:43:

(i) SEQUENCE CHARACTERISTICS:

(A)	LENGTH:	2396
(B)	TYPE:	nucleic acid
(C)	STRANDEDNESS:	double
(D)	TOPOLOGY:	linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:43

TTTCTAGTTG	CTTTTAGCCA	ATGTCGGATC	AGGTTTTTCA	AGCGACAAAG	50
AGATACTGAG	ATCCTGGGCA	GAGGACATCC	TAGCTCGGTC	AGATTGGGC	100
AGGCTCAAGT	GACCACTGTC	TTAAGSCAGA	AGGGAGTCGG	GGTAGGGTCT	150
GGCTGAACCC	TCAACCGGGG	CTTTTAACTC	AGGGTCTAGT	CCTGGCGCCA	200
AATGGATGGG	ACCTAGAAAA	GGTGACAGAG	TGCGCAGGAC	ACCAGGAAGC	250
TGGTCCCACC	CCTGCGCGGC	TCCCGGGCGC	TCCTTCCCA	GGCTCCGAG	300
GATCTTGGAT	TCTGGCCACC	TCCGCACCCT	TTGGATGGGT	GTGGATGATT	350
TCAAAAGTGG	ACGTGACCGC	GGCGGAGGGG	AAAGCCAGCA	CGGAAATGAA	400
AGAGAGCGAG	GAGGGGAGGG	CGGGGAGGGG	AGGGCGCTAG	GGAGGGACTC	450
CCGGGAGGGG	TGGGAGGGAT	GGAGCGCTGT	GGGAGGGTAC	TGAGTCTTGG	500
CGCCAGAGGC	GAAGCAGGAC	CGGTTCAGG	GGCTTGAGC	CAGCGCGCCG	550
GCTGCCCCAG	CTCTCCCGGC	AGCGGGCGGT	CCAGCCAGGT	GGGATGCTGA	600
GGCTGCTGCT	GCTGTGGCTC	TGGGGGCGGC	TCGGTGCCCT	GGCCCAGGGC	650
GCCCCGCGG	GGACCGCGCC	GACCGACGAC	GTGGTAGACT	TGGAGTTTTA	700
CACCAAGCGG	CGCTCCGAA	GCCTGAGTCC	CTCGTTCCTG	TCCATCACCA	750
TGACGCCAG	CCTGGCCACC	GACCCGCGCT	TCCTCACCTT	CCTGGGCTCT	800
CCAAGGCTCC	GTGCTCTGGC	TAGAGGCTTA	TCTCCTGCAT	ACTTGAGATT	850
TGGCGGCACA	AAGACTGACT	TCCTTATTTT	TGATCCGGAC	AAGGAACCGA	900
CTTCCGAAGA	AAGAAGTTAC	TGGAAATCTC	AAGTCAACCA	TGATATTTGC	950
AGGTCTGAGC	CGGTCTCTGC	TGCGGTGTTG	AGGAACTCC	AGGTGGAATG	1000
GCCCTTCCAG	GAGCTGTTGC	TGCTCCGAGA	GCAGTACCAA	AAGGAGTTCA	1050
AGAACAGCAC	CTACTCAAGA	AGCTCAGTGG	ACATGCTCTA	CAGTTTGGCC	1100
AAGTGCTCGG	GGTTAGACCT	GATCTTTGGT	CTAAATGCGT	TACTACGAAC	1150
CCCAGACTTA	CGGTGGAACA	GCTCCAACGC	CCAGCTTCTC	CTTGACTACT	1200
GCTCTTCCAA	GGGTATATAAC	ATCTCCTGGG	AACTGGGCAA	TGAGCCCAAC	1250
AGTTTCTGGA	AGAAAGCTCA	CATTCTCATC	GATGGGTGTC	AGTTAGGAGA	1300
AGACTTTGTG	GAGTTGCATA	AACCTCTACA	AAGGTCAGCT	TTCCAAAATG	1350
CAAAACTCTA	TGGTCTTGAC	ATCGGTCAGC	CTCGAGGGAA	GACAGTTAAA	1400
CTGCTGAGGA	GTTTCTTGAA	GGCTGGCGGA	GAAGTGATCG	ACTCTCTTAC	1450
ATGGCATCAC	TATTACTTGA	ATGGACGCAT	CGCTACCAA	GAAGATTTTC	1500

31

TGAGCTCTGA TGCCTGGAC ACTTTTATTC TCTCTGTGCA AAAAATCTG 1550
 AAGGTCACTA AAGAGATCAC ACCTGGCAAG AAGGTCTGGT TGGGAGAGAC 1600
 GAGCTCAGCT TACGGTGGCG GTGCACCCTT GCTGTCCAAC ACCTTTGCAG 1650
 CTGGCTTTAT GTGGCTGGAT AAATTGGGCC TGTCAGCCCA GATGGGCATA 1700
 GAAGTCGTGA TGAGGCAGGT GTTCTTCGGA GCAGGCAACT ACCACTTAGT 1750
 GGATGAAAAAC TTTGAGCCCTT TACCTGATTA CTGGCTCTCT CTTCTGTTC 1800
 AGAAACTGGT AGGTCCAGG GTGTTACTGT CAAGAGTGAA AGGCCAGAC 1850
 AGGAGCAAAC TCCGAGTGA TCTCCACTGC ACTAACGTCT ATCACCACG 1900
 ATATCAGGAA GGAGATCTAA CTCTGTATGT CCTGAACCTC CATAATGTCA 1950
 CCAAGCACTT GAAGGTACCG CCTCCGTTGT TCAGGAAACC AGTGGATACG 2000
 TACCTTCTGA AGCCTTCGGG GCCGGATGGA TTACTTTCCA AATCTGTCCA 2050
 ACTGAACGGT CAAATTCTGA AGATGGTGGA TGAGCAGACC CTGCCAGCTT 2100
 TGACAGAAAA ACCTCTCCCC GCAGGAAGTG CACTAAGCCT GCCTGCCTTT 2150
 TCCTATGGTT TTTTGTGCAT AAGAAATGCC AAAATCGCTG CTTGTATATG 2200
 AAAATAAAAG GCATACGGTA CCCCTGAGAC AAAAGCCGAG GGGGGTGTTA 2250
 TTCATAAAAC AAAACCTAG TTTAGGAGGC CACCTCCTTG CCGAGTTCCA 2300
 GAGCTTCGGG AGGGTGGGGT ACACTTCAGT ATTACATTCA GTGTGGTGTT 2350
 CTCTCTAAGA AGAATACTGC AGGTGGTGAC AGTTAATAGC ACTGTG 2396

(2) INFORMATION FOR SEQ ID NO:44:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 535
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:44

Met Leu Arg Leu Leu Leu Leu Trp Leu Trp Gly Pro Leu Gly Ala
 5 10 15
 Leu Ala Gln Gly Ala Pro Ala Gly Thr Ala Pro Thr Asp Asp Val
 20 25 30
 Val Asp Leu Glu Phe Tyr Thr Lys Arg Pro Leu Arg Ser Val Ser
 35 40 45
 Pro Ser Phe Leu Ser Ile Thr Ile Asp Ala Ser Leu Ala Thr Asp
 50 55 60
 Pro Arg Phe Leu Thr Phe Leu Gly Ser Pro Arg Leu Arg Ala Leu
 65 70 75
 Ala Arg Gly Leu Ser Pro Ala Tyr Leu Arg Phe Gly Gly Thr Lys
 80 85 90
 Thr Asp Phe Leu Ile Phe Asp Pro Asp Lys Glu Pro Thr Ser Glu
 95 100 105
 Glu Arg Ser Tyr Trp Lys Ser Gln Val Asn His Asp Ile Cys Arg
 110 115 120
 Ser Glu Pro Val Ser Ala Ala Val Leu Arg Lys Leu Gln Val Glu
 125 130 135
 Trp Pro Phe Gln Glu Leu Leu Leu Leu Arg Glu Gln Tyr Gln Lys
 140 145 150
 Glu Phe Lys Asn Ser Thr Tyr Ser Arg Ser Ser Val Asp Met Leu
 155 160 165
 Tyr Ser Phe Ala Lys Cys Ser Gly Leu Asp Leu Ile Phe Gly Leu
 170 175 180
 Asn Ala Leu Leu Arg Thr Pro Asp Leu Arg Trp Asn Ser Ser Asn
 185 190 195
 Ala Gln Leu Leu Leu Asp Tyr Cys Ser Ser Lys Gly Tyr Asn Ile
 200 205 210
 Ser Trp Glu Leu Gly Asn Glu Pro Asn Ser Phe Trp Lys Lys Ala
 215 220 225
 His Ile Leu Ile Asp Gly Leu Gln Leu Gly Glu Asp Phe Val Glu

32

230	235	240
Leu His Lys Leu Leu Gln Arg Ser Ala Phe Gln Asn Ala Lys Leu		
245	250	255
Tyr Gly Pro Asp Ile Gly Gln Pro Arg Gly Lys Thr Val Lys Leu		
260	265	270
Leu Arg Ser Phe Leu Lys Ala Gly Gly Glu Val Ile Asp Ser Leu		
275	280	285
Thr Trp His His Tyr Tyr Leu Asn Gly Arg Ile Ala Thr Lys Glu		
290	295	300
Asp Phe Leu Ser Ser Asp Ala Leu Asp Thr Phe Ile Leu Ser Val		
305	310	315
Gln Lys Ile Leu Lys Val Thr Lys Glu Ile Thr Pro Gly Lys Lys		
320	325	330
Val Trp Leu Gly Glu Thr Ser Ser Ala Tyr Gly Gly Gly Ala Pro		
335	340	345
Leu Leu Ser Asn Thr Phe Ala Ala Gly Phe Met Trp Leu Asp Lys		
350	355	360
Leu Gly Leu Ser Ala Gln Met Gly Ile Glu Val Val Met Arg Gln		
365	370	375
Val Phe Phe Gly Ala Gly Asn Tyr His Leu Val Asp Glu Asn Phe		
380	385	390
Glu Pro Leu Pro Asp Tyr Trp Leu Ser Leu Leu Phe Lys Lys Leu		
395	400	405
Val Gly Pro Arg Val Leu Leu Ser Arg Val Lys Gly Pro Asp Arg		
410	415	420
Ser Lys Leu Arg Val Tyr Leu His Cys Thr Asn Val Tyr His Pro		
425	430	435
Arg Tyr Gln Glu Gly Asp Leu Thr Leu Tyr Val Leu Asn Leu His		
440	445	450
Asn Val Thr Lys His Leu Lys Val Pro Pro Pro Leu Phe Arg Lys		
455	460	465
Pro Val Asp Thr Tyr Leu Leu Lys Pro Ser Gly Pro Asp Gly Leu		
470	475	480
Leu Ser Lys Ser Val Gln Leu Asn Gly Gln Ile Leu Lys Met Val		
485	490	495
Asp Glu Gln Thr Leu Pro Ala Leu Thr Glu Lys Pro Leu Pro Ala		
500	505	510
Gly Ser Ala Leu Ser Leu Pro Ala Phe Ser Tyr Gly Phe Phe Val		
515	520	525
Ile Arg Asn Ala Lys Ile Ala Ala Cys Ile		
530	535	

(2) INFORMATION FOR SEQ ID NO:45:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2396
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:45

	TT TCT AGT	8
TGC TTT TAG CCA ATG TCG GAT CAG GTT TTT CAA GCG ACA AAG AGA		53
TAC TGA GAT CCT GGG CAG AGG ACA TCC TAG CTC GGT CAG ATT TGG		98
GCA GGC TCA AGT GAC CAG TGT CTT AAG GCA GAA GGG ACT CGG GGT		143
AGG GTC TGG CTG AAC CCT CAA CCG GGG CTT TTA ACT CAG GGT CTA		188
GTC CTG GCG CCA AAT GGA TGG GAC CTA GAA AAG GTG ACA GAG TGC		233
GCA GGA CAC CAG GAA GCT GGT CCC ACC CCT GCG CGG CTC CCG GGC		278

33

GCT CCC TCC CCA GGC CTC CGA GGA TCT TGG ATT CTG GCC ACC TCC	323
GCA CCC TTT GGA TGG GTG TGG ATG ATT TCA AAA GTG GAC GTG ACC	368
GCG GCG GAG GGG AAA GCC AGC ACG GAA ATG AAA GAG AGC GAG GAG	413
GGG AGG GCG GGG AGG GGA GGG CGC TAG GGA GGG ACT CCC GGG AGG	458
GGT GGG AGG GAT GGA GCG CTG TGG GAG GGT ACT GAG TCC TGG CGC	503
CAG AGG CGA AGC AGG ACC GGT TGC AGG GGG CTT GAG CCA GCG CGC	548
CGG CTG CCC CAG CTC TCC CGG CAG CGG GCG GTC CAG CCA GGT GGG	593
ATG CTG AGG CTG CTG CTG TGG CTC TGG GGG CCG CTC GGT GCC	638
Met Leu Arg Leu Leu Leu Trp Leu Trp Gly Pro Leu Gly Ala	
5 10 15	
CTG GCC CAG GGC GCC CCC GCG GGG ACC GCG CCG ACC GAC GAC GTG	683
Leu Ala Gln Gly Ala Pro Ala Gly Thr Ala Pro Thr Asp Asp Val	
20 25 30	
GTA GAC TTG GAG TTT TAC ACC AAG CGG CCG CTC CGA AGC GTG AGT	728
Val Asp Leu Glu Phe Tyr Thr Lys Arg Pro Leu Arg Ser Val Ser	
35 40 45	
CCC TCG TTC CTG TCC ATC ACC ATC GAC GCC AGC CTG GCC ACC GAC	773
Pro Ser Phe Leu Ser Ile Thr Ile Asp Ala Ser Leu Ala Thr Asp	
50 55 60	
CCG CGC TTC CTC ACC TTC CTG GGC TCT CCA AGG CTC CGT GCT CTG	818
Pro Arg Phe Leu Thr Phe Leu Gly Ser Pro Arg Leu Arg Ala Leu	
65 70 75	
GCT AGA GGC TTA TCT CCT GCA TAC TTG AGA TTT GGC GGC ACA AAG	863
Ala Arg Gly Leu Ser Pro Ala Tyr Leu Arg Phe Gly Gly Thr Lys	
80 85 90	
ACT GAC TTC CTT ATT TTT GAT CCG GAC AAG GAA CCG ACT TCC GAA	908
Thr Asp Phe Leu Ile Phe Asp Pro Asp Lys Glu Pro Thr Ser Glu	
95 100 105	
GAA AGA AGT TAC TGG AAA TCT CAA GTC AAC CAT GAT ATT TGC AGG	953
Glu Arg Ser Tyr Trp Lys Ser Gln Val Asn His Asp Ile Cys Arg	
110 115 120	
TCT GAG CCG GTC TCT GCT GCG GTG TTG AGG AAA CTC CAG GTG GAA	998
Ser Glu Pro Val Ser Ala Ala Val Leu Arg Lys Leu Gln Val Glu	
125 130 135	
TGG CCC TTC CAG GAG CTG TTG CTG CTC CGA GAG CAG TAC CAA AAG	1043
Trp Pro Phe Gln Glu Leu Leu Leu Leu Arg Glu Gln Tyr Gln Lys	
140 145 150	
GAG TTC AAG AAC AGC ACC TAC TCA AGA AGC TCA GTG GAC ATG CTC	1088
Glu Phe Lys Asn Ser Thr Tyr Ser Arg Ser Ser Val Asp Met Leu	
155 160 165	
TAC AGT TTT GCC AAG TGC TCG GGG TTA GAC CTG ATC TTT GGT CTA	1133
Tyr Ser Phe Ala Lys Cys Ser Gly Leu Asp Leu Ile Phe Gly Leu	
170 175 180	
AAT GCG TTA CTA CGA ACC CCA GAC TTA CGG TGG AAC AGC TCC AAC	1178
Asn Ala Leu Leu Arg Thr Pro Asp Leu Arg Trp Asn Ser Ser Asn	

34

185	190	195	
GCC CAG CTT CTC CTT GAC TAC TGC TCT TCC AAG GGT TAT AAC ATC			1223
Ala Gln Leu Leu Leu Asp Tyr Cys Ser Ser Lys Gly Tyr Asn Ile			
200	205	210	
TCC TGG GAA CTG GGC AAT GAG CCC AAC AGT TTC TGG AAG AAA GCT			1268
Ser Trp Glu Leu Gly Asn Glu Pro Asn Ser Phe Trp Lys Lys Ala			
215	220	225	
CAC ATT CTC ATC GAT GGG TTG CAG TTA GGA GAA GAC TTT GTG GAG			1313
His Ile Leu Ile Asp Gly Leu Gln Leu Gly Glu Asp Phe Val Glu			
230	235	240	
TTG CAT AAA CTT CTA CAA AGG TCA GCT TTC CAA AAT GCA AAA CTC			1358
Leu His Lys Leu Leu Gln Arg Ser Ala Phe Gln Asn Ala Lys Leu			
245	250	255	
TAT GGT CCT GAC ATC GGT CAG CCT CGA GGG AAG ACA GTT AAA CTG			1403
Tyr Gly Pro Asp Ile Gly Gln Pro Arg Gly Lys Thr Val Lys Leu			
260	265	270	
CTG AGG AGT TTC CTG AAG GCT GGC GGA GAA GTG ATC GAC TCT CTT			1448
Leu Arg Ser Phe Leu Lys Ala Gly Gly Glu Val Ile Asp Ser Leu			
275	280	285	
ACA TGG CAT CAC TAT TAC TTG AAT GGA CGC ATC GCT ACC AAA GAA			1493
Thr Trp His His Tyr Tyr Leu Asn Gly Arg Ile Ala Thr Lys Glu			
290	295	300	
GAT TTT CTG AGC TCT GAT GCG CTG GAC ACT TTT ATT CTC TCT GTG			1538
Asp Phe Leu Ser Ser Asp Ala Leu Asp Thr Phe Ile Leu Ser Val			
305	310	315	
CAA AAA ATT CTG AAG GTC ACT AAA GAG ATC ACA CCT GGC AAG AAG			1583
Gln Lys Ile Leu Lys Val Thr Lys Glu Ile Thr Pro Gly Lys Lys			
320	325	330	
GTC TGG TTG GGA GAG ACG AGC TCA GCT TAC GGT GGC GGT GCA CCC			1628
Val Trp Leu Gly Glu Thr Ser Ser Ala Tyr Gly Gly Gly Ala Pro			
335	340	345	
TTG CTG TCC AAC ACC TTT GCA GCT GGC TTT ATG TGG CTG GAT AAA			1673
Leu Leu Ser Asn Thr Phe Ala Ala Gly Phe Met Trp Leu Asp Lys			
350	355	360	
TTG GGC CTG TCA GCC CAG ATG GGC ATA GAA GTC GTG ATG AGG CAG			1718
Leu Gly Leu Ser Ala Gln Met Gly Ile Glu Val Val Met Arg Gln			
365	370	375	
GTG TTC TTC GGA GCA GGC AAC TAC CAC TTA GTG GAT GAA AAC TTT			1763
Val Phe Phe Gly Ala Gly Asn Tyr His Leu Val Asp Glu Asn Phe			
380	385	390	
GAG CCT TTA CCT GAT TAC TGG CTC TCT CTT CTG TTC AAG AAA CTG			1808
Glu Pro Leu Pro Asp Tyr Trp Leu Ser Leu Leu Phe Lys Lys Leu			
395	400	405	

GTA GGT CCC AGG GTG TTA CTG TCA AGA GTG AAA GGC CCA GAC AGG	1853
Val Gly Pro Arg Val Leu Leu Ser Arg Val Lys Gly Pro Asp Arg	
410 415 420	
AGC AAA CTC CGA GTG TAT CTC CAC TGC ACT AAC GTC TAT CAC CCA	1898
Ser Lys Leu Arg Val Tyr Leu His Cys Thr Asn Val Tyr His Pro	
425 430 435	
CGA TAT CAG GAA GGA GAT CTA ACT CTG TAT GTC CTG AAC CTC CAT	1943
Arg Tyr Gln Glu Gly Asp Leu Thr Leu Tyr Val Leu Asn Leu His	
440 445 450	
AAT GTC ACC AAG CAC TTG AAG GTA CCG CCT CCG TTG TTC AGG AAA	1988
Asn Val Thr Lys His Leu Lys Val Pro Pro Pro Leu Phe Arg Lys	
455 460 465	
CCA GTG GAT ACG TAC CTT CTG AAG CCT TCG GGG CCG GAT GGA TTA	2033
Pro Val Asp Thr Tyr Leu Leu Lys Pro Ser Gly Pro Asp Gly Leu	
470 475 480	
CTT TCC AAA TCT GTC CAA CTG AAC GGT CAA ATT CTG AAG ATG GTG	2078
Leu Ser Lys Ser Val Gln Leu Asn Gly Gln Ile Leu Lys Met Val	
485 490 495	
GAT GAG CAG ACC CTG CCA GCT TTG ACA GAA AAA CCT CTC CCC GCA	2123
Asp Glu Gln Thr Leu Pro Ala Leu Thr Glu Lys Pro Leu Pro Ala	
500 505 510	
GGA AGT GCA CTA AGC CTG CCT GCC TTT TCC TAT GGT TTT TTT GTC	2168
Gly Ser Ala Leu Ser Leu Pro Ala Phe Ser Tyr Gly Phe Phe Val	
515 520 525	
ATA AGA AAT GCC AAA ATC GCT GCT TGT ATA TGA AAA TAA AAG GCA	2213
Ile Arg Asn Ala Lys Ile Ala Ala Cys Ile	
530 535	
TAC GGT ACC CCT GAG ACA AAA GCC GAG GGG GGT GTT ATT CAT AAA	2258
ACA AAA CCC TAG TTT AGG AGG CCA CCT CCT TGC CGA GTT CCA GAG	2303
CTT CGG GAG GGT GGG GTA CAC TTC AGT ATT ACA TTC AGT GTG GTG	2348
TTC TCT CTA AGA AGA ATA CTG CAG GTG GTG ACA GTT AAT AGC ACT	2393
GTG	2396

(2) INFORMATION FOR SEQ ID NO:46:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 385
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:46

CGGCCGCTGC TGCTGCTGTG GCTCTGGGGG CGGCTCCGTG CCCTGACCCA	50
AGGCACTCCG GCGGGGACCG CGCCGACCAA AGACGTGGTG GACTTGGAGT	100
TTTACACCAA GAGGCTATTC CAAAGCGTGA GTCCCTCGTT CCTGTCCATC	150
ACCATCGAGC CCAGTCTGGC CACCGACCCT CGGTTCTCTA CCTTCTTGAG	200
CTCTCCACGG CTTCGAGCCC TGTCTAGAGG CTTATCTCCT GCGTACTTGA	250
GATTTGCGG CACCAAGACT GACTTCCTTA TTTTGTATCC CAACAACGAA	300
CCCACCTCTG AAGAAAGAAG TTACTGGCAA TCTCAAGACA ACAATGATAT	350

36

TTGCGGGTCT GACCGGGTCT CCGCTGACGT GTTGA

385

(2) INFORMATION FOR SEQ ID NO:47:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 541
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:47

AAATCAGGAC ATATCCTTCA CTTATTTGCC TCTTGGTCAT ATTGGAGGCA 50
TTTGTATTCA TTTTAAATAA CCTCAAAT AGTGCATGCA AAGTGCTAAG 100
CGTCATTGCG CACATGGTGC CATTAACTGT CACCACCTGC AGTGGTCTAC 150
TTAGAGAACA CCGCACTGGA TGTTAACTT GAAGCGCGTG CCCC GCCCTC 200
CCGAGGCTCT GGATCCAGCG TTGAAGCTTG CCCC GCCCTC CCGAGGCTCT 250
GGATCCAGCA CTGGAGCATG CCCC GCCCTC CCGAGGCTCT GGAGCTTGCT 300
AAGGAGTCCG CTCCCTACCG CTGGGGTTTT GCTTTATTCT TATGAATGAC 350
ACCCCTGACC GCTTTCGTCT CAGGGGTACT GTAATGCCTT TTATTTTCAT 400
ATACAAGCTG CGATTTTGGC ATTTCTTATG ACAAAAAACC CATAGGAAAA 450
GGCGGGCAGC CTTAGTGAGC TTCCTGCGGG GAGAGGTTTT TCTGTTAGAG 500
CTGGCANGGT CTGCTCATCG ACCATCTTCA GGCCTCGTGC C 541

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/03542

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) :C12N 15/56, 15/63, 1/21, 9/24, 15/11

US CL :536/23.1, 23.2; 435/200, 325, 252.3, 320.1; 424/94.61

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 536/23.1, 23.2; 435/200, 325, 252.3, 320.1; 424/94.61

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EAST, MEDLINE, BIOSIS, CAPLUS, SCISEARCH, EMBASE, JAPIO, PATOWEP, PATOSWO search terms: heparanase, gene or sequence

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X - Y	US 5,362,641 A (FUKE et al.) 08 November 1994, see entire document.	21-25 ----- 1-20, 26-28
X, P	US 5,968,822 A (PECKER et al.) 19 October 1999, see entire document.	1-28
X - Y	WO 95/04158 A1 (THE UPJOHN COMPANY) 09 February 1995, see entire document.	21-25 ----- 1-24, 26-28
X, P	WO 99/11798 A1 (INSIGHT STRATEGY & MARKETING LTD.) 11 March 1999, see entire document.	1-28

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* "A" "B" "L" "O" "P"	Special categories of cited documents: document defining the general state of the art which is not considered to be of particular relevance earlier document published on or after the international filing date document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) document referring to an oral disclosure, use, exhibition or other means document published prior to the international filing date but later than the priority date claimed	"T" "X" "Y" "A"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art document member of the same patent family
--------------------------------------	---	--------------------------	--

Date of the actual completion of the international search

12 JUNE 2000

Date of mailing of the international search report

24 JUL 2000

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

Richard Hutson
RICHARD HUTSON

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/03542

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☒ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/03542

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I, claim(s) 1-7, 19, 20 and 28, drawn to a nucleic acid encoding a polypeptide having heparanase activity.

Group II, claim(s) 8-18, drawn to antisense oligonucleotides of a polynucleotide which encodes a polypeptide having heparanase activity.

Group III, claim(s) 21-25, drawn to polypeptide having heparanase activity.

Group IV, claim(s) 26, drawn to a method of identifying a chromosome region harboring a heparanase gene.

Group V, claim(s) 27, drawn to a method of eliciting anti-heparanase antibodies in vivo.

The species listed above do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, the species lack the same or corresponding special technical features for the following reasons: The listed inventions share a technical relationship of a polypeptide having heparanase activity, but this does not constitute a special technical feature because Fuks et al. (Fuks et al. US Patent No: 5,362,641) teach a polypeptide having heparanase activity.